



Pfizer Healthcare Ireland
9 Riverwalk, National Digital Park,
Citywest Business Campus, Dublin 24. D24 NT20.
Telephone: 01-4676500 Facsimile: 01-4676501
Freephone: 1800-460-900
www.pfizer.ie

PLEASE READ

**IMPORTANT MEDICINE
SAFETY INFORMATION**

APPROVED
BY THE



Medroxyprogesterone acetate: Risk of meningioma and measures to minimise this risk

10 October 2024

Dear Healthcare Professional,

Pfizer in agreement with the European Medicines Agency and the HPRA would like to inform you of the following:

Summary

- There is an increased risk of developing meningioma with high doses of medroxyprogesterone acetate (all injectable and ≥ 100 mg oral formulations), primarily after prolonged use (several years).
- For contraception or non-oncological indications:
 - Medicines containing high doses of medroxyprogesterone acetate are contraindicated in patients with meningioma or a history of meningioma.
 - If meningioma is diagnosed in a patient treated with high doses of medroxyprogesterone acetate, treatment must be stopped.
- For oncological indications:
 - If a meningioma is diagnosed in a patient treated with high doses of medroxyprogesterone acetate, the need to continue the treatment should be carefully reconsidered, on a case-by-case basis taking into account individual benefits and risks.
- Patients treated with high doses of medroxyprogesterone acetate should be monitored for signs and symptoms of meningioma in accordance with clinical practice.

Background on the safety concern

Medroxyprogesterone acetate is available in both injectable and oral formulations for gynaecological (including contraception and endometriosis) and oncological indications. A table attached to this letter shows the formulations and indications available in the European Union/EEA.

¹ Roland N, Neumann A, Hoisnard L, Duranteau L, Froelich S, Zureik M et al. Use of progestogens and the risk of intracranial meningioma: national case-control study BMJ 2024; 384 :e078078 doi:10.1136/bmj-2023-078078.

In Ireland, medroxyprogesterone acetate as Depo-Provera 150 mg/ml Suspension for Injection is indicated for contraception.

Meningioma is a rare, most frequently benign tumour that forms from the meninges. Clinical signs and symptoms of meningioma may be non-specific and include changes in vision, hearing loss or ringing in the ears, loss of smell, headaches that worsen with time, memory loss, seizures or weakness in the extremities. While meningiomas are usually benign, their location may lead to serious consequences and may require surgery.

Based on results from a French epidemiological case-control study¹, an association between medroxyprogesterone acetate and meningioma has been observed. This study was based on data from the French National health data system (SNDS – Système National des Données de Santé) and included a population of 18,061 women who had intracranial surgery for meningioma. Each case was matched to five controls per year of birth and area of residence (90,305 controls). The exposure to medroxyprogesterone acetate 150 mg/3ml injectable was compared between women who had intracranial surgery for meningioma and women without meningioma. Analyses showed an excess risk of meningioma with the use of medroxyprogesterone acetate 150 mg/3 ml (9/18,061 cases (0.05%) vs. 11/90,305 controls (0.01%), odds ratio (OR) 5.55 (95% CI 2.27 to 13.56)). This excess risk seems to be driven by prolonged use (≥ 3 years) of medroxyprogesterone acetate 150 mg/3 ml. Although the relative risk of meningioma is significantly increased with the use of high dose medroxyprogesterone acetate, the absolute risks are very small.

No new safety concern regarding a risk of meningioma associated with the use of low dose (<100 mg) medroxyprogesterone and combination products containing medroxyprogesterone has been identified at this moment and therefore the recommendations do not apply for lower doses of oral formulations of MPA.

The product information for all relevant medroxyprogesterone acetate containing medicines will be updated accordingly and meningioma will be added as an adverse reaction with a frequency 'not known'.

Call for reporting

Healthcare professionals are asked to report any suspected adverse reactions in patients taking medicines containing medroxyprogesterone acetate via HPRA Pharmacovigilance, website: www.hpra.ie

Company contact point

For further information contact Medical Information at Pfizer Healthcare Ireland, The Watermarque Building, Ringsend Road, Dublin 4, D04 K7N3, Ireland. Telephone 1800 633 363. Email: medical.information@pfizer.com

Yours sincerely,



Orlaith Gavan

Country Medical Director

PFIZER CONFIDENTIAL

A table of presentations and pharmaceutical forms of MPA licensed in EEA is attached to this Annex 1.

Formulation	Route of Administration and Strengths	Indication
DMPA injectable suspension	IM 50mg/ml; Injection, suspension (150mg/3ml; Injection, suspension) 150mg/ml; Injection, suspension 500mg/3.3ml; Injection, suspension	Contraception Endometriosis Menopausal vasomotor symptoms Recurrent and/or metastatic (breast/endometrial/renal) cancer
	SC 104mg/0.65ml; Injection, suspension	Contraception
MPA tablets	Oral 100mg; tablet 200mg; tablet 250mg; tablet 400mg; tablet 500mg; tablet	Endometriosis Menopausal vasomotor symptoms Diagnosis of primary amenorrhea Diagnosis and treatment of secondary amenorrhea Dysfunctional (anovulatory) uterine bleeding Opposition of endometrial effects of estrogen in menopausal women being treated with estrogen (HT) Recurrent and/or metastatic (breast/endometrial/renal) cancer Metastatic prostate cancer Anorexia and cachexia syndrome