

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Benakor 5 mg tablets for dogs

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

Active substance:

Benazepril hydrochloride 5 mg

Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Colorcon Pigment Blend 22870 yellow (Iron oxides, E172)	0.5 mg
Microcrystalline cellulose (E460)	
Lactose anhydrous	
Silica colloidal anhydrous (E551)	
Sodium cyclamate (E952)	
Sodium starch glycolate Type A	
Magnesium stearate (E470b)	

Yellow oblong divisible tablets, with a break mark on both sides.

3. CLINICAL INFORMATION

3.1 Target species

Dogs.

3.2 Indications for use for each target species

Treatment of congestive heart failure.

3.3 Contraindications

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Do not use in cases of hypotension, hypovolaemia, hyponatraemia or acute renal failure.

Do not use in cases of cardiac output failure due to aortic or pulmonary stenosis.

Do not use during pregnancy or lactation (section 3.7).

3.4 Special warnings

None.

3.5 Special precautions for use

Special precautions for safe use in the target species:

No evidence of renal toxicity of the veterinary medicinal product has been observed in dogs during clinical trials, however, as is routine in cases of chronic kidney disease, it is recommended to monitor plasma creatinine, urea and erythrocyte counts during therapy.

The efficacy and safety of the veterinary medicinal product has not been established in dogs below 2.5 kg body weight.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash hands after use.

In case of accidental oral ingestion, seek medical advice immediately and show the label or the package leaflet to the physician.

Pregnant women should take special care to avoid accidental oral exposure because angiotensin converting enzyme (ACE) inhibitors have been found to affect the unborn child during pregnancy in humans.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Dogs:

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Vomiting ^a Fatigue Incoordination
Undetermined frequency (cannot be estimated from the available data)	Elevated creatinine ^b

^a Transient

^b In dogs with chronic kidney disease. At the start of therapy, a moderate increase in plasma creatinine concentration after administration of ACE inhibitors is compatible with the reduction of glomerular hypertension induced by these agents. This is therefore, in the absence of other symptoms, not necessarily a reason to discontinue treatment.

In double-blind clinical trials in dogs with congestive heart failure, benazepril hydrochloride was well tolerated with an incidence of adverse reactions lower than observed in placebo-treated dogs.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

Do not use during pregnancy or lactation. The safety of the veterinary medicinal product has not been established during pregnancy or lactation.

Laboratory studies in rats have shown evidence of embryotoxic effects (foetal urinary tract malformation) at maternally nontoxic doses.

Fertility:

The safety of the veterinary medicinal product has not been established in breeding animals.

3.8 Interaction with other medicinal products and other forms of interaction

In dogs with congestive heart failure, benazepril hydrochloride has been given in combination with digoxin, diuretics, pimobendan and anti-arrhythmic veterinary medicinal products without demonstrable adverse interactions.

In humans, the combination of ACE inhibitors and Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) can lead to reduced antihypertensive efficacy or impaired renal function. The combination of benazepril hydrochloride and other antihypertensive agents (e.g. calcium channel blockers, β -blockers or diuretics), anaesthetics or sedatives may lead to additive hypotensive effects. Therefore, concurrent use of NSAIDs or other medications with a hypotensive effect should be considered with care. Renal function and signs of hypotension (lethargy, weakness etc.) should be monitored closely and treated as necessary.

Interactions with potassium preserving diuretics like spironolactone, triamterene or amiloride cannot be ruled out. It is recommended to monitor plasma potassium levels when using benazepril hydrochloride in combination with a potassium sparing diuretic because of the risk of hyperkalaemia.

3.9 Administration routes and dosage

Oral use.

The veterinary medicinal product should be given orally once daily, with or without food. The duration of treatment is unlimited.

The veterinary medicinal product should be administered orally at a minimum dose of 0.25 mg (range 0.25-0.5) benazepril hydrochloride/kg body weight once daily, according to the following table:

Weight of dog (kg)	5 mg tablets	
	Standard dose	Double dose
>5-10	0.5 tablet	1 tablet
>10-20	1 tablet	2 tablets

The dose may be doubled, still administered once daily, to a minimum dose of 0.5 mg/kg (range 0.5-1.0), if judged clinically necessary and advised by the veterinary surgeon.

In case of using halved tablets: Put the remaining half of a divided tablet back in the blister pocket and store it in a dry place below 25 °C. Use the remaining tablet half for the next administration.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

Benazepril hydrochloride reduced erythrocyte counts in normal dogs when dosed at 150 mg/kg body weight once daily for 12 months, but this effect was not observed at the recommended dose during clinical trials in dogs.

Transient reversible hypotension may occur in cases of accidental overdose. Therapy should consist of intravenous infusion of warm isotonic saline.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QC09AA07

4.2 Pharmacodynamics

Benazepril hydrochloride is a pro-drug hydrolysed in vivo to its active metabolite, benazeprilat. Benazeprilat is a highly potent and selective inhibitor of ACE, thus preventing the conversion of inactive angiotensin I to active angiotensin II and thereby also reducing synthesis of aldosterone. Therefore, it blocks effects mediated by angiotensin II and aldosterone, including vasoconstriction of both arteries and veins, retention of sodium and water by the kidney and remodelling effects (including pathological cardiac hypertrophy and degenerative renal changes).

The veterinary medicinal product causes long-lasting inhibition of plasma ACE activity in dogs, with more than 95% inhibition at peak effect and significant activity (>80% in dogs) persisting 24 hours after dosing.

The veterinary medicinal product reduces the blood pressure and volume load on the heart in dogs with congestive heart failure.

4.3 Pharmacokinetics

After oral administration of benazepril hydrochloride, peak levels of benazepril are attained rapidly (T_{max} 1.1 hours in dogs) and decline quickly as the active substance is partially metabolised by liver enzymes to benazeprilat. The systemic bioavailability is incomplete (~13% in dogs) due to incomplete absorption (38% in dogs) and first pass metabolism. In dogs, peak benazeprilat concentrations (C_{max} of 384.16 ng/ml after a dose of 1.6 mg/kg benazepril hydrochloride) are achieved with a T_{max} of 1.1 hours.

Benazeprilat concentrations decline biphasically: the initial fast phase ($t_{1/2}$ =1.7 hours in dogs) represents elimination of free drug, while the terminal phase ($t_{1/2}$ =19 hours in dogs) reflects the release of benazeprilat that was bound to ACE, mainly in the tissues.

Benazepril and benazeprilat are extensively bound to plasma proteins (85-90%), and in tissues are found mainly in the liver and kidney.

There is no significant difference in the pharmacokinetics of benazeprilat when benazepril hydrochloride is administered to fed or fasted dogs. Repeated administration of benazepril hydrochloride leads to slight accumulation of benazeprilat ($R=1.47$ in dogs with 0.5 mg/kg), steady state being achieved within a few days (4 days in dogs).

Benazeprilat is excreted 54% via the biliary and 46% via the urinary route in dogs. The clearance of benazeprilat is not affected in dogs with impaired renal function and therefore no adjustment of benazepril hydrochloride dose is required in cases of renal insufficiency.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Not applicable.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale:

PVC/PE/PVDC-Aluminium blister: 1 year.

Aluminium/Aluminium blister: 2 years.

Tablet halves should be used within one day.

5.3 Special precautions for storage

Do not store above 25 °C.

Store in the original package.

Store tablet halves in the original blister in the original package.

5.4 Nature and composition of immediate packaging

PVC/PE/PVDC/Alu-foil blister

Cardboard box of 1, 2, 3, 4, 5, 6 or 7 blisters of 14 tablets.

or

Alu/Alu-foil blister

Cardboard box of 1, 2, 3, 4, 5, 6 or 7 blisters of 14 tablets.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Dechra Regulatory B.V.,

7. MARKETING AUTHORISATION NUMBER(S)

VPA22622/046/001

8. DATE OF FIRST AUTHORISATION

24 October 2008

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

04 March 2025

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).