

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Milpro 16 mg/40 mg film-coated tablets for cats

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

Active substances:

Milbemycin oxime 16 mg
Praziquantel 40 mg

Excipients:

Qualitative composition of excipients and other constituents	Quantitative composition if that information is essential for proper administration of the veterinary medicinal product
Core:	
Microcrystalline cellulose	
Croscarmellose sodium	
Magnesium stearate	
Povidone	
Silica hydrophobic colloidal	
Coat:	
Natural Poultry liver flavour	
Hypromellose	
Microcrystalline cellulose	
Macrogol stearate	
Allura Red AC (E129)	0.1 mg
Titanium Dioxide (E171)	0.5 mg

Oval shaped, red to pink, meat flavoured tablets with a score on both sides.
The tablets can be divided into halves.

3. CLINICAL INFORMATION

3.1 Target species

Cats (weighing at least 2 kg).

3.2 Indications for use for each target species

In cats: treatment of mixed infections by immature and adult cestodes (tapeworms) and adult nematodes (roundworms) of the following species:

Cestodes:

Echinococcus multilocularis,
Dipylidium caninum,

Taenia spp.,
Nematodes:
Ancylostoma tubaeforme,
Toxocara cati

The veterinary medicinal product can also be used in the prevention of heartworm disease (*Dirofilaria immitis*), if concomitant treatment against cestodes is indicated.

3.3 Contraindications

Do not use in cats weighing less than 2 kg
Do not use in cases of hypersensitivity to the active substances or to any of the excipients.

3.4 Special warnings

It is recommended to treat all the animals living in the same household concomitantly.

In order to develop an effective worm control programme local epidemiological information and the living conditions of the cat should be taken into account and therefore it is recommended to seek professional advice.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

When *Dipylidium caninum* infection is present, concomitant treatment against intermediate hosts, such as fleas and lice, should be considered to prevent re-infection.

3.5 Special precautions for use

Special precautions for safe use in the target species:

No studies have been performed with severely debilitated cats or individuals with seriously compromised kidney or liver function. The veterinary medicinal product is not recommended for such animals or only according to the benefit-risk assessment by the responsible veterinarian.

Studies have shown that treatment of dogs with a high number of circulating microfilariae can sometimes lead to the appearance of hypersensitivity reactions, such as pale mucous membranes, vomiting, trembling, laboured breathing or excessive salivation. These reactions are associated with the release of proteins from dead or dying microfilariae and are not a direct toxic effect of the veterinary medicinal product. In the absence of data on cats with microfilaraemia, its use should be according to a benefit risk assessment by the attending veterinarian.

The tablets are flavoured. In order to avoid any accidental ingestion, store tablets out of reach of the animals.

As per good veterinary practice, animals should be weighed to ensure accurate dosing. Ensure cats and kittens weighing between 0.5 kg and ≤ 2 kg receive the appropriate tablet strength (4 mg MBO/10 mg praziquantel) and the appropriate dose (1/2 or 1 tablet) for the corresponding weight band (1/2 tablet for cats weighing 0.5 to 1 kg ; 1 tablet for cats weighing >1 to 2 kg – 1 tablet).

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not handle this veterinary medicinal product in case of hypersensitivity to the active substances or to any of the excipients.

Wash hands after use.

Part tablets should be returned to the open blister pack and stored in the carton.

In case of accidental ingestion of the tablets, particularly by a child, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Not applicable.

Other precautions:

Echinococcosis represents a hazard for humans. As Echinococcosis is a notifiable disease to the World Organisation for Animal Health (WOAH), specific guidelines on the treatment and follow-up, and on the safeguard of persons, need to be obtained from the relevant competent authority.

3.6 Adverse events

Cats (weighing at least 2 kg).

Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Hypersensitivity reaction ¹ Systemic disorders ¹ (e.g. Lethargy) Neurological disorders ¹ (e.g. Ataxia, Muscle tremor) Digestive tract disorders ¹ (e.g. Emesis, Diarrhoea)
--	--

¹ Especially in young cats.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation:

In a study, this combination of active substances was demonstrated to be well tolerated in breeding queens, including during pregnancy and lactation.

As a specific study with this veterinary medicinal product has not been performed, use during pregnancy and lactation only according to the benefit-risk assessment by the responsible veterinarian.

3.8 Interaction with other medicinal products and other forms of interaction

The concurrent use of the combination praziquantel/milbemycin oxime with selamectin is well tolerated. No interactions were observed when the recommended dose of the macrocyclic lactone selamectin was administered during treatment with the combination at the recommended dose. In the absence of further studies, caution should be taken in the case of concurrent use of the veterinary medicinal product and other macrocyclic lactones. Also no such studies have been performed with reproducing animals.

3.9 Administration routes and dosage

Oral use.

To ensure a correct dosage, body weight should be determined as accurately as possible.

Minimum recommended dose rate: 2 mg of milbemycin oxime and 5 mg of praziquantel per kg are given once orally as a single dose.

The veterinary medicinal product should be administered with or after some food.

The veterinary medicinal product is a small size tablet.

To aid with administration, the veterinary medicinal product has been coated with a meat flavour.

The tablets can be divided into halves.

Depending on the bodyweight of the cat, the practical dosing is as follows:

Weight	Tablets
2 – 4 kg	1/2 tablet
>4 – 8 kg	1 tablet
>8 – 12 kg	1½ tablets

The veterinary medicinal product can be inserted into a programme for prevention of heartworm disease if at the same time treatment against tapeworms is indicated. The veterinary medicinal product has a duration of heartworm prevention of one month. For prevention of heartworm disease the use of a monosubstance is preferred.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

In a study conducted with the veterinary medicinal product administered at 1X, 3X and 5X the therapeutic dose, and for a duration which exceed the therapeutic indication, i.e. 3 times at 15 day-intervals, signs uncommonly reported at the recommended dose (see section 3.6) have been observed at 5-fold the therapeutic dose after the second and third treatments. These signs disappeared spontaneously within a day.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance

Not applicable.

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QP54AB51

4.2 Pharmacodynamics

Milbemycin oxime belongs to the group of macrocyclic lactones, isolated from the fermentation of *Streptomyces hygroscopicus* var. *aureolacrimosus*. It is active against mites, against larval and adult stages of nematodes as well as against larvae of *Dirofilaria immitis*. The activity of milbemycin is related to its action on invertebrate neurotransmission: Milbemycin oxime, like avermectins and other milbemycins, increases nematode and insect membrane permeability to chloride ions via glutamate-gated chloride ion channels (related to vertebrate GABA_A and glycine receptors). This leads to hyperpolarisation of the neuromuscular membrane and flaccid paralysis and death of the parasite.

Praziquantel is an acylated pyrazino-isoquinoline derivative. Praziquantel is active against cestodes and trematodes. It modifies the permeability for calcium (influx of Ca²⁺) in the membranes of the parasite inducing an imbalance in the membrane structures, leading to membrane depolarisation and almost instantaneous contraction of the musculature (tetany), rapid vacuolization of the syncytial tegument and subsequent tegumental disintegration (blebbing), resulting in easier expulsion from the gastrointestinal tract or death of the parasite.

4.3 Pharmacokinetics

In the cat, praziquantel reaches peak plasma concentrations within 1-4 hours after oral administration. The half life of elimination is around 3 hours.

In the dog, there is rapid hepatic biotransformation, principally to monohydroxylated derivatives. The principal route of elimination in the dog is renal.

After oral administration in the cat, milbemycin oxime reaches peak plasma concentrations within 2-4 hours. The half life of elimination is around 32 to 48 hours.

In the rat, metabolism appears to be complete although slow, since unchanged milbemycin oxime has not been found in urine or feces. Main metabolites in the rat are monohydroxylated derivatives, attributable to hepatic biotransformation. In addition to relatively high liver concentrations, there is some concentration in fat, reflecting its lipophilicity.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

Not applicable.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

Shelf life after first opening the immediate packaging (for half tablets): 6 months.

5.3 Special precautions for storage

This veterinary medicinal product does not require any special temperature storage conditions.

Half tablets should be stored in the original blister and be used for the next administration.

Keep the blister in the outer carton.

5.4 Nature and composition of immediate packaging

Aluminium/ Aluminium blister pack (Oriented polyamide/Aluminium/Polyvinyl chloride sealed to Aluminium film).

Available pack sizes:

Cardboard box of 2 tablets containing 1 blister of 2 tablets

Cardboard box of 4 tablets containing 2 blisters of 2 tablets

Cardboard box of 24 tablets containing 12 blisters of 2 tablets

Cardboard box of 48 tablets containing 24 blisters of 2 tablets

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

The veterinary medicinal product should not enter water courses as milbemycin oxime may be dangerous for fish and other aquatic organisms.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

VIRBAC

7. MARKETING AUTHORISATION NUMBER(S)

VPA10988/096/002

8. DATE OF FIRST AUTHORISATION

15/08/2014

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

10/07/2025

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

