1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Felipex 5 mg/ml Oral Solution for Cats

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One mL contains:

Active substance:
Thiamazole 5 mg

Excipients:

Qualitative composition of excipients and other constituents	**Quantitative composition if that information is essential for proper administration of the veterinary medicinal product**
Sodium Benzoate (E211)	1.5 mg
Hypromellose	-
Povidone K30	-
Glycerol	-
Disodium phosphate dihydrate	-
Sodium dihydrogen phosphate dihydrate	-
Citric acid	-
Simethicon Emulsion	-
Honey Flavour	-
Purified water	

Colourless to light yellow, opaque solution.

3. CLINICAL INFORMATION

3.1 Target species

Cats.

3.2 Indications for use for each target species

For the stabilisation of hyperthyroidism prior to surgical thyroidectomy.
For the long-term treatment of feline hyperthyroidism.

3.3 Contraindications

Do not use in cats suffering from systemic disease such as primary liver disease or diabetes mellitus.
Do not use in cats showing signs of autoimmune disease.
Do not use in animals with disorders of white blood cells, such as neutropenia and lymphopenia.
Do not use in animals with platelet disorders and coagulopathies (particularly thrombocytopenia).
Do not use in pregnant or lactating females. Please refer to section 3.7.
Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

3.4 Special warnings

In order to enhance stabilisation of the hyperthyroid patient the same feeding and dosing schedule should be used daily.

3.5 Special precautions for use

Special precautions for safe use in the target species:

- If more than 10 mg of thiamazole per day (2 ml of the veterinary medicinal product) is required, animals should be monitored particularly carefully.

- Use of the veterinary medicinal product in cats with renal dysfunction should be subject to careful benefit-risk assessment by the clinician. Due to the effect thiamazole can have on reducing the glomerular filtration rate, the effect of therapy on renal function should be monitored closely as deterioration of an underlying condition may occur.

- Haematology must be monitored due to risk of leucopenia or haemolytic anaemia.
- Any animal that suddenly appears unwell during therapy, particularly if they are febrile, should have a blood sample taken for routine haematology and biochemistry.
- Neutropenic animals (neutrophil counts $<2.5 \times 10^9$/litre) should be treated with prophylactic bactericidal antibacterial drugs and supportive therapy.

- As thiamazole can cause haemoconcentration, cats should always have access to drinking water. Please refer to section 3.9 for monitoring instructions.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- As thiamazole is a suspected human teratogen and it is excreted in breast milk, women of child-bearing age and lactating women must wear non-permeable single-use gloves when administering the veterinary medicinal product, or handling litter/vomit of treated cats.

- If you are pregnant, think you may be pregnant or are attempting to conceive, you should not administer the veterinary medicinal product or handle the litter/vomit of treated cats.

- This veterinary medicinal product can cause allergic reactions after dermal contact as well as skin and eye irritation. Avoid skin and eye contact including hand-to-eye contact. People with known hypersensitivity (allergy) to thiamazole or sodium benzoate should avoid contact with the veterinary medicinal product. In case of accidental skin and/or eye contact, rinse exposed skin and/or eyes immediately with clean running water. If irritation and/or allergic symptoms, such as a skin rash, swelling of the face, lips or eyes or difficulty in breathing, develop, seek medical advice immediately and show the package leaflet or the label to the physician.

- Thiamazole may cause gastrointestinal disturbances, vomiting, headache, fever, arthralgia (joint pain), pruritus (itching) and pancytopaenia (decrease in blood cells and platelets). Avoid dermal and oral exposure, including hand-to-mouth contact, especially by children.

- Do not eat, drink or smoke while handling the veterinary medicinal product, vomit or used litter of treated animals.

- Do not leave filled syringes unattended. Replace the cap immediately after filling the syringe. Following administration of the veterinary medicinal product, any residues remaining on the tip of the dosing syringe should be wiped clean with a tissue. The contaminated tissue should be immediately disposed of. The used syringe should be stored with the veterinary medicinal product in the original carton.

- Wash hands with soap and water after administration of the product and handling the vomit of or litter used by treated animals. Wash any spillages or spatter from skin immediately.

- In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Special precautions for the protection of the environment:

Not applicable.

3.6 Adverse events

Cat:

Uncommon (1 to 10 animals / 1,000 animals treated):	Vomiting[1], Anorexia[1], Inappetence[1], Lethargy[1] Pruritus[1,2], Excoriation[1,2] Prolonged bleeding[1,3,4], Icterus[1,4], Hepatopathy[1] Eosinophilia[1], Lymphocytosis[1], Neutropenia[1], Lymphopenia[1], Leucopenia[1,8], Agranulocytosis[1] Thrombocytopenia[1, 6, 7], Haemolytic anaemia[1]
Rare (1 to 10 animals / 10,000 animals treated):	Autoimmune disorder (serum anti-nuclear antibodies)[5,7], Anaemia[5,7]
Very rare (<1 animal / 10,000 animals treated, including isolated reports):	Lymphadenopathy[5,7]

[1] These side-effects resolve within 7-45 days after cessation of thiamazole therapy.
[2] Severe and of the head and neck.
[3] Sign of a bleeding diathesis.
[4] Associated with hepatopathy.
[5] Immunological side-effect.
[6] Occurs uncommonly as a haematological abnormality and rarely as an immunological side effect.
[7] Treatment should be stopped immediately and alternative therapy considered following a suitable period for recovery.
[8] Slight

Adverse events have been reported following long-term control of hyperthyroidism. In many cases, signs may be mild and transitory and not a reason for withdrawal of treatment. The more serious effects are mainly reversible when medication is stopped.

Following long-term treatment with thiamazole in rodents, an increased risk of neoplasia in the thyroid gland has been shown to occur, but no evidence is available in cats.

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

3.7 Use during pregnancy, lactation or lay

Pregnancy and lactation
Laboratory studies in rats and mice have shown evidence of teratogenic and embryotoxic effects of thiamazole. The safety of the veterinary medicinal product was not assessed in pregnant or lactating cats. Do not use in pregnant or lactating females.

3.8 Interaction with other medicinal products and other forms of interaction

Concurrent treatment with phenobarbital may reduce the clinical efficacy of thiamazole.

Thiamazole is known to reduce the hepatic oxidation of benzimidazole wormers and may lead to increases in their plasma concentrations when given concurrently.

Thiamazole is immunomodulatory, therefore this should be taken into account when considering vaccination programmes.

3.9 Administration routes and dosage

Oral use.
The veterinary medicinal product should be administered directly into the mouth of the cat.
Do not administer in food as efficacy of the veterinary medicinal product when administered via this route has not been established.

For the stabilisation of feline hyperthyroidism prior to surgical thyroidectomy and for the long-term treatment of feline hyperthyroidism, the recommended starting dose is 5 mg of thiamazole per day (1 ml of the veterinary medicinal product per day) .

The total daily dose should be divided into two equal doses and administered morning and evening. If, for reasons of compliance, once daily dosing is preferable, then this is acceptable, although a twice-daily dose (2.5 mg of thiamazole or 0.5 ml of the veterinary medicinal product, given twice) may be more efficacious in the short term. In order to enhance stabilisation of the hyperthyroid patient the same dosing schedule relative to feeding should be used daily.

Use the syringe provided in the package. The syringe is graduated in 0.25 mg increments up to 5 mg and fits on the bottle. In order to administer the dose accurately, remove the cap from the bottle, insert the syringe tip into the adapter of the bottle, invert the bottle to draw up the required dose, return the bottle to an upright position and remove the syringe from the bottle and administer the veterinary medicinal product directly into the cat's mouth.

Haematology, biochemistry and serum total T4 should be assessed before initiating treatment and after 3 weeks, 6 weeks, 10 weeks, 20 weeks, and thereafter every 3 months. At each of the recommended monitoring intervals, the dose should be titrated to effect according to the total T4 and to clinical response to treatment. Dose adjustments should be made in increments of 2.5 mg of thiamazole (0.5 ml of the veterinary medicinal product) and the aim should be to achieve the lowest possible dose rate. In cats that require particularly small dose adjustments, increments of 1.25 mg of thiamazole (0.25 ml of the veterinary medicinal product) can be used.

If more than 10 mg of thiamazole per day is required (2 ml of the veterinary medicinal product) animals should be monitored particularly carefully.

The dose administered should not exceed 20 mg of thiamazole per day (4 ml of the veterinary medicinal product).
For long-term treatment of hyperthyroidism, the animal should be treated for life.

3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)

In tolerance studies in young healthy cats, the following dose-related clinical signs occurred at doses of up to 30 mg/animal/day: anorexia, vomiting, lethargy, pruritus and haematological and biochemical

abnormalities such as neutropenia, lymphopenia, reduced serum potassium and phosphorus levels, increased magnesium and creatinine levels and the occurrence of anti-nuclear antibodies. At a dose of 30 mg/day some cats showed signs of haemolytic anaemia and severe clinical deterioration. Some of these signs may also occur in hyperthyroid cats treated at doses of up to 20 mg per day.

Excessive doses in hyperthyroid cats may result in signs of hypothyroidism. This is however unlikely, as hypothyroidism is usually corrected by negative feedback mechanisms. Please refer to Section 3.6: Adverse events.

If overdosage occurs, stop treatment and give symptomatic and supportive care.

3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance.

Not applicable.

3.12 Withdrawal periods

Not applicable.

4. PHARMACOLOGICAL INFORMATION

4.1 ATCvet code: QH03BB02

4.2 Pharmacodynamics

Thiamazole acts by blocking the biosynthesis of thyroid hormone in vivo. The primary action is to inhibit binding of iodide to the enzyme thyroid peroxidase, thereby preventing the catalysed iodination of thyroglobulin and T_3 and T_4 synthesis.

4.3 Pharmacokinetics

Following oral dosing in healthy cats, thiamazole is rapidly and completely absorbed with a bioavailability of >75%. However, there is a considerable variation between animals. Elimination of the drug from cat plasma is rapid with a half-life of 2.3 to 8.7 hours. Peak plasma levels occur approximately 0.5 to 1.5 hours after dosing. The mean C_{max} is 1.4 µg/ml.

In rats, thiamazole has been shown to be poorly bound to plasma protein (5%); 40% was bound to red blood cells. The metabolism of thiamazole in cats has not been investigated, however, in rats thiamazole is rapidly metabolised in the thyroid gland. About 64% of the administered dose being eliminated in the urine and only 7.8% excreted in faeces. This is in contrast with man where the liver is important for the metabolic degradation of the compound. The drug residence time in the thyroid gland is assumed to be longer than in the plasma.

From man and rats it is known that the drug can cross the placenta and concentrates in the foetal thyroid gland. There is also a high rate of transfer into breast milk.

5. PHARMACEUTICAL PARTICULARS

5.1 Major incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

5.2 Shelf life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.
Shelf life after first opening the immediate packaging: 6 months.

5.3 Special precautions for storage

Store in the original container in order to protect from light.
This veterinary medicinal product does not require any special temperature storage conditions.

5.4 Nature and composition of immediate packaging

Polyethylene terephthalate (PET) amber bottles with low density polyethylene (LDPE) cap inserts and high density polyethylene (HDPE/LDPE) or Polypropylene (PP)/HDPE child resistant caps with tamper evident seals in a cardboard box.
The veterinary medicinal product is supplied with a 1 ml measuring syringe graduated in 0.25 mg increments up to 5 mg, made of HDPE plunger and a PP barrel.

Package size:
Cardboard box with 1 bottle of 30 mL oral solution and 1 oral syringe of 1ml graduated in 0.25 mg.
Cardboard box with 1 bottle of 100 mL oral solution and 1 oral syringe of 1 mL graduated in 0.25 mg.

Not all pack sizes may be marketed.

5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

6. NAME OF THE MARKETING AUTHORISATION HOLDER

Chanelle Pharmaceuticals Manufacturing Limited.

7. MARKETING AUTHORISATION NUMBER(S)

VPA10987/186/001

8. DATE OF FIRST AUTHORISATION

17/07/2026

9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS

21/09/2026

10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the Union Product Database (https://medicines.health.europa.eu/veterinary).