

IRISH MEDICINES BOARD ACT 1995

EUROPEAN COMMUNITIES (ANIMAL REMEDIES) (No. 2) REGULATIONS 2007

(S.I. No. 786 of 2007)

VPA: **10021/045/001**
Case No: 7005869

The Irish Medicines Board in exercise of the powers conferred on it by Animal Remedies (No. 2) Regulations (S.I. No. 786 of 2007) hereby grants to:

Bayer Limited

Animal Health Division, The Atrium, Dublin 18, Ireland

an authorisation, subject to the provisions of the said Regulations and the general conditions of the attached authorisation, in respect of the Veterinary Medicinal Product:

Top Drop 40 spot-on solution for small cats

The particulars of which are set out in Part 1 and Part 2 of the said Schedule. The authorisation is also subject to any special conditions as may be specified in the said Schedule.

The authorisation,unless revoked, shall continue in force from **15/04/2009**.

Signed on behalf of the Irish Medicines Board

A person authorised in that behalf by the said Board.

(NOTE: This authorisation replaces any previous authorisation in respect of this product which is now null and void.)

Part II

Summary of Product Characteristics

1 NAME OF THE VETERINARY MEDICINAL PRODUCT

Top Drop 40 spot-on solution for small cats

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Active substance:

Imidacloprid 40 mg/pipette (0.4 ml of a 10 % solution)

Excipient(s):

Butylhydroxytoluene (E 321) 0.4mg/pipette

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Spot-on solution

Clear yellow to slightly brownish solution

4 CLINICAL PARTICULARS

4.1 Target Species

Cats

4.2 Indications for use, specifying the target species

For the prevention and treatment of flea infestations on cats of less than 4 kg body weight.

For cats of 4 kg body weight and greater use Top Drop for large cats.

Fleas on cats are killed within one day following treatment. One treatment prevents further flea infestation for three to four weeks. The product can be used as part of a treatment strategy for flea allergy dermatitis.

4.3 Contraindications

Do not treat unweaned kittens of less than 8 weeks of age.

Do not use in case of hypersensitivity to any of the ingredients contained in this product.

4.4 Special warnings for each target species

None.

4.5 Special precautions for use

Special precautions for use in animals

This product is for topical use and should not be administered orally.

Care should be taken to avoid the contents of the pipette coming into contact with the eyes or mouth of the recipient animal.

Do not allow recently treated animals to groom each other.

Avoid overdose in sick or debilitated kittens.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands thoroughly after use.

Wash off any skin contamination with soap and water.

People with known skin sensitivity may be particularly sensitive to this product.

Avoid contact of the product with the eyes or mouth.

If the product gets into eyes accidentally, the eyes should be thoroughly flushed with water. If skin or eye irritation persists, or the product is accidentally swallowed, obtain medical attention.

Do not eat, drink or smoke during application.

4.6 Adverse reactions (frequency and seriousness)

The product is bitter tasting and salivation may occasionally occur if the cat licks the application site immediately after treatment. This is not a sign of intoxication and disappears within some minutes without treatment (see also section 4.9 *Amounts to be administered and administration route*).

On very rare occasions skin reactions such as hair loss, redness, itching and skin lesions may occur. Agitation has also been reported. On exceptional occasions, excessive salivation and nervous signs such as incoordination, tremors and depression have been reported.

4.7 Use during pregnancy, lactation or lay

No primary embryotoxic, teratogenic or reproductive toxic effects have been observed during the studies with imidacloprid on rats and rabbits. Studies on pregnant and lactating queens together with their offspring are limited. Evidence so far suggests that no adverse effects are to be expected in these animals.

4.8 Interaction with other medicinal products and other forms of interaction

No incompatibility has been observed between this product at twice the recommended dose and the following commonly used veterinary products: lufenuron, pyrantel and praziquantel. The compatibility of the product was also demonstrated with a wide range of routine treatments under field conditions including vaccination.

4.9 Amounts to be administered and administration route

Dosage and Treatment Schedule

Cat (kg bw)	Product	Number of Pipettes	Imidacloprid (mg/kg bw)
< 4 kg	Top Drop for Small Cats	1 x 0.4 ml	minimum of 10
≥ 4kg	Top Drop for Large Cats	1 x 0.8 ml	minimum of 10

Re-infestation from emergence of new fleas in the environment may continue to occur for six weeks or longer after treatment is initiated. More than one treatment may therefore be required, depending on the level of fleas in the environment. To aid in environmental challenge, the additional use of a suitable environmental treatment against adult fleas and their developing stages is recommended.

The product remains effective if the animal becomes wet, for example after exposure to heavy rain. However, re-treatment may become necessary, depending on the presence of fleas in the environment. In these cases do not re-treat more frequently than once weekly.

Method of Administration

Remove one pipette from the package. Hold pipette in an upright position, twist and pull off cap. Use reversed cap to twist and remove seal from pipette.



Part the hair on the cat's neck at the base of the skull until the skin is visible. Place the tip of the pipette on the skin and squeeze firmly several times to empty the contents directly onto the skin.



The product is bitter tasting and salivation may occasionally occur if the cat licks the application site immediately after treatment. This is not a sign of intoxication and disappears within some minutes without treatment. Application at the base of the skull will minimize the opportunity for the cat to lick the product.

Apply only to undamaged skin. Do not allow recently treated animals to groom each other.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

No adverse clinical signs were produced using doses of five times the therapeutic level weekly for eight consecutive weeks.

In rare cases of overdose or licking of treated fur, nervous system disorders (such as twitching, tremors, ataxia, mydriasis, miosis, lethargy) can occur.

Poisoning following inadvertent oral uptake in animals is unlikely. In this event, treatment should be symptomatic under veterinary medical attention. There is no known specific antidote but administration of activated charcoal may be beneficial.

4.11 Withdrawal Period(s)

Not applicable.

5 PHARMACOLOGICAL or IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: antiparasitic agent

ATC vet code: QP53AX17

5.1 Pharmacodynamic properties

Imidacloprid, 1-(6-Chloro-3-pyridylmethyl)-N-nitro-imidazolidin-2-ylideneamine* is an ectoparasiticide belonging to a group of chloronicotinyl compounds. Chemically, it is more accurately described as a chloronicotinyl nitroguanidine.

The substance has a high affinity for the nicotinic acetylcholine receptors in the post-synaptic region of the central nervous system (CNS). The ensuing inhibition of cholinergic transmission in insects results in paralysis and death. Due to the weak nature of the interaction with mammalian nicotinic receptor sites and the postulated poor penetration through the blood/brain barrier in mammals, it has virtually no effect on the mammalian CNS. The minimal pharmacological activity in mammals is supported by safety studies involving systemic administration of sub-lethal doses to rabbits, mice and rats.

In further studies, in addition to the adulticide flea efficacy of imidacloprid, a larvicidal flea efficacy in the surroundings of the treated pet has been demonstrated. Larval stages in the pet's surroundings are killed following contact with a treated animal.

* CAS-No. 138261-41-3

5.2 Pharmacokinetic properties

The product is indicated for cutaneous administration. Following topical application in cats, the solution is quickly distributed over the animal. Acute dermal studies in the rat and target animal overdose and serum kinetic studies have established that systemic absorption is very low, transient and not relevant for the clinical efficacy. This has been further demonstrated by a study in which fleas were not killed after having fed on previously treated animals once the animal's skin and fur had been cleaned of all active material.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Butylhydroxytoluene E321

Benzyl alcohol

Propylene carbonate

6.2 Incompatibilities

None known.

6.3 Shelf-life

Shelf life of the veterinary medicinal product as packaged for sale
5 years

6.4 Special precautions for storage

No special storage conditions required.
Store away from food, drink and animal feeding stuffs.

6.5 Nature and composition of immediate packaging

Pack sizes	0.4 ml solution per pipette Blister pack containing 2, 3, 4, or 6 unit dose pipettes
------------	---

Container	White polypropylene pipettes with caps
-----------	--

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials

Any unused product or waste materials should be disposed of in accordance with national requirements.

7 MARKETING AUTHORISATION HOLDER

Bayer Ltd,
Animal Health Division,
The Atrium,
Blackthorn Road,
Dublin 18,
Ireland.

8 MARKETING AUTHORISATION NUMBER(S)

VPA 10021/45/1

9 DATE OF THE FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

21st November 2007

10 DATE OF REVISION OF THE TEXT

15th April 2009