

## Summary of Product Characteristics

### 1 NAME OF THE VETERINARY MEDICINAL PRODUCT

Advantage Flea Control 40 mg Spot-on solution for small cats.

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 0.4 ml pipette contains:

**Active substance:**

Imidacloprid 40 mg

**Excipient(s):**

Butylhydroxytoluene (E 321)	0.4 mg
Benzyl alcohol (E 1519)	332.8 mg

For the full list of excipients, see section 6.1.

### 3 PHARMACEUTICAL FORM

Spot-on solution.

Clear yellow to slightly brownish solution.

### 4 CLINICAL PARTICULARS

#### 4.1 Target Species

Cats.

#### 4.2 Indications for use, specifying the target species

For the prevention and treatment of flea infestations on cats of less than 4 kg body weight.

One treatment prevents further flea infestation for three to four weeks.

#### 4.3 Contraindications

Do not treat unweaned kittens of less than 8 weeks of age.

Do not use in animals that are known to be hypersensitive to the active substance or any of the excipients.

#### 4.4 Special warnings for each target species

Re-infestation from emergence of new fleas in the environment may continue to occur for six weeks or longer after treatment is initiated. More than one treatment may therefore be required, depending on the level of fleas in the environment. To aid reduction in environmental challenge, the additional use of a suitable environmental treatment against adult fleas and their developing stages is recommended. In order to reduce further the environmental challenge, it is recommended that all dogs, cats and rabbits in the household are treated with a suitable product.

If there are several pets in the household, it is advisable to treat all pets at the same time. Fleas can be detected by parting the pet's coat and examining skin for signs of flea activity or by combing coat with a fine-toothed flea comb. Frequent scratching or excessive grooming can also be signs of flea infestation. Fleas from pets often infest pet's basket, bedding and regular resting areas, such as carpets and soft furnishings which should be vacuumed regularly and treated periodically with an insecticide suitable for household use, preferably containing an I.G.R. (Insect Growth Regulator) to control flea egg development and help break the flea life cycle. Wash and change pet bedding regularly.

The product remains effective if the animal becomes wet, for example after exposure to heavy rain. However, re-treatment may become necessary, depending on the presence of fleas in the environment. In these cases do not treat more frequently than once weekly.

#### 4.5 Special precautions for use

##### Special precautions for use in animals

This product is for topical use and should not be administered orally.

Care should be taken to avoid the contents of the pipette coming into contact with the eyes or mouth of the recipient animal.

Do not allow recently treated animals to groom each other.

##### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands thoroughly after use.

Wash off any skin contamination with soap and water.

People with known skin sensitivity may be particularly sensitive to this product.

Avoid contact of the product with the eyes or mouth.

After application, do not stroke or groom pets until application site is dry.

If the product gets into eyes accidentally, the eyes should be thoroughly flushed with water. If skin or eye irritation persists, or the product is accidentally swallowed, obtain medical attention.

Do not eat, drink or smoke during application.

##### Other precautions

Keep treated pets away from materials such as leather, soft furnishings, plastics etc until site of application is completely dry as contact may damage these surfaces.

#### **4.6 Adverse reactions (frequency and seriousness)**

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reactions during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals)
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

The product is bitter tasting and salivation may occasionally occur if the animal licks the application site immediately after treatment. This is not a sign of intoxication and disappears within some minutes without treatment (see also section 4.9 *Amounts to be administered and administration route*).

In very rare occasions skin reactions such as hair loss, redness, itching and skin lesions may occur. Agitation, excessive salivation and nervous signs such as incoordination, tremors and depression have also been reported.

#### **4.7 Use during pregnancy, lactation or lay**

No primary embryotoxic, teratogenic or reproductive toxic effects have been observed during the studies with imidacloprid on rats and rabbits. Studies on pregnant and lactating queens together with their offspring are limited. Evidence so far suggests that no adverse effects are to be expected in these animals.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

No incompatibility has been observed between this product at twice the recommended dose and the following commonly used veterinary products: lufenuron, pyrantel and praziquantel. The compatibility of the product was also demonstrated with a wide range of routine treatments under field conditions including vaccination.


## 4.9 Amounts to be administered and administration route

### *Dosage and Treatment Schedule*


Product	Cat (kg bw)	Number of Pipettes	Imidacloprid (mg/kg bw)
Advantage Flea Control 40 mg Spot-on solution for small cats	< 4 kg	1 x 0.4 ml	minimum of 10
Cats weighing $\geq$ 4kg bodyweight: use Advantage Flea Control 80 mg Spot-on solution for large cats			

### *Method of Administration*

Remove one pipette from the package. Hold pipette in an upright position, twist and pull off cap. Use reversed cap to twist and remove seal from pipette.



Part the hair on the cat's neck at the base of the skull until the skin is visible. Place the tip of the pipette on the skin and squeeze firmly several times to empty the contents directly onto the skin.



The product is bitter tasting and salivation may occasionally occur if the cat licks the application site immediately after treatment. This is not a sign of intoxication and disappears within some minutes without treatment. Application at the base of the skull will minimize the opportunity for the cat to lick the product.

Apply only to undamaged skin.

## 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

No adverse clinical signs were produced using doses of five times the therapeutic level weekly for eight consecutive weeks.

In rare cases of overdose or licking of treated fur, nervous system disorders (such as twitching, tremors, ataxia, mydriasis, miosis, lethargy) can occur.

Poisoning following inadvertent oral uptake in animals is unlikely. In this event, treatment should be symptomatic under veterinary medical attention. There is no known specific antidote but administration of activated charcoal may be beneficial.

## 4.11 Withdrawal Period(s)

Not applicable.

## 5 PHARMACOLOGICAL or IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: antiparasitic products, ectoparasiticides, imidacloprid  
ATCvet code: QP53AX17

### 5.1 Pharmacodynamic properties

Imidacloprid, 1-(6-Chloro-3-pyridylmethyl)-N-nitro-imidazolidin-2-ylideneamine is an ectoparasiticide belonging to a group of chloronicotinyl compounds. Chemically, it is more accurately described as a chloronicotinyl nitroguanidine.

The substance has a high affinity for the nicotinic acetylcholine receptors in the post-synaptic region of the central nervous system (CNS). The ensuing inhibition of cholinergic transmission in insects results in paralysis and death. Due to the weak nature of the interaction with mammalian nicotinic receptor sites and the postulated poor penetration through the blood/brain barrier in mammals, it has virtually no effect on the mammalian CNS. The minimal pharmacological activity in mammals is supported by safety studies involving systemic administration of sub-lethal doses to rabbits, mice and rats.

In further studies, in addition to the adulticide flea efficacy of imidacloprid, a larvicidal flea efficacy in the surroundings of the treated pet has been demonstrated. Larval stages in the pet's surroundings are killed following contact with a treated animal.

### 5.2 Pharmacokinetic properties

The product is indicated for cutaneous administration. Following topical application in cats, the solution is quickly distributed over the animal. Acute dermal studies in the rat and target animal overdose and serum kinetic studies have established that systemic absorption is very low, transient and not relevant for the clinical efficacy. This has been further demonstrated by a study in which fleas were not killed after having fed on previously treated animals once the animal's skin and fur had been cleaned of all active material.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Butylhydroxytoluene E321  
Benzyl alcohol E1519  
Propylene carbonate

### 6.2 Incompatibilities

None known.

### 6.3 Shelf-life

Shelf life of the veterinary medicinal product as packaged for sale:  
5 years

### 6.4 Special precautions for storage

This veterinary medicinal product does not require any special temperature storage conditions.  
Keep the blister in the outer carton.

## **6.5 Nature and composition of immediate packaging**

Pack sizes     0.4 ml solution per pipette  
                  Blister pack containing 1, 2, 3, 4, or 6 unit dose pipettes

Container     White polypropylene pipettes with caps

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Bayer Limited  
Animal Health Division  
The Atrium  
Dublin 18  
Ireland

## **8 MARKETING AUTHORISATION NUMBER(S)**

VPA 10021/066/001

## **9 DATE OF THE FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

2nd May 2014

## **10 DATE OF REVISION OF THE TEXT**