

## **Package leaflet: Information for the user**

### **Paralief Max 1000mg Effervescent Tablets**

Paracetamol

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you**

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist or nurse has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after 3 days.

#### **What is in this leaflet:**

1. What Paralief Max effervescent tablets are and what they are used for
2. What you need to know before you take Paralief Max effervescent tablets
3. How to take Paralief Max effervescent tablets
4. Possible side effects
5. How to store Paralief Max effervescent tablets
6. Contents of the pack and other information

#### **1. What Paralief Max effervescent tablets are and what they are used for**

This medicine contain paracetamol, which belongs to a group of medicines called analgesics and antipyretics that relieve mild to moderate pain and fever.

It can be used to relieve headache, migraine, neuralgia, toothache, period pain, rheumatic aches and pains, sore throat and the symptoms of colds and flu.

You must talk to a doctor if you do not feel better or if you feel worse after 3 days.

#### **2. What you need to know before you take Paralief Max effervescent tablets**

##### **Do not take Paralief Max effervescent tablets**

- If you are allergic to paracetamol or any of the ingredients of this medicine (listed in section 6).

#### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Paralief Max effervescent tablets:

- If you are suffering from kidney problems.
- If you are suffering from liver problems including liver problems due to excessive alcohol consumption.
- If you have Gilbert's syndrome (mild jaundice).
- If you have haemolytic anaemia (abnormal breakdown of red blood cells).
- If you are an asthmatic and sensitive to aspirin (acetylsalicylic acid).
- If you are suffering from dehydration or chronic malnutrition.
- If you are on paracetamol containing medicines.
- If you have fever after paracetamol therapy.
- If you suffer from glucose-6-phosphate dehydrogenase deficiency (enzyme deficiency).
- If you are elderly.
- If you are taking other drugs that affect the liver (see Other medicines and Paralief Max Effervescent Tablets)

#### **Other important information**

Do not drink alcohol (e.g. wine, beer, spirits) whilst taking this medicine.

This medicine is not suitable for patients weighting less than 50 kg.

Do not use paracetamol unless prescribed by your doctor if you have an addiction to alcohol or liver damage. If this applies to you, you must not take these tablets with alcohol. If you are already taking other pain medication containing paracetamol do not use these tablets without first speaking to your doctor or a pharmacist.

Never take more Paralief Max effervescent tablets than recommended. A higher dose does not increase pain relief; instead it can cause severe liver damage. The symptoms of liver damage occur first after a few days. It is important therefore that you contact your doctor as soon as possible if you have taken more Paralief Max effervescent tablets than recommended in this leaflet.

### **Children and adolescents**

Do not use Paralief Max effervescent tablets in children and adolescents below 16 years of age and under 50 kg body weight.

### **Other medicines and Paralief Max effervescent tablets**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Tell your doctor before you take Paralief Max effervescent tablets if you are taking:

- Drugs to thin the blood (anti-coagulant drugs e.g. warfarin, other coumarins)
- Medicines used to relieve sickness (e.g. metoclopramide, domperidone)
- Medicines used to treat high cholesterol (cholestyramine)
- Probenecid (medicine used to treat high levels of uric acid in the blood stream, i.e. gout)
- Medicine to treat fever or mild pain (aspirin (acetylsalicylic acid), salicylamide)
- Drugs such as rifampicin (drug that helps to fight infection) and some antiepileptics such as carbamazepine, phenytoin, phenobarbital, primidone (drugs used to treat or cure convulsions)
- Medicines used to treat epilepsy (lamotrigine)
- Medicines to treat tuberculosis (isoniazid)
- Flucloxacillin (antibiotic), due to a serious risk of blood and fluid abnormality (high anion gap metabolic acidosis) that must have urgent treatment and which may occur particularly in case of severe renal impairment, sepsis (when bacteria and their toxins circulate in the blood leading to organ damage), malnutrition, chronic alcoholism, and if the maximum daily doses of paracetamol are used.

Paracetamol may extend the duration of action of chloramphenicol

### **Effects of Paralief Max effervescent tablets on laboratory tests**

Uric acid and blood sugar tests may be affected.

### **Paralief Max effervescent tablets with food**

There is no significant effect on absorption of paracetamol when taken with a meal.

### **Pregnancy and breast-feeding**

If necessary, Paralief Max effervescent tablets can be used during pregnancy. You should use the lowest possible dose that reduces your pain and/or your fever and use it for the shortest time possible. Contact your doctor or pharmacist if the pain and/or fever are not reduced or if you need to take the medicine more often.

Paracetamol passes into breast milk. Therapeutic doses of paracetamol can be administered during breast-feeding.

### **Driving and using machines**

Paralief Max effervescent tablets have no influence on the ability to drive and use machines.

### **Paralief Max effervescent tablets contain sodium and aspartame**

This medicine contains 533.51 mg sodium (main component of cooking/table salt) in each effervescent tablet. This is equivalent to 26.68% of the recommended maximum daily dietary intake of sodium for an adult.

This medicine contains 10 mg aspartame in each effervescent tablet. Aspartame is a source of phenylalanine. It may be harmful if you have phenylketonuria (PKU), a rare genetic disorder in which phenylalanine builds up because the body cannot remove it properly.

### **3. How to take Paralief Max effervescent tablets**

This presentation is reserved for use in adults and in adolescents aged 16 years and above.

Always take this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure.

Paralief Max effervescent tablets are for oral administration and may be taken with or without food. Place the tablets in a full glass of water. Allow it to completely dissolve. Then drink the solution straight away.

If you have doubts on the correct dose of Paralief Max effervescent tablets to use please consult your doctor.

#### Adults and adolescents (aged 16 years and above) weighing more than 50 kg

Take one Paralief Max 1000 mg effervescent tablet (1000 mg paracetamol) every 6 hours as needed, up to a maximum of 4 tablets (4000 mg paracetamol) in 24 hours.

Paracetamol 1000 mg tablets are not suitable for:

- Children and adolescents below 16 years of age
- Adults or adolescents of 16 to 18 years and weighing less than 50 kg.

If the pain persists for more than 5 days or the fever lasts for more than 3 days, or gets worse or other symptoms appear, you should stop the treatment and consult a doctor.

Do not take Paralief Max effervescent tablets for more than 10 days without consulting the doctor.

If complaints persist or worsen, you should seek medical advice. Do not exceed the stated dose.

#### Kidney problems:

In case of kidney problems your doctor may decide to reduce the dose. The daily dose should not exceed 2 g/day unless directed by your physician.

In moderate kidney problems: The usual dose is 500 mg repeated every 6 hours.

In severe kidney problems: The usual dose is 500 mg repeated every 8 hours.

#### Liver problems:

In case of problems with your liver please consult your doctor. Your doctor may decide to reduce the dose.

#### The Elderly

Normal adult dosage is usually appropriate.

You may need a lower dose if you

- are unable to move,
- have liver or kidney problems

In chronic alcoholics, dehydrated or malnourished a dose of 2000 mg per day should not be exceeded

Do not divide the 1000 mg tablet into two equal halves for the lower doses. Paracetamol tablets of lower strengths are available in the market.

**Do not give to adolescents of 16 years of age and younger.**

**CONTAINS PARACETAMOL.**

**Do not take any other paracetamol-containing products.**

**Do not exceed the stated dose.**

**Immediate medical advice should be sought in the event of overdosage, because of the risk of irreversible liver damage.**

**If you take more Paralief Max effervescent tablets than you should**

Symptoms of paracetamol overdose in the first 24 hours may include paleness, nausea (feeling sick), vomiting, lack of desire to eat and stomach pain. Talk to a doctor at once if you take too much of this medicine even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.

**If you forget to take Paralief Max effervescent tablets**

If you forget to take a dose, take it as soon as you remember, unless it is almost time for your next dose. Remember to leave at least four hours between doses. Never double-up on a dose to make up for the one you have missed.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist or nurse.

#### **4. Possible side effects**

Like all medicines this medicine can cause side effects, although not everybody gets them.

**You should stop taking Paralief Max effervescent tablets and see your doctor immediately if you experience symptoms like:**

- Swelling in the face, tongue or throat, difficulty swallowing, red and itchy swellings on the skin and difficulty in breathing.
- Serious skin reactions causing rash or skin peeling or sores or blisters.

**Other side effects includes:**

**Rare (may affect up to 1 in 1,000 people):**

- Bleeding problems or clotting disorders, decreased formation of cells, severe decrease in white blood cells which may lead to severe infections, frequent infections due to poorly functioning white blood cells or decrease in white blood cells, reduction in blood platelets, which increases the risk of bleeding or bruising abnormal breakdown of red blood cells, which may cause weakness or pale skin, decrease in blood count, reduced neutrophil count in blood.
- Allergies (excluding swelling on the face, mouth, hands).
- Depression, confusion, sensing unreal things.
- Tremor, headache.
- Abnormal vision.
- Abnormal accumulation of fluid under the skin.
- Stomach pain, diarrhoea, nausea, vomiting, bleeding.
- Abnormal liver function, liver failure, death of liver cells, jaundice.
- Dizziness, feeling of general discomfort or uneasiness, fever, drowsiness, drug interaction.
- Overdose and poisoning.

**Very rare (may affect up to in 10,000 people):**

- Damage caused to the liver.
- Immediate severe allergic reaction (hypersensitivity reaction requiring discontinuation of treatment).
- Low levels of glucose in the blood.
- Cloudy urine and kidney disorders.
- Allergic reaction of the skin.
- Accumulation of fluid in the voice box.
- Decrease in red blood cells.
- Severe kidney impairment.
- Kidney disorder.
- Blood in urine.
- Inability to urinate.
- Uneasiness.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

HPRRA Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Paralief Max effervescent tablets**

- Keep this medicine out of the sight and reach of children.
- Do not use Paralief Max effervescent tablets after expiry date which is stated on the label after EXP. The expiry date refers to last day of that month.
- This medicinal product does not require any special temperature conditions.
- Store in the original package to protect from light and moisture.
- The solution is stable up to 8 hours below 25°C after dissolving the tablet, hence solution should be consumed within 8 hours.
- Do not use the product if you notice visible signs of deterioration, like brown or black spots, bulging of tablets or discoloration.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**6. Contents of the pack and other information**

**What Paralief Max effervescent tablet contains**

The active substance is paracetamol. Each effervescent tablet contains 1000 mg of paracetamol.

The other ingredients are:

Citric acid (anhydrous) (E330), povidone, sodium bicarbonate (E500), sodium saccharin (E954), sodium carbonate (anhydrous) (E500), simeticone (E900), polysorbate 80 (E443), aspartame (E951).

**What Paralief Max effervescent tablets look like and contents of the pack**

Paralief Max 1000 mg effervescent tablets are white to off white coloured circular flat-faced, bevelled tablets plain on both sides. Diameter: 25.2 mm.

Laminate blister strips packed into cardboard cartons.

Pack size(s) for strip pack: 8, 10, 12, 16, 20, 32, 40, 50, 60, 100 units.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer**

Clonmel Healthcare Ltd, Waterford road, Clonmel, Co. Tipperary, Ireland

**Manufacturer**

Cipla Europe NV, De Keyserlei 58-60 bus 19, Antwerp, 2018, Belgium

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Ireland (IE): Paralief Max1000 mg effervescent tablets

Spain (ES): Paracetamol Cipla 1g comprimidos efervescente EFG

**The leaflet was last revised in May 2024.**