

Part II

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

PanOxyl 10% Lotion

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Benzoyl peroxide hydrous equivalent to Benzoyl Peroxide anhydrous 10.0% w/w.

For excipients, see 6.1.

3 PHARMACEUTICAL FORM

Cutaneous Emulsion

A smooth white lotion free from gritty particles.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

PanOxyl 10% Lotion is indicated for the treatment of acne vulgaris.

4.2 Posology and method of administration

Apply to the affected area once daily. Washing with soap and water prior to application greatly enhances the efficacy of the preparation.

4.3 Contraindications

Patients with a known hypersensitivity to benzoyl peroxide should not use this product.

4.4 Special warnings and precautions for use

Avoid contact with the eyes, mouth and other mucous membranes. Care should be taken when applying the product to the neck and other sensitive areas.

The product may bleach dyed fabrics.

4.5 Interaction with other medicinal products and other forms of interaction

None.

4.6 Pregnancy and lactation

There are no restrictions on the use of the product during pregnancy or lactation.

4.7 Effects on ability to drive and use machines

None.

4.8 Undesirable effects

In normal use, a mild burning sensation will probably be felt on first application and a moderate reddening and peeling of the skin will occur within a few days. During the first few weeks of treatment a sudden increase in peeling will occur in most patients; this is not harmful and will normally subside in a day or two if treatment is temporarily discontinued.

4.9 Overdose

Not applicable.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Benzoyl peroxide has a sebostatic and keratolytic activity counteracting the hyperkeratinisation and excessive sebum secretion associated with acne vulgaris. It has antibacterial activity against *Propionibacterium acnes*, the organism implicated in acne vulgaris. The use of benzoyl peroxide in the treatment of acne is well established.

5.2 Pharmacokinetic properties

Published pharmacokinetic studies have demonstrated that benzoyl peroxide penetrates the skin following topical application, undergoing rapid metabolism to benzoic acid, which is quickly eliminated, without accumulation. A lack of systemic effects following topical administration of benzoyl peroxide has been clearly shown.

5.3 Preclinical safety data

Not applicable. Benzoyl peroxide has been in wide-spread use for many years. The relevant information is given in section 4 of the Summary of Product Characteristics.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Macrogol 1000 monostearate
Stearic acid
Glycerol monostearate
Isopropyl palmitate
Propylene glycol
Zinc soap
Purified water

6.2 Incompatibilities

Not applicable.

6.3 Shelf Life

3 years.

6.4 Special precautions for storage

Store below 25° C.

6.5 Nature and contents of container

Low density polyethylene bottles of 30ml.

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Stiefel Laboratories (UK) Ltd
Holtspur Lane
Wooburn Green
High Wycombe
Buckinghamshire HP10 0AU
England

8 MARKETING AUTHORISATION NUMBER

PA 144/11/10

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 19 July 1978

Date of last renewal: 19 July 2003

10 DATE OF REVISION OF THE TEXT

August 2006