

**Predmycin-P Liquifilm® 5mg/ml + 3.5mg/ml + 10,000U/ml Eye Drops Suspension**  
(Prednisolone Acetate)  
(Neomycin)  
(Polymyxin B Sulphate)  
Eyedrops Suspension

**PATIENT INFORMATION LEAFLET**

**WHAT DOES THE MEDICINE CONTAIN?**

This medicine contains 0.5% w/v (5 mg/ml) of Prednisolone Acetate, Neomycin Sulphate equivalent to Neomycin base 0.35% w/v (3.5 mg/ml) and 10000 units/ml of Polymyxin B Sulphate, which are the three active ingredients. It also contains Polyvinyl alcohol (Liquifilm®), Thiomersal, Sodium Acetate Trihydrate, Polysorbate 80, Propylene Glycol and Purified Water.

**TYPE OF MEDICINE**

Predmycin P® is an eye drop suspension and each bottle holds 5ml in a 10ml bottle.

**WHO IS RESPONSIBLE FOR THE PRODUCT?**

It is distributed by: Cahill May Roberts, Chapelizod, Dublin 20, Ireland

It is manufactured by: Allergan Pharmaceuticals Ireland, Westport, Co. Mayo, Ireland

Product Authorisation Holder: Allergan Pharmaceuticals Ireland, Castlebar Road, Westport, Co. Mayo, Ireland

**WHAT IS THE MEDICINE USED FOR?**

Predmycin P® is used in the treatment of blepharitis and conjunctivitis including those in which the inflammation is due to micro-organisms sensitive to the anti-infective.

**IMPORTANT INFORMATION ABOUT PREDMYCIN-P**

This medicinal product contains thiomersal as a preservative and it is possible that you may experience an allergic reaction.

**Contact your doctor if you experience blurred vision or other visual disturbances.**

**WHEN SHOULD THE MEDICINE NOT BE USED?**

Predmycin P® should not be used if you are allergic to any of its ingredients.

Predmycin P® should not be used while wearing contact lenses.

**BEFORE YOU USE THIS MEDICINE**

Tell your doctor if

You are pregnant or are breast feeding.

You have any known allergies.

Please tell your doctor if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Some medicines may increase the effects of PREDMYCIN-P and your doctor may wish to monitor you carefully if you are taking these medicines (including some medicines for HIV: ritonavir, cobicistat).

**DOSAGE**


The following information represents the average dose for this ophthalmic preparation. If your dosage is different, do not change it unless your doctor tells you to.

Intraocular instillation into the conjunctival sac or topical application to the eyelids, 1 drop applied every 3-4 hours.

Keep using this medication for the full time of treatment even if the symptoms have disappeared.

**Missed Dose** - If you miss a dose of this medicine, apply it as soon as possible. However, if it is almost time for your next dose, skip the missed dose and go back to your regular schedule.

**HOW TO USE THE MEDICINE**



**THINGS TO NOTE:**

1. Try not to touch the tip of the bottle against your eye or anything else
2. Put the cap back on immediately after use

---

**WHAT SHOULD YOU DO IF YOU USE TOO MUCH?**

If you accidentally place too many drops in your eye(s), wash with water and apply your next dose at the normal time.

Prolonged use may induce increased intraocular pressure, corneal thinning and perforation and subcapsular lenticular opacities.

If the condition does not improve within a few days, or becomes worse, check with your doctor.

**WHAT SIDE EFFECTS CAN THE MEDICINE HAVE?**

Side effects include local irritation, redness of the eye, and less frequently allergic reactions. This medicinal product contains thiomersal as a preservative and it is possible that you may experience an allergic reaction. Prolonged use may cause pigmentation of the lens and Keratopathy. Blurred vision (frequency not known).

**Reporting of side effects**

If you get any side effects, talk to your doctor , pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

HPRA Pharmacovigilance  
Earlsfort Terrace  
IRL - Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.hpra.ie](http://www.hpra.ie)  
e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

**HOW SHOULD YOU STORE THE MEDICINE?**

Keep the bottle in a safe place out of the reach and sight of children.

Do not store above 25°C.

**Shake well before use.**

Protect from freezing.

Store in an upright position.

**HOW LONG SHOULD YOU KEEP THE MEDICINE?**

Do not use the medicine if the tamper proof seal is broken.

The medicine should not be used after the expiry date.

(This is printed on both the bottle label and the bottom of the carton that the bottle is packed in).

Discard 28 days after first opening. Do not use any leftover medicine for future eye problems without first checking with your doctor.