

PACKAGE LEAFLET: INFORMATION FOR THE USER		
Hyoscine Hydrobromide 600 micrograms/ml Solution for Injection		
Hyoscine Hydrobromide (Referred to as Hyoscine Injection in this leaflet)		
D06592.00000	Read all of this leaflet carefully before you are given Hyoscine Injection because it contains important information for you. - Keep this leaflet, you may need to read it again - If you have any further questions, please ask your doctor or nurse - If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.	Other medicines and Hyoscine Injection Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular: - medicines used to treat an irregular heart beat e.g. disopyramide and quinidine. - medicines use to treat respiratory conditions such as COPD (Chronic Obstructive Pulmonary Disease) and asthma e.g. tiotropium and ipratropium. - medicines used to treat depression such as tricyclic and tetracyclic antidepressants. - medicines used to treat hay fever and other allergies e.g. antihistamines. - medicines used to relieve the symptoms of nausea and vomiting e.g. metoclopramide and domperidone. - medicines used to control nausea (feeling sick) and vomiting (being sick) and to treat mental illness e.g. phenothiazines, butyrophenones and clozapine. - medicines used for Parkinson's disease and influenza e.g. amantadine. Hyoscine Injection may also reduce the effects of the following medicines: - levodopa - Used to treat Parkinson's disease - Nitrate (GTN)tablets - Used under the tongue for angina - Parasympathomimetics - Used to treat conditions suchas glaucoma, depression, asthma, heart conditions and retention of urine. Using Hyoscine Injection with alcohol You are advised not to drink alcohol whilst being treated with Hyoscine Injection, as it may cause drowsiness. Pregnancy, breast-feeding and fertility If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine Use of Hyoscine Injection during pregnancy may cause breathing problems in the newborn infant. Use during pregnancy and breast feeding is not recommended. Driving and using machines Hyoscine Injection may cause drowsiness, dizziness or blurred vision. If affected, do not drive or operate any tools or machinery after you have been given Hyoscine Injection. Hyoscine contains Sodium This medicinal product contains 3.54mg/ml of sodium per dose. This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free. 3. How Hyoscine Injection is given This medicine is an injection which will be given to you by a doctor. Your doctor will determine the dose you require. It will be given under the skin or into a muscle. Adults Treatment of nausea (feeling sick), vomiting (being sick), vertigo (feeling dizzy and light headed), loss of balance and motion sickness: 200 micrograms as a single dose. Preventing excessive production of saliva if you have been given a general anaesthetic: 200-600 micrograms given 30-60 minutes before the anaesthetic. Use in children Treatment of nausea (feeling sick), vomiting (being sick), vertigo (feeling dizzy and light headed), loss of balance and motion sickness: 6 micrograms per kg of body weight as a single dose. Preventing excessive production of saliva if you have been given a general anaesthetic: 15 micrograms per kg of body weight given 30-60 minutes before the anaesthetic. Use in Children (under 4 months) Not recommended. Continued Overleaf

If you think you have been given too much Hyoscine Injection This medicine is given to you by your doctor so it is unlikely you will receive too much. Your doctor has information on how to recognise and treat an overdose. If you experience any of the following please speak to your doctor or nurse immediately: - dilated pupils - a rapid heartbeat - rapid breathing - abnormally high temperature - restlessness - excitement - confusion - hallucinations (seeing or hearing things that aren't there)	
4. Possible side effects Like all medicines, this medicine can cause side effects, although not everybody gets them. Stop taking your medicine and see a doctor immediately if you notice any of the following side effects as you may need urgent medical treatment: - Serious allergic reaction, signs include skin rash, itching or hives, swelling of the face, lips, tongue or other parts of the body, difficulty breathing, feeling faint or dizzy. Cases including anaphylactic shock with a fatal outcome have been reported. - collapse in hot weather due to decreased sweating The most common side effects associated with hyoscine are: - drowsiness - dry mouth - dizziness - blurred vision - difficulty passing water Other side effects include: - headache - loss of consciousness - slow heart beat - excitement - psychotic disorder; this is an illness affecting mental health. You may experience delusions (believing things which are not true), hallucinations (seeing and possibly hearing or smelling things that are not really there) or lose touch with reality. - confusion/delirium - hallucinations - dilation of the pupils - sensitivity to light - angle-closure glaucoma (red, painful eye) - heart problems - racing heart, and/or palpitations - irregular heart beat - flushing - decrease in blood pressure - constipation - nausea - vomiting - difficulty in swallowing - dyshidrosis: also known as `pompholyx' or `dyshidrotic eczema', is a skin condition which small, fluid-filled blisters appear on the fingers, palms of the hand, and sometimes on the soles of the feet. The blisters can be very itchy and painful and when they begin to heal and dry up, the skin can become cracked or scaly - skin dryness - pain at the site of injection (particularly after intramuscular use) - thirst - Neuroleptic malignant syndrome: this is a serious and potentially life-threatening condition. The signs may include a high fever, fast heartbeat, unstable blood pressure, sweating and tremor. Other signs are faster breathing, muscle stiffness, agitation and reduced consciousness. The level of an enzyme called creatine phosphokinase may be raised in your blood which suggests damage to muscles.	
Reporting of side effects If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system: Ireland HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2 Tel: +353 1 6764971 Fax: +353 1 6762517 Website: www.hpra.ie e-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.	
5. How to store Hyoscine Injection Keep this medicine out of the sight and reach of children. You should not be given this medicine if it has passed the expiry date shown on the label. The expiry date refers to the last day of that month. The doctor or nurse will check this. Do not store above 25°C. Keep the ampoule in the outer carton in order to protect from light.	
6. Contents of the pack and other information What Hyoscine Injection contains The active ingredient is Hyoscine Hydrobromide each ml contains 600 micrograms of Hyoscine Hydrobromide. The other ingredients are sodium chloride, dilute hydrobromic acid and water for injection. What Hyoscine Injection looks like and contents of the pack Hyoscine Injection is a clear, colourless solution supplied in clear glass 1ml ampoules. It is supplied in cartons containing ten ampoules. Marketing Authorisation Holder: Ethypharm 194, Bureaux de la Colline, Bâtiment D 92213, Saint-Cloud Cedex, France Manufacturer: Fannin Limited Fannin House, South County Business Park, Dublin 18, D18 Y0C9, Ireland This leaflet was last revised in: December 2025 If you would like any more information, or would like the leaflet in a different format, please contact Medical Information at the above address.	
M MARTINDALE PHARMA Bampton Road, Harold Hill, Romford, RM3 8UG, UK	
D06592.00000	



DEVELOPMENT ARTWORK

Component Code: D06592

Prod: Hyoscine Hydrobromide
600mcg/ml ampoules PIL IE

Paper Size: 170 x 296mm

Version Control	Date	By
Version A:	18/12/25	LB
Version B:		
Version C:		
Version D:		
Version E:		
Version F:		
Version G:		
Version H:		
Version I:		
Version K:		
Version L:		
Version M:		
Version N:		
Version O:		

me&you
The Old Printworks, High Street, Otford,
Sevenoaks, Kent. TN14 5PQ
t: +44 (0) 1732 743 455
e: pharma@meandyou.co.uk
www.meandyou.co.uk