

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Gericarb 100mg Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 100 mg of Carbamazepine.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Tablet

White flat, bevelled edged tablets, marked “CB” breakline “100” on one side and blank on the reverse. The breakline is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

An anticonvulsant in the management of epilepsy (generalised tonic-clonic and partial seizure types).

An analgesic in the relief of pain of trigeminal neuralgia, and the lancinating component of deafferentation pain such as in glossopharyngeal neuralgia, peripheral diabetic neuropathy, tabetic lightning pain, superior laryngeal neuralgia, stump pain, phantom limb pain, post hepatic neuralgia.

Management of alcohol withdrawal symptoms.

Treatment of mania and prophylaxis of manic-depressive illness, especially in patients unresponsive to Lithium.

4.2 Posology and method of administration

Posology

Gericarb is given orally in tablet form, usually in two to four divided doses.

Epilepsy

Treatment should be initiated with a low daily dose, gradually increasing dosage regimen is used and this should be adjusted to suit the needs of the individual patient to achieve adequate control of seizures.

Monitoring of the plasma level of carbamazepine will ensure that the optimum dose is prescribed.

In the treatment of epilepsy, the dose of carbamazepine usually requires total plasma-carbamazepine concentrations of about 4 to 12 micrograms/mL (17 to 50 micromoles/litre) (see section 4.4).

If possible a sole anti-epileptic agent should be used but if polytherapy is necessary, the same incremental dosage pattern should be followed (see section 4.5).

Adults

Gericarb should be taken in a number of divided doses.

Initially: 100 to 200 mg once or twice a day, followed by a slow increase until the best response is obtained, often 800 mg to 1200 mg daily in divided doses. In some cases, 1600 mg or even 2000mg daily in divided doses may be necessary.

Elderly

Due to the potential for drug interactions, the dosage of Gericarb should be selected with caution in elderly patients.

Children and adolescents

Usual dosage is 10-20mg/kg bodyweight daily in divided doses.

Age up to 1 year: The usual total daily dose is 100 mg to 200 mg in divided doses.

1-5 years: The usual total daily dose is 200 mg to 400 mg in divided doses.

5-10 years: The usual total daily dose is 400mg to 600mg in divided doses.

10-15 years: The usual total daily dose is 600mg to 1000mg in divided doses.

>15 years of age: 800 to 1200 mg daily (same as adult dose).

Maximum recommended dose:

Up to 6 years of age: 35 mg/kg/day

6-15 years of age: 1000 mg/day

>15 years of age: 1200 mg/day.

Gericarb Tablets are not recommended for children aged 5 years and under

Trigeminal Neuralgia and other forms of Deafferentation Pain

The individual dosage requirements of carbamazepine may vary considerably, depending on the age and weight of the patient. It is recommended that the initial dose is small but in some patients a high dose early in treatment is needed. In elderly patients, a dose of 100mg twice daily is recommended.

The usual dose is 200 mg 3 to 4 times daily but may be increased gradually until a satisfactory clinical response is obtained, which in some instances necessitates 1600mg daily.

The initial dosage of 200 to 400 mg should be slowly raised daily until freedom from pain is achieved (normally at 200 mg 3 to 4 times daily). The dosage should then be gradually reduced to the lowest possible maintenance level.

Maximum recommended dose is 1200 mg/day. When pain relief has been obtained, attempts should be made to gradually discontinue therapy, until another attack occurs.

Alcohol Withdrawal

The dose should be adjusted to suit the needs of the individual patient. The usual daily dose is 600 to 800mg in divided doses but 1200-1600mg daily may be required in delirium tremens with subsequent reductions.

Treatment of mania and prophylaxis of manic depressive illness:

Initial starting dose of 100 – 200mg daily, in divided doses, increasing gradually until symptoms are controlled or a total daily dose of 1600 mg in divided doses is reached. The usual dosage range is 400 – 600 mg daily, in divided doses.

Special populations

Renal impairment / Hepatic impairment

No data are available on the pharmacokinetics of carbamazepine in patients with impaired hepatic or renal function.

Han Chinese and Thai populations

Before deciding to initiate treatment, patients of Han Chinese and Thai origin should whenever possible be screened for HLA-B*1502 as this allele strongly predicts the risk of severe carbamazepine-associated SJS (See information on genetic testings and cutaneous reactions in section 4.4).

Method of Administration

Gericarb may be taken during, after or between meals. The tablets should be taken with a little liquid, e.g. a glass of water.

4.3 Contraindications

Gericarb Tablets are contraindicated in the following:

- Patients with a history of hypersensitivity to carbamazepine or structurally related drugs e.g. tricyclic anti-depressants
- Patients with a hypersensitivity to any of the excipients (see section 6.1)
- Patients with atrioventricular conduction defects
- Patients with a history of bone marrow depression
- Patients with history of hepatic porphyrias (e.g. acute intermittent porphyria, variegate porphyria, porphyria cutanea tarda)

- Patients who are currently on monoamine oxidase inhibitor (MAOIs) therapy or within 2 weeks of such treatment (see section 4.5)
- Herbal preparations containing St. John's Wort (*Hypericum perforatum*) must not be used while taking Gericarb due to the risk of decreased plasma concentrations and reduced clinical effect of Gericarb (see section 4.5)

4.4 Special warnings and precautions for use

Gericarb should be given only under medical supervision. Gericarb should be prescribed only after a critical benefit-risk appraisal and under close monitoring in patients with a history of cardiac, hepatic, or renal damage, adverse haematological reactions to other drugs, or interrupted courses of therapy with Gericarb.

Haematological effects

Agranulocytosis and aplastic anaemia have been associated with Gericarb; however, due to the very low incidence of these diseases, meaningful risk estimates for Gericarb are difficult to obtain. The overall risk in the general untreated population has been estimated at 4.7 persons per million per year for agranulocytosis and 2.0 persons per million per year for aplastic anaemia.

Decreased platelet or white blood cell counts occur occasionally to frequently in association with the use of Gericarb. Nonetheless, complete pre-treatment blood counts, including platelets (and possibly reticulocytes and serum iron), should be obtained as a baseline and periodically thereafter.

If the white blood cell or platelet count is definitely low or decreased during treatment, the patient and the complete blood count should be closely monitored (see section 4.8). However, treatment with Gericarb should be discontinued if the patient develops leucopenia, which is severe, progressive or accompanied by clinical manifestations, e.g. fever or sore throat. Gericarb should be discontinued if any evidence of significant bone marrow depression appears.

Patients and their relatives should be made aware of early toxic signs and symptoms indicative of a potential haematological problem, as well as symptoms of dermatological or hepatic reactions. If reactions such as fever, sore throat, rash, ulcers in the mouth, easy bruising, petechial or purpuric haemorrhage appear, the patient should be advised to consult his physician immediately.

Cutaneous reactions

Serious and sometimes fatal cutaneous reactions, including toxic epidermal necrolysis (TEN; also known as Lyell's syndrome) and Stevens-Johnson syndrome (SJS), have been reported very rarely with Gericarb. Patients should be advised of the signs and symptoms and monitored closely for skin reactions. These reactions are estimated to occur in 1-6 per 10000 new users in countries with mainly Caucasian populations, but the risk in some Asian countries is estimated to be about 10 times higher. Patients with serious dermatological reactions may require hospitalization, as these conditions may be life-threatening and may be fatal. Most of the SJS/TEN cases appear in the first few months of treatment with Gericarb. The highest risk for occurrence of SJS or TEN is within first few weeks of treatment. If signs and of SJS, Lyell's syndrome/TEN (e.g. progressive skin rash often with blisters or mucosal lesions) appear, Gericarb should be withdrawn at once and alternative therapy should be considered. The best results in managing SJS and TEN come from early diagnosis and immediate discontinuation of any suspect drug. Early withdrawal is associated with a better prognosis. If the patient has developed SJS or TEN with the use of Gericarb, it must not be re-started in this patient at any time.

There is growing evidence of the role of different HLA alleles in predisposing patients to immune-mediated adverse reactions (see section 4.2).

Association with HLA-B*1502 allele

Retrospective studies in patients of Han Chinese and Thai origin found a strong correlation between the risk of developing SJS/TEN skin reactions associated with carbamazepine and the presence in these patients of the Human Leukocyte Antigen (HLA)-B*1502 allele. The frequency of HLA-B*1502 allele ranges from 2 to 12% in Han Chinese populations and is about 8% in Thai populations. Allele frequencies up to about 2% and 6% have been reported in Korea and India, respectively. Higher reporting rates of SJS (rare rather than very rare) are reported in some countries in Asia (e.g. Taiwan, Malaysia and the Philippines) in which there is a higher frequency of the HLA-B*1502 allele in the population (e.g. above 15% in the Philippines and some Malaysian populations). There are some data that suggest an increased risk of serious carbamazepine-associated TEN/SJS in other Asian populations.

Because of the prevalence of this allele in other Asian populations (e.g. above 15% in the Philippines and Malaysia), testing genetically at risk populations for the presence of HLA-B*1502 may be considered.

The frequency of the HLA-B*1502 allele is negligible in persons of European descent, several African populations, indigenous peoples of the Americas, Hispanic populations sampled and in Japanese (<1%).

The allele frequencies listed here represent the percentage of chromosomes in the specified population that carry the allele of interest, meaning that the percentage of patients who carry a copy of the allele on at least one of their two chromosomes (i.e., the “carrier frequency”) is nearly twice as high as the allele frequency. Therefore, the percentage of patients who may be at risk is nearly twice the allele frequency.

Whenever possible, screening for the presence of HLA-B*1502 allele should be carried out in patients with ancestry in genetically at-risk populations, prior to initiating treatment with carbamazepine (see section 4.2). (If testing for the presence of the HLA-B*1502 allele should be performed, high-resolution “HLA-B*1502 genotyping” is recommended. The test is positive if either one or two HLA-B*1502 alleles are detected and negative if no HLA-B*1502 alleles are detected). If these individuals test positive, carbamazepine should not be started unless there is no other therapeutic option. Tested patients who are found to be negative for HLA-B*1502 have a low risk of SJS, although the reactions may still very rarely occur.

HLA-B*1502 may be a risk factor for the development of SJS/TEN in Chinese patients taking other anti-epileptic drugs (AED) associated with SJS/TEN. Consideration should therefore be given to avoiding use of other drugs associated with SJS/TEN in HLA-B*1502 positive patients, when alternative therapies are otherwise equally acceptable. Screening is not generally recommended in patients from populations in which the prevalence of HLA-B*1502 is low. Screening is generally not recommended for any current Gericarb users, as the risk of SJS/TEN is largely confined to the first few months of therapy, regardless of HLA-B*1502 status.

The identification of subjects carrying the HLA-B*1502 allele and the avoidance of carbamazepine therapy in these subjects has been shown to decrease the incidence of carbamazepine-induced SJS/TEN.

Association with HLA-A*3101 allele

There are some data that suggest Human Leukocyte Antigen *(HLA)-A*3101 is associated with an increased risk of carbamazepine induced cutaneous adverse drug reactions including SJS, TEN, Drug rash with eosinophilia (DRESS) or less severe acute generalized exanthematous pustulosis (AGEP) and maculopapular rash (see section 4.8) in people of European descent and the Japanese.

The frequency of the HLA-A*3101 allele varies widely between ethnic populations and its frequency is about 2 to 5% in European populations and about 10% in the Japanese population.

The frequency of this allele is estimated to be less than 5% in the majority of Australian, Asian, African and North American populations with some exceptions within 5-12%. Prevalence above 15% has been estimated in some ethnic groups in South America (Argentina and Brazil), North America (US Navajo and Sioux, and Mexico Sonora Seri) and Southern India (Tamil Nadu) and between 10%-15% in other native ethnicities in these same regions and about 10% in Japanese populations.

The allele frequencies listed here represent the percentage of chromosomes in the specified population that carry the allele of interest, meaning that the percentage of patients who carry a copy of the allele on at least one of their two chromosomes (i.e., the “carrier frequency”) is nearly twice as high as the allele frequency. Therefore, the percentage of patients who may be at risk is nearly twice the allele frequency.

The presence of HLA-A*3101 allele may increase the risk for carbamazepine induced cutaneous reactions (mostly less severe) from 5.0% in general populations to 26.0% among subjects of European ancestry, whereas its absence may reduce the risk from 5.0% to 3.8%.

Testing for the presence of HLA-A*3101 allele should be considered in patients with ancestry in genetically at-risk populations (for example, patients of the Japanese and Caucasian populations, patients who belong to the indigenous populations of the Americas, Hispanic populations, people of southern India, and people of Arabic descent), prior to initiating treatment with Gericarb. (If testing for the presence of the HLA-A*3101 allele is performed, high-resolution “HLA-A*3101 genotyping” is recommended.

The test is positive if either one or two HLA- A*3101 alleles are detected and negative if no HLA- A*3101 alleles are detected).

The use of Gericarb should be avoided in patients who are found to be positive for HLA-A*3101, unless the benefits clearly outweigh the risks. Screening is generally not recommended for any current carbamazepine users, as the risk of SJS/TEN, AGEP, DRESS and maculopapular rash is largely confined to the first few months of therapy, regardless of HLA-A*3101 status.

Limitation of genetic screening

Genetic screening results must never substitute for appropriate clinical vigilance and patient management. Many Asian patients positive for HLA-B*1502 and treated with Gericarb will not develop SJS/TEN and patients negative for HLA-B*1502 of any ethnicity can still develop SJS/TEN. Similarly many patients positive for HLA-A*3101 and treated with Gericarb will not develop SJS, TEN, DRESS, AGEP or maculopapular rash and patients negative for HLA-A*3101 of any ethnicity can still develop these severe cutaneous adverse reactions. The role of other possible factors in the development of, and morbidity from these severe cutaneous adverse reactions such as AED dose, compliance, concomitant medications, co-morbidities, and the level of dermatologic monitoring have not been studied.

Other dermatologic reactions

Mild skin reactions e.g. isolated macular or maculopapular exanthemata, can also occur and are mostly transient and not hazardous, and they usually disappear within a few days or weeks, either during the continued course of treatment or following a decrease in dosage. However, since it may be difficult to differentiate the early signs of more serious skin reactions from mild transient reactions, the patient should be kept under close surveillance with consideration given to immediately withdrawing the drug should the reaction worsen with continued use.

The HLA-A*3101 allele has been found to be associated with less severe adverse cutaneous reactions from carbamazepine and may predict the risk of these reactions from carbamazepine, such as anticonvulsant hypersensitivity syndrome or non-serious rash (maculopapular eruption). However, the HLA-B*1502 allele has not been found to predict the risk of these aforementioned skin reactions.

Hypersensitivity

Gericarb may trigger hypersensitivity reactions, including Drug Rash with Eosinophilia and Systemic Symptoms (DRESS), a delayed multi-organ hypersensitivity disorder with fever, rash, vasculitis, lymphadenopathy, pseudo lymphoma, arthralgia, leukopenia, eosinophilia, hepato-splenomegaly, abnormal liver function tests and vanishing bile duct syndrome, (including destruction and disappearance of intrahepatic bile ducts), that may occur in various combinations. Other organs may also be affected (e.g. lungs, kidneys, pancreas, myocardium, colon) (see section 4.8).

The HLA-A*3101 allele has been found to be associated with the occurrence of hypersensitivity syndrome, including maculopapular rash.

Patients who have exhibited hypersensitivity reactions to carbamazepine should be informed that approximately 25 to 30 % of these patients may experience hypersensitivity reactions with oxcarbazepine. Cross-hypersensitivity can occur between carbamazepine and phenytoin.

In general, if signs and symptoms suggestive of hypersensitivity reactions occur, Gericarb should be withdrawn immediately.

Seizures

Gericarb should be used with caution in patients with mixed seizures, which include absences, either typical or atypical. In all these conditions, Gericarb may exacerbate seizures. In case of exacerbation of seizures, Gericarb should be discontinued.

An increase in seizure frequency may occur during switchover from an oral formulation to suppositories.

Abrupt withdrawal of Gericarb may precipitate seizures.

Hepatic function

Liver function tests should also be performed before commencing treatment and periodically thereafter, particularly in patients with a history of liver disease and in elderly patients. The drug should be withdrawn immediately in cases of aggravated liver dysfunction or acute liver disease.

Some liver function tests in patients receiving carbamazepine may be found to be abnormal, particularly gamma glutamyl transferase. This is probably due to hepatic enzyme induction. Enzyme induction may also produce modest elevations in alkaline phosphatase. These enhancements of hepatic metabolising capacity are not an indication of the withdrawal of carbamazepine.

Severe hepatic reactions to carbamazepine occur very rarely. The development of signs and symptoms of liver dysfunction or active liver disease should be urgently evaluated and treatment with Gericarb suspended pending the outcome of the evaluation.

Renal function

Baseline and periodic complete urinalysis and BUN determinations are recommended.

Hyponatremia

Hyponatremia is known to occur with carbamazepine. In patients with pre-existing renal conditions associated with low sodium or in patients treated concomitantly with sodium-lowering medicinal products (e.g. diuretics, medicinal products associated with inappropriate ADH secretion), serum sodium levels should be measured prior to initiating carbamazepine therapy. Thereafter, serum sodium levels should be measured after approximately two weeks and then at monthly intervals for the first three months during therapy, or according to clinical need. These risk factors may apply especially to elderly patients. If hyponatraemia is observed, water restriction is an important counter-measurement if clinically indicated.

Hypothyroidism

Carbamazepine may reduce serum concentrations of thyroid hormones through enzyme induction requiring an increase in dose of thyroid replacement therapy in patients with hypothyroidism. Hence thyroid function monitoring is suggested to adjust the dosage of thyroid replacement therapy.

Anticholinergic effects

Gericarb has shown mild anticholinergic activity; patients with glaucoma and urinary retention should therefore be warned and advised regarding possible hazards.

Psychiatric effects

The possibility of activation of a latent psychosis, and in elderly patients the possibility of agitation or confusion, especially when high doses of Gericarb are administered should be borne in mind.

Suicidal ideation and behaviour

Suicidal ideation and behaviour have been reported in patients treated with antiepileptic agents in several indications. A meta-analysis of randomised placebo controlled trials of anti-epileptic drugs has also shown a small increased risk of suicidal ideation and behaviour. The mechanism of this risk is not known and the available data do not exclude the possibility of an increased risk for carbamazepine. Therefore patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

Endocrinology

The induction of hepatic enzymes by carbamazepine may reduce the activity of the hormones contained in the combined oral-contraceptive pill. This may appear clinically as breakthrough bleeding or spotting. Breakthrough bleeding has been reported in women taking Gericarb while using hormonal contraceptives; the reliability of oral contraceptives may be adversely affected by Gericarb and women of childbearing age should be advised to consider using alternative forms of birth control while taking Gericarb.

Monitoring plasma levels

Although correlations between dosage and plasma levels of carbamazepine and between plasma levels and clinical efficacy or tolerability are rather tenuous, monitoring of the plasma levels may be useful in the following situations: dramatic increase in seizure frequency; during pregnancy; when treating children or adolescents; in suspected absorption disorders; for verification of compliance; in suspected toxicity where more than one drug is being used (see section 4.5).

Dose reduction and withdrawal

Abrupt withdrawal of Gericarb may precipitate seizures, therefore carbamazepine should be withdrawn gradually over a 6-month period. If treatment with Gericarb has to be withdrawn abruptly, the switch to another anti-epileptic drug should if necessary be effected under the cover of a suitable drug.

There have been a few cases of neonatal seizures and / or respiratory depression associated with maternal Gericarb and other concomitant anticonvulsant drug use. A few cases of neonatal vomiting, diarrhoea and / or decreased feeding have also been reported in association with maternal Gericarb use. These reactions may represent a neonatal withdrawal syndrome.

Interactions

Co-administration of inhibitors of CYP3A4 or inhibitors of epoxide hydrolase with carbamazepine can induce adverse reactions (increase of carbamazepine or carbamazepine-10,11 epoxide plasma concentrations respectively). The dosage of Gericarb should be adjusted accordingly and/or the plasma levels monitored.

Co-administration of CYP3A4 inducers with carbamazepine may decrease carbamazepine plasma concentrations and its therapeutic effect, while discontinuation of a CYP3A4 inducer may increase carbamazepine plasma concentrations. The dosage of Gericarb may have to be adjusted.

Carbamazepine is a potent inducer of CYP3A4 and other phase I and phase II enzyme systems in the liver, and may therefore reduce plasma concentrations of co-medications mainly metabolized by CYP3A4 by induction of their metabolism (see section 4.5).

Female patients of childbearing potential should be warned that the concurrent use of Gericarb with hormonal contraceptives may render this type of contraceptive ineffective (see sections 4.5 and 4.6). Alternative non-hormonal forms of contraception are recommended when using Gericarb.

4.5 Interaction with other medicinal products and other forms of interaction

Cytochrome P4503A4 (CYP3A4) is the main enzyme catalysing formulation of the active metabolite carbamazepine 10, 11 epoxide. Co-administration of inhibitors of CYP3A4 may result in increased carbamazepine plasma concentrations, which could induce adverse reactions. Co-administration of CYP3A4 inducers might increase the rate of Gericarb metabolism, thus leading to a potential decreases in the carbamazepine serum level and potential decrease in the therapeutic effect. Similarly, discontinuation of a CYP3A4 inducer may decrease the rate of metabolism of carbamazepine, leading to an increase in carbamazepine plasma levels.

Carbamazepine is a potent inducer of CYP3A4 and other phase I and phase II enzyme systems in the liver, and may therefore reduce plasma concentrations of co-medications mainly metabolized by CYP3A4 by induction of their metabolism.

Human microsomal epoxide hydrolase has been identified as the enzyme responsible for the formation of the 10,11-transdiol derivative from carbamazepine-10,11 epoxide. Co-administration of inhibitors of human microsomal epoxide hydrolase may result in increased carbamazepine-10,11 epoxide plasma concentrations.

Interactions resulting in a contraindication

The use of Gericarb is contraindicated in combination with monoamine-oxidase inhibitors (MAOIs); before administering Gericarb MAOIs should be discontinued for a minimum of 2 weeks, or longer if the clinical situation permits (see section 4.3).

Agents that may raise carbamazepine plasma levels:

Since raised plasma carbamazepine levels may result in adverse reactions (e.g. dizziness, drowsiness, ataxia, diplopia), the dosage of Gericarb should be adjusted accordingly and /or the plasma levels monitored when used concomitantly with the substances described below:

Analgesics: anti-inflammatory drugs: dextropropoxyphene, ibuprofen.

Androgens: danazol.

Antibiotics: macrolide antibiotics (e.g. erythromycin, troleandomycin, josamycin, clarithromycin), ciprofloxacin.

Antidepressants: possibly desipramine, fluoxetine, fluvoxamine, nefazodone, paroxetine, trazodone, viloxazine.

Antiepileptics: stiripentol, vigabatrin.

Antifungals: azoles (e.g. itraconazole, ketoconazole, fluconazole, voriconazole). Alternative anti-convulsants may be recommended in patients treated with voriconazole or itraconazole.

Antihistamines: loratadine, terfenadine.

Antipsychotics: olanzapine.

Antituberculosis: isoniazid.

Antivirals: protease inhibitors for HIV treatment (e.g. ritonavir).

Carbonic anhydrase inhibitors: acetazolamide.

Cardiovascular drugs: diltiazem, verapamil.

Gastrointestinal drugs: possibly cimetidine, omeprazole.

Muscle relaxants: oxybutynin, dantrolene.

Platelet aggregation inhibitors: ticlopidine.

Other interactions: grapefruit juice, nicotinamide (only in high dosage).

Agents that may raise the active metabolite carbamazepine 10, 11-epoxide plasma levels

Since raised plasma carbamazepine-10, 11-epoxide levels may result in adverse reactions (e.g. dizziness, drowsiness, ataxia, diplopia), the dosage of Gericarb should be adjusted accordingly and /or the plasma levels monitored when used concomitantly with the substances described below:

Loxapine, quetiapine, primidone, progabide, valproic acid, valnoctamide and valpromide.

Agents that may decrease carbamazepine plasma levels:

The dose of Gericarb may have to be adjusted when used concomitantly with the substances described below.

Antiepileptics: felbamate, methsuximide, oxcarbazepine, phenobarbital, phenoximide, phenytoin (to avoid phenytoin intoxication and subtherapeutic concentrations of carbamazepine it is recommended to adjust the plasma concentration of phenytoin to 13 micrograms /mL before adding carbamazepine to the treatment) and fosphenytoin, primidone, and, although the data are partly contradictory, possibly also clonazepam

Antineoplastics: cisplatin or doxorubicin.

Antimalarials: mefloquine, may antagonise the anticonvulsant effect of carbamazepine

Antituberculosis: rifampicin.

Bronchodilators or anti-asthma drugs: theophylline, aminophylline.

Dermatological drugs: isotretinoin has been reported to alter the bioavailability and/or clearance of carbamazepine and carbamazepine 10,11 epoxide. Carbamazepine levels should be monitored.

Other interactions: herbal preparations containing St John's wort (*Hypericum perforatum*)

Effect of Carbamazepine on plasma levels of concomitant agents:

Plasma or whole blood concentrations of carbamazepine can be reduced by concomitant use of the herbal preparation St John's wort (*Hypericum perforatum*). This is due to induction of drug metabolising enzymes by St John's wort. Herbal preparations containing St John's wort should therefore not be combined with Gericarb. The inducing effect may persist for at least 2 weeks after cessation of treatment with St. John's wort. If a patient is already taking St John's wort check carbamazepine blood levels and stop St John's wort. Carbamazepine levels may increase on stopping St John's wort. The dose of carbamazepine may need adjusting.

Carbamazepine may lower the plasma level, or diminish - or even abolish - the activity of certain drugs. The dosage of the following drugs may have to be adjusted to clinical requirements:

Analgesics, anti-inflammatory agents: buprenorphine, methadone, paracetamol, (long term administration of carbamazepine and paracetamol (acetaminophen) may be associated with hepatotoxicity), phenazone (antipyrine), tramadol.

Antibiotics: doxycycline, rifabutin.

Anticoagulants: oral anticoagulants (e.g. warfarin, phenprocoumon, dicoumarol and acenocoumarol).

Antidepressants: bupropion, citalopram, mianserin, nefazodone, sertraline, trazodone, tricyclic antidepressants (e.g. imipramine, amitriptyline, nortriptyline, clomipramine).

Antiemetics: aprepitant

Antiepileptics: clobazam, clonazepam, ethosuximide, felbamate, lamotrigine, oxcarbazepine, primidone, tiagabine, topiramate, valproic acid, zonisamide. To avoid phenytoin intoxication and subtherapeutic concentrations of carbamazepine it is recommended to adjust the plasma concentration of phenytoin to 13 micrograms /mL before adding carbamazepine to the treatment. There have been rare reports of an increase in plasma mephenytoin levels.

Antifungals: itraconazole, voriconazole. Alternative anti-convulsants may be recommended in patients treated with voriconazole or itraconazole.

Antihelmintics: praziquantel, albendazole.

Antineoplastics: imatinib, cyclophosphamide, lapatinib, temsirolimus.

Antipsychotics: clozapine, haloperidol and bromperidol, olanzapine, quetiapine, risperidone, ziprasidone, aripiprazole, paliperidone.

Antivirals: protease inhibitors for HIV treatment (e.g. indinavir, ritonavir, saquinavir).

Anxiolytics: alprazolam, midazolam.

Bronchodilators or anti-asthma drugs: theophylline.

Contraceptives: hormonal contraceptives (alternative contraceptive methods should be considered).

Cardiovascular drugs: digoxin, calcium channel blockers (dihydropyridine group) e.g. felodipine, simvastatin, atorvastatin, lovastatin, cerivastatin, ivabradine.

Corticosteroids: corticosteroids (e.g. prednisolone, dexamethasone).

Drugs used in erectile dysfunction: tadalafil.

Immunosuppressants: ciclosporin, everolimus, tacrolimus, sirolimus.

Thyroid agents: levothyroxine.

Other drug interactions: products containing oestrogens and/or progesterones (gestrinone, tribolone, toremifene)

Plasma phenytoin levels have been reported both to be raised and to be lowered by carbamazepine, and plasma mephenytoin levels have been reported in rare instances to increase.

Combinations that require specific consideration:

Co-administration of carbamazepine and paracetamol may reduce the bioavailability paracetamol / acetaminophen.

Concomitant use of carbamazepine and levetiracetam has been reported to increase carbamazepine-induced toxicity.

Concomitant use of carbamazepine and isoniazid has been reported to increase isoniazid hepatotoxicity.

Combined use of carbamazepine and lithium or metoclopramide on the one hand, and carbamazepine and neuroleptics (haloperidol, thioridazine) on the other, may lead to increased neurological adverse reactions (with the latter combination even in the presence of 'therapeutic plasma levels').

Concomitant medication with carbamazepine and some diuretics (hydrochlorothiazide, furosemide) may lead to symptomatic hyponatraemia.

Carbamazepine may antagonise the effects of non-depolarising muscle relaxants (e.g. pancuronium); their dosage should be raised and patients monitored closely for a more rapid recovery from neuromuscular blockade than expected.

Carbamazepine, like other psychoactive drugs, may reduce alcohol tolerance; it is therefore advisable for the patient to abstain from alcohol.

Interference with serological testing

Carbamazepine may result in false positive perphenazine concentrations in HPLC analysis due to interference.

Carbamazepine and the 10,11-epoxide metabolite may result in false positive tricyclic antidepressant concentration in fluorescence polarized immunoassay method.

4.6 Fertility, pregnancy and lactation

Pregnancy

Offspring of epileptic mothers are known to be more prone to developmental disorders, including malformations. Although conclusive evidence from controlled studies with carbamazepine monotherapy is lacking, developmental disorders and malformations, including spina bifida and also other congenital anomalies, e.g. craniofacial defects, cardiovascular malformations, hypospadias and anomalies involving various body systems, have been reported in association with Gericarb. Based on data in a North American pregnancy registry, the rate of major congenital malformations, defined as a structural abnormality with surgical, medical, or cosmetic importance, diagnosed within 12 weeks of birth was 3.0% (95% CI 2.1 to 4.2%) among mothers exposed to carbamazepine monotherapy in the first trimester and 1.1% (95% CI 0.35 to 2.5%) among pregnant women not taking any antiepileptic drug (relative risk 2.7, 95% CI 1.1 to 7.0).

Taking these data into consideration:

- Pregnant women with epilepsy should be treated with special care
- If women receiving Gericarb become pregnant or plan to become pregnant, or if the problem of initiating treatment with Gericarb arises during pregnancy, the drug's potential benefits must be carefully weighed against its possible hazards, particularly in the first 3 months of pregnancy.
- In women of childbearing age Gericarb should, wherever possible, be prescribed as monotherapy, because the incidence of congenital abnormalities in the offspring of women treated with a combination of antiepileptic drugs is greater than in those of mothers receiving the individual drugs as monotherapy. The risk of malformations following exposure to carbamazepine as polytherapy may vary depending on the specific drugs used and may be higher in polytherapy combinations that include valproate.
- Minimum effective doses should be given and monitoring of plasma levels is recommended. The plasma concentration could be maintained in the lower side of the therapeutic range 4 to 12 micrograms/mL provided seizure control is maintained. There is evidence to suggest that the risk of malformation with carbamazepine may be dose-dependent i.e. at a dose < 400mg per day, the rates of malformation were lower than with higher doses of carbamazepine.
- Patients should be counselled regarding the possibility of an increased risk of malformations and given the opportunity of antenatal screening.
- During pregnancy, an effective antiepileptic treatment must not be interrupted, since the aggravation of the illness is detrimental to both the mother and the foetus.

Monitoring and prevention

Folic acid deficiency is known to occur in pregnancy. Antiepileptic drugs have been reported to aggravate folic acid deficiency. This deficiency may contribute to the increased incidence of birth defects in the offspring of treated epileptic women. Folic acid supplementation is recommended before and during pregnancy.

In the neonate

In order to prevent bleeding disorders in the offspring, it has also been recommended that vitamin K1 be given to the mother during the last weeks of pregnancy as well as to the neonate.

There have been a few cases of neonatal seizures and/or respiratory depression associated with maternal Gericarb and other concomitant anticonvulsant drug use. A few cases of neonatal vomiting, diarrhoea and/or decreased feeding have also been reported in association with maternal Gericarb use. These reactions may represent a neonatal withdrawal syndrome.

Women of child-bearing potential and contraceptive measures

Due to enzyme induction, Gericarb may result in a failure of the therapeutic effect of oral contraceptive drugs containing oestrogen and/or progesterone. Women of child bearing potential should be advised to use alternative contraceptive methods while on treatment with Gericarb.

Breast-feeding

Although carbamazepine passes into the breast milk in concentrations of about 25-60% of the plasma level, this is not believed to present a significant hazard to the infant, which is likely to receive at most 10% of an appropriate therapeutic dose of carbamazepine for an infant with epilepsy. As with all drugs, the benefits of breast-feeding should be weighed against the remote possibility of an adverse effect occurring in the infant. Mothers taking Gericarb may breastfeed their infants, provided the infant is observed for possible adverse reactions (e.g. excessive somnolence, allergic skin reaction). There have been some reports of cholestatic hepatitis in neonates exposed to carbamazepine during antenatal and or during breast feeding. Therefore breast-fed infants of mothers treated with carbamazepine should be carefully observed for adverse hepatobiliary effects.

Fertility

There have been very rare reports of impaired male fertility and/or abnormal spermatogenesis.

4.7 Effects on ability to drive and use machines

The patients’ ability to drive or operate machinery may be impaired by medical condition resulting in seizures and adverse reactions including dizziness, drowsiness ataxia, diplopia, impaired accommodation and blurred vision reported with Gericarb, especially in the early stages of treatment. Patients should be warned of the possible hazard when driving or operating machinery.

4.8 Undesirable effects

Particularly at the start of treatment with Gericarb, or if the initial dosage is too high or when treating elderly patients, certain types of adverse reaction occur very commonly or commonly, e.g. CNS adverse reactions (dizziness, headache, ataxia, drowsiness, fatigue, diplopia); gastrointestinal disturbances nausea, vomiting), and allergic skin reactions.

The dose-related adverse reactions usually abate within a few days, either spontaneously or after a transient dosage reduction. The occurrence of CNS adverse reactions may be a manifestation of relative overdosage or significant fluctuation in plasma levels. In such cases it is advisable to monitor the plasma levels and divide the daily dosage into smaller (i.e. 3-4) fractional doses.

There have been reports of decreased bone mineral density, osteopenia, osteoporosis and fractures in patients on long-term therapy with carbamazepine. The mechanism by which Gericarb affects bone metabolism has not been identified.

Tabulated summary of adverse drug reactions compiled from clinical trials and from spontaneous reports
Adverse drug reactions from clinical trials are listed by MedDRA system organ class. Within each system organ class, the adverse drug reactions are ranked by frequency, with the most frequent reactions first. Within each frequency grouping, adverse drug reactions are presented in order of decreasing seriousness.

In addition, the corresponding frequency category for each adverse drug reaction is based on the following convention (CIOMS III): very common (≥ 1/10); common (≥ 1/100, < 1/10); uncommon (≥ 1/1000, < 1/100); rare (≥ 1/10 000, < 1/1000); very rare (< 1/10 000).

Table 1

Blood and lymphatic system disorders		
Very common:	Leukopenia.	
Common:	thrombocytopenia, eosinophilia.	
Rare:	Leukocytosis, lymphadenopathy.	
Very rare:	Agranulocytosis, aplastic anaemia, pancytopenia, pure red cell aplasia, anaemia, anaemia megaloblastic, reticulocytosis haemolytic anaemia.	
Immune system disorders		
Rare:	A delayed multi-organ hypersensitivity disorder (of serum sickness type) with fever, rashes, vasculitis, lymphadenopathy, pseudo lymphoma, arthralgia, leukopenia, eosinophilia, hepato-splenomegaly and abnormal liver function tests, occurring in various combinations. Other organs may also be affected (e.g. lungs, kidneys, pancreas, myocardium, colon). Treatment must be discontinued immediately if such hypersensitivity reactions occur.	
Very rare:	Anaphylactic reaction, angioedema, hypogammaglobulinaemia.	

Endocrine disorders		
Common:	Oedema, fluid retention, weight increase, hyponatraemia and blood osmolarity decreased due to an antidiuretic hormone (ADH)-like effect leading in rare cases to water intoxication accompanied by lethargy, vomiting, headache, confusional state, neurological disorders.	
Very rare:	Galactorrhoea, gynecomastia.	
Metabolism and nutrition disorders		
Rare:	Folate deficiency, decreased appetite.	
Very rare:	Porphyria acute (acute intermittent porphyria and variegate porphyria), porphyria non-acute (porphyria cutanea tarda).	
Psychiatric disorders		
Rare:	Hallucinations (visual or auditory), depression, restlessness, aggression, agitation, confusional state.	
Very rare:	Activation of psychosis.	
Nervous system disorders		
Very common:	Dizziness, ataxia, somnolence.	
Common:	Headache, diplopia.	
Uncommon:	Abnormal involuntary movements (e.g. tremor, asterixis, dystonia, tics); nystagmus.	
Rare:	Dyskinesia, eye movement disturbances, speech disorders (e.g. dysarthria, slurred speech), choreoathetosis, neuropathy peripheral, paraesthesia paresis.	
Very rare:	Neuroleptic malignant syndrome, aseptic meningitis with myoclonus and peripheral eosinophilia, dysgeusia.	
Eye disorders		
Common:	Accommodation disorders (e.g. blurred vision).	
Very rare:	Lenticular opacities, conjunctivitis.	
Ear and labyrinth disorders		
Very rare:	Hearing disorders, e.g. tinnitus, hyperacusis, hypoacusis, change in pitch perception.	
Cardiac disorders		
Rare:	Cardiac conduction disorders.	
Very rare:	Bradycardia, arrhythmia, atrioventricular block with syncope, cardiac failure congestive, coronary artery disease aggravated.	
Vascular disorders		
Rare:	Hypertension or hypotension.	
Very rare:	Circulatory collapse, embolism (e.g. pulmonary embolism), thrombophlebitis	
Respiratory, thoracic and mediastinal disorders		
Very rare:	Pulmonary hypersensitivity characterized e.g. by fever, dyspnoea, pneumonitis or pneumonia.	

Gastrointestinal disorders	
Very common:	Nausea, vomiting.
Common:	Dry mouth; with suppositories, rectal irritation may occur.
Uncommon:	Diarrhoea, constipation.
Rare:	Abdominal pain.
Very rare:	Glossitis, stomatitis, pancreatitis.
Hepatobiliary disorders	
Rare:	Hepatitis of cholestatic, parenchymal (hepatocellular) or mixed type, vanishing bile duct syndrome (destruction and disappearance of the intrahepatic bile duct), jaundice.
Very rare:	Granulomatous hepatitis, hepatic failure.
Skin and subcutaneous tissue disorders	
Very common:	Dermatitis allergic, urticaria which may be severe.
Uncommon:	Exfoliative dermatitis.
Rare:	Systemic lupus erythematosus, pruritus.
Very rare:	Severe cutaneous adverse reactions (SCARs) e.g. Stevens-Johnson syndrome (SJS)*, toxic epidermal necrolysis (TEN) (see section 4.4), , photosensitivity reaction, erythema multiforme erythema nodosum, pigmentation disorder, purpura, acne, hyperhidrosis, hair loss, hirsutism
Musculoskeletal, connective tissue and bone disorders	
Rare	Muscular weakness
Very rare:	Bone metabolism disorders (decrease in plasma calcium and blood 25-hydroxycholecalciferol) leading to osteomalacia / osteoporosis, arthralgia, myalgia, muscle spasms.
Renal and urinary disorders	
Very rare:	Tubulointerstitial nephritis, renal failure, renal impairment (e.g. albuminuria, haematuria, oliguria, and blood urea increased/azotemia), urinary frequency, urinary retention.
Reproductive system	
Very rare:	Sexual dysfunction/impotence, spermatogenesis abnormal (with decreased sperm count and/or motility).
General disorders and administration site conditions	
Very common:	Fatigue
Investigations	
Very common:	Gamma-glutamyl transferase increased (due to hepatic enzyme induction), usually not clinically relevant.
Common:	Blood alkaline phosphatase increased.
Uncommon:	Transaminases increased.
Very rare:	Intraocular pressure increased, blood cholesterol increased, high density lipoprotein increased, blood triglycerides increased.

Thyroid function test abnormal: decreased L-Thyroxin (free thyroxine, thyroxine, tri-iodothyronine) and increased blood thyroid stimulating hormone, usually without clinical manifestations, blood prolactin increased.

* In some Asian countries also reported as rare. See also section 4.4.

Additional adverse drug reactions from spontaneous reports (frequency not known)

The following adverse drug reactions have been derived from post-marketing experience with Gericarb via spontaneous case reports and literature cases. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency which is therefore categorized as not known. Adverse drug reactions are listed according to system organ classes in MedDRA. Within each system organ class, ADRs are presented in order of decreasing seriousness.

Infections and infestations: Reactivation of Human herpesvirus 6 infection.

Blood and lymphatic system disorders: Bone marrow failure.

Nervous system disorders: Sedation, memory impairment.

Gastrointestinal disorders: Colitis.

Immune system disorders: Drug Rash with Eosinophilia and Systemic Symptoms (DRESS).

Skin and subcutaneous tissue disorders: Acute Generalized Exanthematous Pustulosis (AGEP), lichenoid keratosis, onychomadesis.

Musculoskeletal and connective tissue disorders: Fracture.

Investigations: Bone density decreased.

There is increasing evidence regarding the association of genetic markers and the occurrence of cutaneous ADRs such as SJS, TEN, DRESS, AGEP and maculopapular rash. In Japanese and European patients, these reactions have been reported to be associated with the use of carbamazepine and the presence of the HLA-A*3101 allele. Another marker, HLA-B*1502 has been shown to be strongly associated with SJS and TEN among individuals of Han Chinese, Thai and some other Asian ancestry (see sections 4.2 and 4.4 for further information).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517; Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

By reporting side effects you can help provide more information on the safety of this medicine.

4.9 Overdose

Signs and symptoms:

The presenting signs and symptoms of overdosage involve the central nervous, cardiovascular or respiratory systems and the adverse drug reactions mentioned under section 4.8.

Central nervous system:

CNS depression; disorientation, depressed level of consciousness, somnolence, agitation, hallucination, coma; blurred vision, slurred speech, dysarthria, nystagmus, ataxia, dyskinesia, initially hyper-reflexia, later hyporeflexia; convulsions, psychomotor disturbances, myoclonus, hypothermia, mydriasis.

Respiratory system:

Respiratory depression, pulmonary oedema.

Cardiovascular system:

Tachycardia, changes in blood pressure (hypotension and at times hypertension), cardiac arrhythmias, conduction disturbance with widening of QRS complex; syncope, in association with cardiac arrest.

Gastrointestinal system:

Vomiting, delayed gastric emptying, reduced bowel motility.

Musculoskeletal system:

There have been some cases which reported rhabdomyolysis in association with carbamazepine toxicity.

Renal function:

Retention of urine, oliguria or anuria; fluid retention, water intoxication due to ADH-like effect of carbamazepine.

Laboratory findings:

Hyponatraemia, possibly metabolic acidosis, possibly hyperglycaemia, increased muscle creatinine phosphokinase.

Treatment:

There is no specific antidote to Gericarb.

Management according to the patient's clinical condition. Possible admission to hospital. Measurement of the plasma level to confirm carbamazepine poisoning and to ascertain the size of the overdose. Evacuation of the stomach, gastric lavage, and administration of activated charcoal. Delay in evacuating the stomach may result in delayed absorption, leading to relapse during recovering from intoxication. Supportive medical care in an intensive care unit with cardiac monitoring and careful correction of electrolyte imbalance, if required.

Special recommendations:

Charcoal haemoperfusion has been recommended. Haemodialysis is the effective treatment modality in the management of the carbamazepine overdose.

Relapse and aggravation of symptomatology on the 2nd and 3rd day after overdose, due to delayed absorption, should be anticipated.

5 PHARMACOLOGICAL PROPERTIES**5.1 Pharmacodynamic properties****Pharmacotherapeutic group:**

Antiepileptic, neurotropic, and psychotropic agent. (ATC Code: N03 AF01).

Dibenzazepine derivative.

Mechanism of action:

As an antiepileptic agent its spectrum of activity embraces: partial seizures (simple and complex) with and without secondary generalization; generalized tonic-clonic seizures, as well as combinations of these types of seizures.

The mechanism of action of carbamazepine, the active substance of Gericarb, has only been partially elucidated. Carbamazepine stabilizes hyperexcited nerve membranes, inhibits repetitive neuronal discharges, and reduces synaptic propagation of excitatory impulses. It is conceivable that prevention of repetitive firing of sodium-dependent action potentials in depolarized neurons via use- and voltage-dependent blockade of sodium channels may be its main mechanism of action.

Whereas reduction of glutamate release and stabilization of neuronal membranes may account mainly for the antiepileptic effects, the depressant effect on dopamine and noradrenaline turnover could be responsible for the antimanic properties of carbamazepine.

5.2 Pharmacokinetic properties

Absorption:

Carbamazepine is absorbed almost completely but relatively slowly from the tablets. The conventional tablets yield mean peak plasma concentrations of the unchanged substance within 12 hours, following single oral doses. With the oral suspension, mean peak plasma concentrations are attained within 2 hours. With respect to the amount of active substance absorbed, there is no clinically relevant difference between the oral dosage forms. After a single oral dose of 400 mg carbamazepine (tablets) the mean peak concentration of unchanged carbamazepine in the plasma is approx. 4.5 micrograms/mL.

The bioavailability of carbamazepine in various oral formulations has been shown to lie between 85-100%.

Different preparations of carbamazepine may vary in bioavailability; to avoid reduced effect or risk of breakthrough seizures or excessive side effects, it may be prudent to avoid changing the formulation.

Ingestion of food has no significant influence on the rate and extent of absorption, regardless of the dosage form of Gericarb.

Steady-state plasma concentrations of carbamazepine are attained within about 1 to 2 weeks, depending individually upon auto-induction by carbamazepine and hetero-induction by other enzyme-inducing drugs, as well as on pretreatment status, dosage, and duration of treatment.

The steady-state plasma concentrations of carbamazepine considered as 'therapeutic range' vary considerably interindividually: for the majority of patients a range between 4 to 12 micrograms/mL corresponding to 17 to 50 micromol/L has been reported. Concentrations of carbamazepine-10,11-epoxide (pharmacologically active metabolite): about 30% of carbamazepine levels .

Distribution:

Carbamazepine is bound to serum proteins to the extent of 70 to 80%. The concentration of unchanged substance in cerebrospinal fluid and saliva reflects the non-protein bound portion in the plasma (20 to 30%). Concentrations in breast milk were found to be equivalent to 25 to 60% of the corresponding plasma levels.

Carbamazepine crosses the placental barrier. Assuming complete absorption of carbamazepine, the apparent volume of distribution ranges from 0.8 to 1.9 L/kg.

Biotransformation / metabolism:

Carbamazepine is metabolised in the liver where the epoxide pathway of biotransformation is the most important one, yielding the 10,11-transdiol derivative and its glucuronide as the main metabolites. Cytochrome P450 3A4 has been identified as the major isoform responsible for the formation of the pharmacologically active carbamazepine-10,11 epoxide from carbamazepine. Human microsomal epoxide hydrolase has been identified as the enzyme responsible for the formation of the 10,11-transdiol derivative from carbamazepine-10,11 epoxide . 9-Hydroxy-methyl-10-carbamoyl acridan is a minor metabolite related to this pathway. After a single oral dose of carbamazepine about 30% appears in the urine as end-products of the epoxide pathway. Other important biotransformation pathways for carbamazepine lead to various monohydroxylated compounds, as well as to the N-glucuronide of carbamazepine produced by UGT2B7.

Elimination:

Elimination half life of unchanged carbamazepine averages approx. 36 hours following a single oral dose, whereas after repeated administration it averages only 16-24 hours (auto-induction of the hepatic mono-oxygenase system), depending on the duration of the medication. In patients receiving concomitant treatment with other liver-enzyme inducing drugs (e.g. phenytoin, phenobarbitone), half-life values averaging 9 to 10 hours have been found.

The mean elimination half-life of the 10,11-epoxide metabolite in the plasma is about 6 hours following single oral doses of the epoxide itself.

After administration of a single oral dose of 400 mg carbamazepine, 72% is excreted in the urine and 28% in the faeces. In the urine, about 2% of the dose is recovered as unchanged drug and about 1% as the pharmacologically active 10,11-epoxide metabolite.

Special Populations

Patients with hepatic and renal impairment

No data is available on the pharmacokinetics of carbamazepine in patients with impaired hepatic or renal function. In advanced hepatic disease carbamazepine metabolism may be impaired.

Elderly

There is no indication of altered pharmacokinetics of carbamazepine in elderly patients as compared with young adults but its metabolism may be affected by hepatic dysfunction.

Children

Owing to enhanced carbamazepine elimination, children may require higher doses of carbamazepine (in mg/kg) than adults.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, single and repeated dose toxicity, genotoxicity and carcinogenic potential. However, the animal studies were insufficient to rule out a teratogenic effect of carbamazepine.

Carcinogenicity

In rats treated with carbamazepine for 2 years, there was an increased incidence of hepatocellular tumors in females and benign testicular tumors in males.

However, there is no evidence that these observations are of any relevance to the therapeutic use of carbamazepine in humans.

Genotoxicity

Carbamazepine was not found to be genotoxic in various standard bacterial and mammalian mutagenicity studies.

Reproductive toxicity

The cumulative evidence from various animal studies in mice, rats and rabbits indicates that carbamazepine has no or only minor teratogenic potential at doses relevant to man. However, the animal studies were insufficient to rule out a teratogenic effect of carbamazepine. In a reproduction study in rats, nursing offspring demonstrated a reduced weight gain at a maternal dosage level of 192 mg/kg/day.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Microcrystalline Cellulose
 Pregelatinised Maize Starch
 Colloidal Anhydrous Silica
 Talc
 Sodium Starch Glycollate (Type A)
 Magnesium Stearate

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Do not store above 25°C.

Store in the original container.

6.5 Nature and contents of container

Polypropylene containers with polyethylene caps and optional polyethylene filler in packs of 100, 250 or 500 tablets.

Not all pack sizes may be marketed

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Generics (UK) Limited
12 Station Close
Potters Bar
Hertfordshire EN6 1TL
United Kingdom

8 MARKETING AUTHORISATION NUMBER

PA0405/016/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 02 February 1989

Date of last renewal: 02 February 2009

10 DATE OF REVISION OF THE TEXT

August 2015