

Part II

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Iliadin Mini Paediatric 0.025% w/v Nasal Drops, solution.

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Oxymetazoline Hydrochloride 0.025% w/v

For excipients, see 6.1

3 PHARMACEUTICAL FORM

Nasal drops, solution.

A clear, colourless solution.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

For the relief of nasal congestion associated with disorders of the upper respiratory tract including infective and allergic rhinitis, sinusitis, nasopharyngitis and coryza

4.2 Posology and method of administration

For children under 6 years

Instil half the contents of one mini unit into each nostril: repeat every 8-12 hours if necessary.

Intranasally

4.3 Contraindications

Use in patients with known hypersensitivity to sympathomimetics.

Use in patients receiving monoamine oxidase inhibitors or within fourteen days of stopping such treatment.

Use in acute coronary disease, cardiac asthma, thyrotoxicosis, hypertension, closed-angle glaucoma or urinary retention.

4.4 Special warnings and special precautions for use

Use with great care in patients suffering from angina and digitalized patients.

This product should be given with care to patients with prostatic enlargement, as it may increase difficulty in micturition.

Prolonged use may cause rebound congestion and drug induced rhinitis.

Use with caution in diabetic patients as this product may cause an increase in blood sugar.

The physician or pharmacist should reassure himself/herself sympathomimetic containing preparations are not simultaneously administered by several routes i.e. orally and topically (nasal, aural and eye preparations).

When considering use in infants and young children under 2 years old, great caution should be exercised. Serious side effects including malignant neuroleptic syndrome, hypertension, central nervous system irritation with screaming attacks, muscular rigidity and convulsions have been reported in this age group. However, the likelihood of their occurrence is reduced when using the controlled dosage units of Iliadin Mini Paediatric.

This product should not be used for more than 5-7 consecutive days without medical advice.

Care should be taken not to exceed the stated dose

This product should not be used by patients already taking medication without first consulting their physician.

4.5 Interaction with other medicinal products and other forms of interaction

Concurrent use with halogenated anaesthetic agents may provoke or worsen ventricular arrhythmias.

The effects of this product may be altered by guanethidine, reserpine, methyldopa, phenothiazines and tricyclic anti depressants.

4.6 Pregnancy and lactation

Iliadin Drops should not be used in pregnancy unless considered essential by the physician.

4.7 Effects on ability to drive and use machines

None

4.8 Undesirable effects

Side effects may include local stinging, irritation and dryness.
Prolonged use may cause rebound vasodilatation and chemical rhinitis.

4.9 Overdose

No experience of overdose, but supportive measures would be the appropriate treatment.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Oxymetazoline is a potent sympathomimetic agent which has at least ten times the potency of xylometazoline, a greater therapeutic index and greater duration of action. Acute toxicity studies in mice have indicated no unusual aspects of lethality.

5.2 Pharmacokinetic properties

Tritiated oxymetazoline has been shown to be rapidly absorbed in dogs following intranasal administration. Approximately 30% are excreted in the urine within 72 hours and faecal excretion is about 10% within 120 hours. The drug is approximately 30% bound to human plasma proteins.

5.3 Preclinical safety data

There are no preclinical data of relevance to the prescriber, which are additional to that already updated in other sections of the SPC.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium Dihydrogen Phosphate Dihydrate
Disodium Phosphate Dihydrate
Benzalkonium Chloride Solution
Sodium Hydroxide
Water for injection.

6.2 Incompatibilities

Not applicable.

6.3 Shelf Life

Unopened: 3 years

After first opening the container: Use immediately after opening. Discard any unused contents.

6.4 Special precautions for storage

Do not store above 25°C.

6.5 Nature and contents of container

0.32ml single dose polyethylene pipette mini-unit containers manufactured from low density natural polythene and heat sealed.
10 single dose units per pack.

6.6 Instructions for use and handling

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Seven Seas Limited
Trading as
Merck Consumer Health
Hedon Road
Marfleet
Kingston-Upon-Hull HU9 5NJ
United Kingdom

8 MARKETING AUTHORISATION NUMBER

PA 417/12/3

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 1st April 1979

Date of last renewal: 1st April 2004

10 DATE OF REVISION OF THE TEXT

February 2005

