

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Remifentanil 5 mg powder for concentrate for solution for injection/infusion

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

5 mg powder for concentrate for solution for injection/infusion

1 vial contains 5 mg remifentanil (as remifentanil hydrochloride).

After reconstitution the solution contains 1 mg/ml remifentanil (as hydrochloride), if prepared as recommended (see section 6.6)

Excipient(s) with known effect: sodium 1.15 mg

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Powder for concentrate for solution for injection/infusion

Lyophilized white to slightly yellow cake or powdery mass.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Remifentanil is indicated as an analgesic agent for use during induction and/or maintenance of general anaesthesia.

Remifentanil is indicated for provision of analgesia in mechanically ventilated intensive care patients 18 years of age and over.

4.2 Posology and method of administration

Remifentanil should be administered only in a setting fully equipped for the monitoring and support of respiratory and cardiovascular function and by persons specifically trained in the use of anaesthetic drugs and the recognition and management of the expected adverse effects of potent opioids, including respiratory and cardiac resuscitation. Such training must include the establishment and maintenance of a patent airway and assisted ventilation.

Continuous infusions of remifentanil must be administered by a calibrated infusion device into a fast flowing IV line or via a dedicated IV line. This infusion line should be connected at, or close to, the venous cannula to minimise the potential dead space (see section 6.6 for additional information, including tables with examples of infusion rates by body weight to help titrate remifentanil to the patient's anaesthetic needs). Care should be taken to avoid obstruction or disconnection of infusion lines and to adequately clear the lines to remove residual remifentanil after use (see section 4.4). IV lines/infusion system should be removed after cessation of use to avoid inadvertent administration.

Remifentanil may be given by target-controlled infusion (TCI) with an approved infusion device incorporating the Minto pharmacokinetic model with covariates for age and lean body mass (LBM).

Congestion of the infusion drips or tearing off should be avoided, and the infusion drips should be sufficiently rinsed in order to remove residual Remifentanil after discontinuation of medication (see also section 4.4).

Remifentanil is for intravenous use only and must not be administered by epidural or intrathecal injection (see section 4.3).

Dilution

Remifentanyl should not be administered without further dilution after reconstitution of the lyophilized powder. See section 6.3 for storage conditions and section 6.6 for recommended diluents and instructions on reconstitution / dilution of the product before administration.

For manually-controlled infusion Remifentanyl can be diluted to concentrations of 20 to 250 micrograms/ml (50 micrograms/ml is the recommended dilution for adults and 20 to 25 micrograms/ml for paediatric patients aged 1 year and over).

For TCI the recommended dilution of Remifentanyl is 20 to 50 micrograms/ml.

General Anaesthesia

The administration of Remifentanyl must be individualised based on the patient's response.

Adults

Administration by manually-controlled infusion (MCI)

DOSING GUIDELINES FOR ADULTS

	Remifentanyl BOLUS INJECTION (micrograms/kg)	Remifentanyl CONTINUOUS INFUSION (micrograms/kg/min)	
		Starting Rate	Range
	1 (within at least 30 seconds)	0.5 to 1	—
	Maintenance of anaesthesia in ventilated patients		
• Nitrous oxide (66 %)	0.5 to 1	0.4	0.1 to 2
• Isoflurane (starting dose 0.5 MAC)	0.5 to 1	0.25	0.05 to 2
• Propofol (Starting dose 100 micrograms /kg/min)	0.5 to 1	0.25	0.05 to 2

When given by bolus injection at induction Remifentanyl should be administered over not less than 30 seconds.

At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended above to avoid an increase of haemodynamic effects (hypotension and bradycardia) of remifentanyl.

No data are available for dosage recommendations for simultaneous use of other hypnotics other than those listed in the table with remifentanyl.

Induction of anaesthesia

Remifentanyl should be co-administered with a hypnotic agent, such as propofol, thiopentone, or isoflurane, for the induction of anaesthesia. Administering Remifentanyl after a hypnotic agent will reduce the incidence of muscle rigidity. Remifentanyl can be administered at an infusion rate of 0.5 to 1 microgram/kg/min, with or without an initial bolus injection of 1 microgram/kg given over not less than 30 seconds. If endotracheal intubation is to occur more than 8 to 10 minutes after the start of the infusion of Remifentanyl, then a bolus injection is not necessary.

Maintenance of anaesthesia in ventilated patients

After endotracheal intubation, the infusion rate of Remifentanyl should be decreased, according to anaesthetic technique, as indicated in the above table. Due to the fast onset and short duration of action of remifentanyl, the rate of administration during anaesthesia can be titrated upward in 25% to 100% increments or downward in 25% to 50% decrements, every 2 to 5 minutes to attain the desired level of μ -opioid response. In response to light anaesthesia, supplemental bolus injections may be administered every 2 to 5 minutes.

Anaesthesia in spontaneously breathing patients

In spontaneously breathing anaesthetised patients with a secured airway respiratory depression is likely to occur. Therefore attention must be given to respiratory effects eventually combined with muscular rigidity. Special care is needed to adjust the dose to the patient requirements and ventilatory support may be required. Adequate facilities should be available for monitoring of patients administered remifentanyl. It is essential that these facilities be fully equipped to handle all degrees of respiratory depression (intubation equipment must be available) and/or muscle rigidity (for more information see section 4.4).

The recommended starting infusion rate for supplemental analgesia in spontaneously breathing anaesthetised patients is 0.04 micrograms/kg/min with titration to effect. A range of infusion rates from 0.025 to 0.1 micrograms/kg/min has been studied.

Bolus injections are not recommended in spontaneously breathing anaesthetised patients.

Remifentanyl should not be used as an analgesic in procedures where patients remain conscious or do not receive any airway support during the procedure.

Concomitant medication

Remifentanyl decreases the amounts or doses of inhalational anaesthetics, hypnotics and benzodiazepines required for anaesthesia (see section 4.5).

Doses of the following agents used in anaesthesia: isoflurane, thiopentone, propofol and temazepam have been reduced by up to 75 % when used concurrently with remifentanyl.

Guidelines for discontinuation/continuation during immediate post-operative period

Due to the very rapid offset of action of remifentanyl no residual opioid activity will be present within 5 to 10 minutes after discontinuation. For those patients undergoing surgical procedures where post-operative pain is anticipated, analgesics should be administered prior to discontinuation of remifentanyl. Sufficient time must be allowed to reach the maximum effect of the longer acting analgesic. The choice of analgesic should be appropriate for the patient's surgical procedure and the level of post-operative care.

In the event that the longer acting analgesia has not reached the appropriate effect before the end of surgery, the administration of remifentanyl can be continued to maintain analgesia during immediate post-operative period until longer acting analgesia has reached the maximum effect.

If remifentanyl is continued post-procedural, it should only be used in a setting fully equipped for the monitoring and support of respiratory and cardiovascular function, under the close supervision of persons specifically trained in the recognition and management of the respiratory effects of potent opioids.

Furthermore it is recommended that patients should be closely monitored post-operatively for pain, hypotension and bradycardia.

Further information about the administration in mechanically ventilated intensive care patients is given in this section.

In spontaneously breathing patients the initial infusion rate of remifentanyl may be decreased to 0.1 micrograms/kg/min and thereafter can be increased or decreased every 5 min in steps of 0.025 micrograms/kg/min to balance the extent of analgesia against the degree of respiratory depression.

In spontaneously breathing patients bolus doses for analgesia are not recommended during the post-operative period.

Administration by Target-Controlled Infusion (TCI)

Induction and maintenance of anaesthesia in ventilated patients

Remifentanyl TCI should be used in association with an intravenous or inhalational hypnotic agent during the induction and maintenance of anaesthesia in ventilated adult patients (see table above for manually controlled infusion). In association with these agents, adequate analgesia for induction of anaesthesia and surgery can generally be achieved with target blood remifentanyl concentrations ranging from 3 to 8 nanograms/ml. Remifentanyl should be titrated to individual patient response. For particularly stimulating surgical procedures target blood concentrations up to 15 nanograms/ml may be required.

At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended to avoid an increase of

haemodynamic effects (hypotension and bradycardia) of remifentanyl (see table above for manually controlled infusion).

The following table provides the equivalent blood remifentanyl concentration using a TCI approach for various manually controlled infusion rates at steady state:

Remifentanyl blood concentrations (nanograms/ml) estimated using the Minto (1997) pharmacokinetic model in a 70 kg, 170 cm, 40 year old male patient for various manually controlled infusion rates (micrograms/kg/min) at steady state

Remifentanyl Infusion Rate (micrograms /kg/min)	Remifentanyl Blood Concentration (nanograms/ml)
0.05	1.3
0.10	2.6
0.25	6.3
0.40	10.4
0.50	12.6
1.0	25.2
2.0	50.5

As there are insufficient data, the administration of remifentanyl by TCI for spontaneous ventilation anaesthesia is not recommended.

Guidelines for discontinuation/continuation into the immediate post-operative period

At the end of surgery when the TCI infusion is stopped or the target concentration reduced, spontaneous respiration is likely to return at calculated remifentanyl concentrations in the region of 1 to 2 nanograms/ml. As with manually controlled infusion, post-operative analgesia should be established before the end of surgery with longer acting analgesics (see also Guidelines for discontinuation / continuation during immediate postoperative period in section above for Manually Controlled Infusion).

As there are insufficient data, the administration of Remifentanyl by TCI for the management of post-operative analgesia is not recommended.

Paediatric patients (1 to 12 years of age)

Co-administration of remifentanyl and an intravenous anaesthetic agent for induction of anaesthesia has not been studied in detail and is therefore not recommended.

Remifentanyl TCI has not been studied in paediatric patients and therefore administration of remifentanyl by TCI is not recommended in these patients.

When given by bolus injection, Remifentanyl should be administered over not less than 30 seconds. Surgery should not commence until at least 5 minutes after the start of the Remifentanyl infusion, if a simultaneous bolus dose has not been given.

For sole administration of nitrous oxide (70%) with Remifentanyl, typical maintenance infusion rates should be between 0.4 and 3 micrograms/kg/min, and although not specifically studied, adult data suggest that 0.4 micrograms/kg/min is an appropriate starting rate.

Paediatric patients should be monitored and the dose titrated to the depth of analgesia appropriate for the surgical procedure.

Maintenance of anaesthesia

The following doses of remifentanyl are recommended for maintenance of anaesthesia:

DOSING GUIDELINES FOR PAEDIATRIC PATIENTS (1 to 12 years of age)

*CONCOMITANT ANAESTHETIC AGENT	Remifentanyl BOLUS INJECTION (micrograms /kg)	CONTINUOUS INFUSION (micrograms/kg/min)	
		Starting Rate	Range
Halothane (starting dose 0.3 MAC)	1	0.25	0.05 to 1.3
Sevoflurane (starting dose 0.3 MAC)	1	0.25	0.05 to 0.9
Isoflurane (starting dose 0.5 MAC)	1	0.25	0.06 to 0.9

*co-administered with nitrous oxide/oxygen in a ratio of 2:1

Concomitant medication

At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane, halothane and sevoflurane should be administered as recommended above to avoid an increase of haemodynamic effects (hypotension and bradycardia) of remifentanyl. No data are available for dosage recommendations for simultaneous use of other hypnotics with remifentanyl (see in section above: Administration by Manually Controlled Infusion (MCI), Concomitant medication).

Guidelines for patient management in the immediate post-operative period

Establishment of alternative analgesia prior to discontinuation of Remifentanyl:

Due to the very rapid offset of action of remifentanyl, no residual activity will be present within 5 to 10 minutes after discontinuation. For those patients undergoing surgical procedures where post-operative pain is anticipated, analgesics should be administered prior to discontinuation of remifentanyl. Sufficient time must be allowed to reach the therapeutic effect of the longer acting analgesic. The choice of agent(s), the dose and the time of administration should be planned in advance and individually tailored to be appropriate for the patient's surgical procedure and the level of post-operative care anticipated (see section 4.4).

Neonates/infants (aged less than 1 year):

There is limited clinical trial experience of remifentanyl in neonates and infants (aged under 1 year old; see section 5.1). The pharmacokinetic profile of remifentanyl in neonates/infants (aged less than 1 year) is comparable to that seen in adults after correction for body weight differences (see section 5.2.). However, because there are insufficient clinical data, the administration of remifentanyl is not recommended for this age group.

Use for Total Intravenous anaesthesia (TIVA):

There is limited clinical trial experience of remifentanyl for TIVA in infants (see section 5.1). However, there are insufficient clinical data to make dosage recommendations.

Special Patient groups

For dosage recommendations for special patient groups (elderly and obese patients, renally and hepatically impaired patients, patients undergoing neurosurgery and ASA III/IV patients; see section below).

Cardiac Surgery

Administration by Manually-Controlled Infusion

For dosage recommendations in patients undergoing cardiac surgery see table below:

DOSING GUIDELINES FOR CARDIAC ANAESTHESIA

INDICATION	Remifentanil BOLUS INJECTION (micrograms /kg)	Remifentanil CONTINUOUS INFUSION (micrograms /kg/min)	
		Starting Rate	Typical infusion rates
Intubation	Not recommended	1	–
Maintenance of anaesthesia			
• Isoflurane (starting dose 0.4 MAC)	0.5 to 1	1	0.003 to 4
• Propofol (Starting dose 50 micrograms /kg/min)	0.5 to 1	1	0.01 to 4.3
Continuation of post-operative analgesia, prior to extubation	Not recommended	1	0 to 1

Induction of anaesthesia

After administration of hypnotic to achieve loss of consciousness, Remifentanil should be administered at an initial infusion rate of 1 micrograms/kg/min. The use of bolus injections of Remifentanil during induction in cardiac surgical patients is not recommended. Endotracheal intubation should not occur until at least 5 minutes after the start of the infusion.

Maintenance period of anaesthesia

After endotracheal intubation the infusion rate of Remifentanil should be titrated according to the patient need. Supplemental bolus doses may also be administered as required. High risk cardiac patients, such as those undergoing valve surgery or with poor left ventricular function, should be administered a maximum bolus dose of 0.5 micrograms/kg.

These dosing recommendations also apply during hypothermic cardiopulmonary bypass (see section 5.2).

Concomitant medication

At the doses recommended above, remifentanil significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended above to avoid an increase of haemodynamic effects (hypotension and bradycardia) of remifentanil. No data are available for dosage recommendations for simultaneous use of other hypnotics with remifentanil (see in section above: Administration by Manually Controlled Infusion (MCI), Concomitant medication).

Guidelines for post-operative supply of patient

Continuation of Remifentanil post-operatively to provide analgesia prior to extubation

It is recommended that the infusion of Remifentanil is maintained at the final intra-operative rate during transfer of patients to the post-operative care area. The patient's level of analgesia and sedation should be closely monitored and the Remifentanil infusion rate adjusted to meet the individual patient's requirements (see Intensive care, below, for further information on management of intensive care patients).

Establishment of alternative analgesia prior to discontinuation of Remifentanyl

Due to the very rapid offset of action of remifentanyl, no residual opioid activity will be present within 5 to 10 minutes after discontinuation. Prior to discontinuation of Remifentanyl, patients must be given alternative analgesic and sedative agents at a sufficient time in advance to allow the therapeutic effects of these agents to become established. It is therefore recommended that the choice of agent(s), the dose and the time of administration are planned, before weaning the patient from the ventilator.

Guidelines for discontinuation of Remifentanyl

Due to the very rapid offset of action of Remifentanyl, hypertension, shivering and aches have been reported in cardiac patients immediately following discontinuation of Remifentanyl (see section 4.8). To minimise the risk of these events, adequate alternative analgesia must be established (as described above), before the Remifentanyl infusion is discontinued. The infusion rate should be reduced by 25% decrements in at least 10-minute intervals until the infusion is discontinued.

During weaning from the ventilator the Remifentanyl infusion should not be increased and only down titration should occur, supplemented as required with alternative analgesics. Haemodynamic changes such as hypertension and tachycardia should be treated with alternative agents as appropriate.

When other opioid agents are administered as part of the regimen for transition to alternative analgesia, the patient must be carefully monitored. The benefit of providing adequate post-operative analgesia must always be balanced against the potential risk of respiratory depression with these agents.

Administration by Target-Controlled Infusion

Induction and maintenance of anaesthesia in ventilated patients

Remifentanyl TCI should be used in association with an intravenous or inhalational hypnotic agent during the induction and maintenance of anaesthesia in ventilated adult patients (see the table under *Cardiac surgery/Administration by Manually-Controlled Infusion/Dosing guidelines for cardiac anaesthesia*). In association with these agents, adequate analgesia for cardiac surgery is generally achieved at the higher end of the range of target blood remifentanyl concentrations used for general surgical procedures. Following titration of remifentanyl to individual patient response, blood concentrations as high as 20 nanograms/ml have been achieved in clinical studies.

At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended above to avoid an increase of haemodynamic effects (hypotension and bradycardia) of remifentanyl (see table under *Cardiac surgery/Administration by Manually-Controlled Infusion/Dosing guidelines for cardiac anaesthesia*).

For information on blood remifentanyl concentrations achieved with manually controlled infusion see Table 6

Guidelines for discontinuation/continuation into the immediate post-operative period

At the end of surgery when the TCI infusion is stopped or the target concentration reduced, spontaneous respiration is likely to return at calculated remifentanyl concentrations in the range of 1 to 2 nanograms/ml. As with manually-controlled infusion, post-operative analgesia should be established before the end of surgery with longer acting analgesics (see under *Cardiac surgery/Administration by Manually-Controlled Infusion/Guidelines for discontinuation of Remifentanyl*).

As there are insufficient data, the administration of Remifentanyl by TCI for the management of post-operative analgesia is not recommended.

Intensive Care – Adults

Remifentanyl can be used in mechanically ventilated intensive care patients. If required, additionally sedating drugs should be applied.

Remifentanyl has been studied in intensive care patients in well controlled clinical trials for up to three days. As patients were not studied beyond three days, no evidence of safety and efficacy for longer treatment has been established. Therefore, a usage longer than three days is not recommended.

Due to the lack of data the administration of remifentanyl by TCI is not recommended for ICU patients.

In adults, it is recommended that Remifentanyl is initiated at an infusion rate of 0.1 micrograms /kg/min (6 micrograms /kg/h) to 0.15 micrograms/kg/min (9 micrograms/kg/h). The infusion rate should be titrated in

increments of 0.025 micrograms/kg/min (1.5 micrograms/kg/h) to achieve the desired level of analgesia. A period of at least 5 minutes should be allowed between dose adjustments. The patient should be carefully monitored, regularly reassessed and the Remifentanyl infusion rate adjusted accordingly. If an infusion rate of 0.2 micrograms/kg/min (12 micrograms/kg/h) is reached and the desired level of sedation is not achieved, it is recommended that dosing with an appropriate sedative agent is initiated (see below). The dose of the sedative agent should be titrated to obtain the desired level of sedation. Further increases to the Remifentanyl infusion rate in increments of 0.025 micrograms/kg/min (1.5 micrograms/kg/h) may be made if additional analgesia is required.

The following table summarises the starting infusion rates and typical dose range for provision of analgesia in individual patients:

DOSING GUIDELINES FOR USE OF REMIFENTANIL WITHIN THE INTENSIVE CARE SETTING

CONTINUOUS INFUSION micrograms/kg/min (micrograms/kg/h)	
Starting Rate	Range
0.1 (6) to 0.15 (9)	0.006 (0.36) to 0.74 (44.4)

Bolus doses of Remifentanyl are not recommended in the intensive care setting.

The use of Remifentanyl will reduce the dosage requirement of any concomitant sedative agents. Typical starting doses for sedative agents, if required, are given below.

RECOMMENDED STARTING DOSE OF SEDATIVE AGENTS, IF REQUIRED

Sedative Agent	Bolus (mg/kg)	Infusion rate (mg/kg/h)
Propofol	Up to 0.5	0.5
Midazolam	Up to 0.03	0.03

To allow separate titration of the respective agents, sedative agents should not be administered as an admixture.

Additional analgesia for ventilated patients undergoing painful procedures

An increase in the existing Remifentanyl infusion rate may be required to provide additional analgesic cover for ventilated patients undergoing stimulating and/or painful procedures such as endotracheal suctioning, wound dressing and physiotherapy. It is recommended that a Remifentanyl infusion rate of at least 0.1 micrograms/kg/min (6 micrograms/kg/h) is maintained for at least 5 minutes prior to the start of the stimulating procedure. Further dose adjustments may be made every 2 to 5 minutes in increments of 25%-50% in anticipation of, or in response to, additional requirement for analgesia. A mean infusion rate of 0.25 micrograms/kg/min (15 micrograms/kg/h), maximum 0.75 micrograms/kg/min (44.4 micrograms/kg/h), has been administered for provision of additional analgesia during stimulating procedures.

Establishment of alternative analgesia prior to discontinuation of Remifentanyl

Due to the very rapid offset of action of remifentanyl, no residual opioid activity will be present within 5 to 10 minutes after discontinuation regardless of the duration of infusion. After administration of remifentanyl the potential for the development of tolerance and hyperalgesia should be attended. Therefore, prior to discontinuation of remifentanyl, patients must be given alternative analgesic and sedative agents at a sufficient time in advance to allow the therapeutic effects of these agents to become established and to prevent hyperalgesia and concomitant haemodynamic changes. It is therefore recommended that the choice of agent(s), the dose and the time of administration are planned prior to discontinuation of remifentanyl. Long acting analgesics or intravenous or local analgesics, which can be controlled by the health care staff or the patient are alternative options for analgesia and should be chosen carefully according to the patients needs.

Prolonged administration of μ -opioid agonists may induce development of tolerance.

Guidelines for extubation and discontinuation of Remifentanyl

In order to ensure a smooth emergence from a remifentanyl-based regimen it is recommended that the infusion rate of Remifentanyl is titrated gradually to 0.1 micrograms/kg/min (6 micrograms/kg/h) over a period up to 1 hour prior to extubation.

Following extubation, the infusion rate should be reduced by 25% decrements in at least 10-minute intervals until the infusion is discontinued. During weaning from the ventilator the Remifentanyl infusion should not be increased and only down titration should occur, supplemented as required with alternative analgesics.

Upon discontinuation of Remifentanyl, the IV cannula should be cleared or removed to prevent subsequent inadvertent administration.

When other opioid agents are administered as part of the regimen for transition to alternative analgesia, the patient must be carefully monitored. The benefit of providing adequate analgesia must always be balanced against the potential risk of respiratory depression.

The following tables 1-5 give guidelines for infusion rates of Remifentanyl for manually-controlled infusion:

Table 1: Remifentanyl – Infusion rates (ml/kg/h)

Medicinal product delivery rate (micrograms /kg/min)	Infusion delivery rate (ml/kg/h) for solution concentrations of			
	20 micrograms/ml	25 micrograms/ml	50 micrograms/ml	250 micrograms/ml
	1 mg/50 ml	1 mg/40 ml	1 mg/20 ml	10 mg/40 ml
0.0125	0.038	0.03	0.015	Not recommended
0.025	0.075	0.06	0.03	Not recommended
0.05	0.15	0.12	0.06	0.012
0.075	0.23	0.18	0.09	0.018
0.1	0.3	0.24	0.12	0.024
0.15	0.45	0.36	0.18	0.036
0.2	0.6	0.48	0.24	0.048
0.25	0.75	0.6	0.3	0.06
0.5	1.5	1.2	0.6	0.12
0.75	2.25	1.8	0.9	0.18
1.0	3.0	2.4	1.2	0.24
1.25	3.75	3.0	1.5	0.3
1.5	4.5	3.6	1.8	0.36
1.75	5.25	4.2	2.1	0.42
2.0	6.0	4.8	2.4	0.48

Table 2: Remifentanyl – Infusion rates (ml/h) for a 20 micrograms /ml solution

Infusion rate (micrograms /kg/min)	Patient weight (kg)						
	5	10	20	30	40	50	60
0.0125	0.188	0.375	0.75	1.125	1.5	1.875	2.25
0.025	0.375	0.75	1.5	2.25	3.0	3.75	4.5
0.05	0.75	1.5	3.0	4.5	6.0	7.5	9.0
0.075	1.125	2.25	4.5	6.75	9.0	11.25	13.5
0.1	1.5	3.0	6.0	9.0	12.0	15.0	18.0
0.15	2.25	4.5	9.0	13.5	18.0	22.5	27.0
0.2	3.0	6.0	12.0	18.0	24.0	30.0	36.0
0.25	3.75	7.5	15.0	22.5	30.0	37.5	45.0
0.3	4.5	9.0	18.0	27.0	36.0	45.0	54.0
0.35	5.25	10.5	21.0	31.5	42.0	52.5	63.0
0.4	6.0	12.0	24.0	36.0	48.0	60.0	72.0

Table 3: Remifentanil – Infusion rates (ml/h) for a 25 micrograms /ml solution

Infusion rate (micrograms /kg/min)	Patient weight (kg)									
	10	20	30	40	50	60	70	80	90	100
0.0125	0.3	0.6	0.9	1.2	1.5	1.8	2.1	2.4	2.7	3.0
0.025	0.6	1.2	1.8	2.4	3.0	3.6	4.2	4.8	5.4	6.0
0.05	1.2	2.4	3.6	4.8	6.0	7.2	8.4	9.6	10.8	12.0
0.075	1.8	3.6	5.4	7.2	9.0	10.8	12.6	14.4	16.2	18.0
0.1	2.4	4.8	7.2	9.6	12.0	14.4	16.8	19.2	21.6	24.0
0.15	3.6	7.2	10.8	14.4	18.0	21.6	25.2	28.8	32.4	36.0
0.2	4.8	9.6	14.4	19.2	24.0	28.8	33.6	38.4	43.2	48.0

Table 4: Remifentanil – Infusion rates (ml/h) for a 50 micrograms ml solution

Infusion rate (micrograms /kg/min)	Patient weight (kg)							
	30	40	50	60	70	80	90	100
0.025	0.9	1.2	1.5	1.8	2.1	2.4	2.7	3.0
0.05	1.8	2.4	3.0	3.6	4.2	4.8	5.4	6.0
0.075	2.7	3.6	4.5	5.4	6.3	7.2	8.1	9.0
0.1	3.6	4.8	6.0	7.2	8.4	9.6	10.8	12.0
0.15	5.4	7.2	9.0	10.8	12.6	14.4	16.2	18.0
0.2	7.2	9.6	12.0	14.4	16.8	19.2	21.6	24.0
0.25	9.0	12.0	15.0	18.0	21.0	24.0	27.0	30.0
0.5	18.0	24.0	30.0	36.0	42.0	48.0	54.0	60.0
0.75	27.0	36.0	45.0	54.0	63.0	72.0	81.0	90.0
1.0	36.0	48.0	60.0	72.0	84.0	96.0	108.0	120.0
1.,25	45.0	60.0	75.0	90.0	105.0	120.0	135.0	150.0
1.5	54.0	72.0	90.0	108.0	126.0	144.0	162.0	180.0
1.75	63.0	84.0	105.0	126.0	147.0	168.0	189.0	210.0
2.0	72.0	96.0	120.0	144.0	168.0	192.0	216.0	240.0

Table 5: Remifentanil – Infusion rates (ml/h) for a 250 micrograms /ml solution

Infusion rate (micrograms/kg/min)	Patient weight (kg)							
	30	40	50	60	70	80	90	100
0.1	0.72	0.96	1.20	1.44	1.68	1.92	2.16	2.40
0.15	1.08	1.44	1.80	2.16	2.52	2.88	3.24	3.60
0.2	1.44	1.92	2.40	2.88	3.36	3.84	4.32	4.80
0.25	1.80	2.40	3.00	3.60	4.20	4.80	5.40	6.00
0.5	3.60	4.80	6.00	7.20	8.40	9.60	10.80	12.00
0.75	5.40	7.20	9.00	10.80	12.60	14.40	16.20	18.00
1.0	7.20	9.60	12.00	14.40	16.80	19.20	21.60	24.00
1.25	9.00	12.00	15.00	18.00	21.00	24.00	27.00	30.00
1.5	10.80	14.40	18.00	21.60	25.20	28.80	32.40	36.00
1.75	12.60	16.80	21.00	25.20	29.40	33.60	37.80	42.00
2.0	14.40	19.20	24.00	28.80	33.60	38.40	43.20	48.00

The following table provides the equivalent blood remifentanil concentration using a TCI approach for various manually-controlled infusion rates at steady state:

Table 6: Remifentanil blood concentrations (nanograms/ml) estimated using the Minto (1997) pharmacokinetic model in a 70 kg, 170 cm, 40 year old male patient for various manually-controlled infusion rates (micrograms/kg/min) at Steady State.

Remifentanil-Infusion rates (micrograms /kg/min)	Remifentanil-blood concentration (nanograms/ml)
0.05	1.3
0.10	2.6
0.25	6.3

0.40	10.4
0.50	12.6
1.0	25.2
2.0	50.5

Intensive Care - Paediatric patients

The use of remifentanyl in paediatric intensive care patients cannot be recommended as there are no data available in this patient population.

Renally-impaired intensive care patients

No adjustments to the doses recommended above are necessary in renally-impaired patients, including those undergoing renal replacement therapy. It should, however, be considered that in patients with impaired renal function the clearance of the carboxylic acid metabolite is reduced (see section 5.2).

Special patient groups

Elderly (over 65 years of age)

General anaesthesia

Caution should be exercised in the administration of remifentanyl in this population. The initial starting dose of Remifentanyl administered to patients over 65 should be half the recommended adult dose and then titrated to the individual patient’s need as an increased sensitivity to the pharmacodynamic effects of remifentanyl has been seen in this patient population.

This dose adjustment refers to the use in all phases of the anaesthesia including onset, maintaining and immediate post operative analgesia.

Because of the increased sensitivity of elderly patients to remifentanyl, when administering remifentanyl by TCI in this population the initial target concentration should be 1.5 to 4 nanograms/ml with subsequent titration according to the individual patient’s response.

Cardiac anaesthesia

Reduction of initial dosage is not required.

Intensive care

Reduction of initial dosage is not required.

Neurosurgery

Limited clinical experience in patients undergoing neurosurgery has shown that no special dosage recommendations are required.

ASA III/IV patients

General anaesthesia

As the haemodynamic effects of potent opioids can be expected to be more pronounced in ASA III/IV patients, caution should be exercised in the administration of remifentanyl in this population. Initial dosage reduction and subsequent titration to effect is therefore recommended.

As there are insufficient data, dosage recommendation cannot be given for children.

For TCI, a lower initial target of 1.5 to 4 nanograms/ml should be used in ASA III or IV patients and subsequently titrated to response.

Cardiac anaesthesia

No initial dose reduction is required.

Obese patients

For manually-controlled infusion it is recommended that for obese patients the dosage of Remifentanyl should be reduced and based upon ideal body weight as the clearance and volume of distribution of remifentanyl are better correlated with ideal body weight than actual body weight.

With the calculation of lean body mass (LBM) used in the Minto model, LBM is likely to be underestimated in female patients with a body mass index (BMI) greater than 35 kg/m² and in male patients with BMI greater than 40 kg/m². To avoid underdosing in these patients, remifentanyl TCI should be titrated carefully to individual response.

Renal impairment

On the basis of investigations carried out to date, a dose adjustment in patients with impaired renal function, including intensive care patients, is not necessary; however, these patients exhibit reduced clearance of carboxylic acid metabolite.

Patients with hepatic impairment

No adjustment of the initial dose, relative to that used in healthy adults, is necessary as the pharmacokinetic profile of remifentanyl is unchanged in this patient population. However, patients with severe hepatic impairment may be slightly more sensitive to the respiratory depressant effects of remifentanyl (see section 4.4). These patients should be closely monitored and the dose of remifentanyl titrated to individual patient need.

4.3 Contraindications

As glycine is present in the formulation, Remifentanyl is contra-indicated for epidural and intrathecal use (see section 5.3).

Remifentanyl is contra-indicated in patients with known hypersensitivity to remifentanyl and other fentanyl analogues or any other component of the preparation (see section 6.1).

Remifentanyl is contra-indicated for use as the sole agent for induction of anaesthesia.

4.4 Special warnings and precautions for use

Remifentanyl should be administered only in a setting fully equipped for the monitoring and support of respiratory and cardiovascular function, and by persons specifically trained in the use of anaesthetic drugs and the recognition and management of the expected adverse effects of potent opioids, including respiratory and cardiac resuscitation. Such training must include the establishment and maintenance of a patent airway and assisted ventilation.

As mechanically ventilated, intensive care patients were not studied beyond three days, no evidence of safety and efficacy for longer treatment has been established. Therefore, a longer usage is not recommended in intensive care patients.

Rapid offset of action/Switch to alternative analgesic treatment

Due to the very rapid offset of action of remifentanyl, patients may emerge rapidly from anaesthesia and no residual opioid activity will be present within 5–10 minutes after the discontinuation of remifentanyl. During administration of remifentanyl as a μ -opioid agonist the potential for the development of tolerance and hyperalgesia should be paid attention to. Therefore, prior to discontinuation of remifentanyl, patients must be given alternative analgesic and sedative agents at a sufficient time in advance to allow the therapeutic effects of these agents to become established and to prevent hyperalgesia and concomitant haemodynamic changes.

For those patients undergoing surgical procedures where post-operative pain is anticipated, analgesics should be administered prior to discontinuation of remifentanyl. Sufficient time must be allowed to reach the maximum effect of the longer acting analgesic. The choice of analgesic should be appropriate for the patient's surgical procedure and the level of post-operative care. When other opioid agents are administered as part of the regimen for transition to alternative analgesia, the benefit of providing adequate post-operative analgesia must always be balanced against the potential risk of respiratory depression with these agents.

Inadvertent administration

A sufficient amount of Remifentanyl may be present in the dead space of the IV line and/or cannula to cause respiratory

depression, apnoea and/or muscle rigidity if the line is flushed with IV fluids or other drugs. This may be avoided by administering Remifentanyl into a fast flowing IV line or via a dedicated IV line which is removed when Remifentanyl is discontinued.

Discontinuation of treatment

Symptoms following withdrawal of remifentanyl including tachycardia, hypertension and agitation have been reported infrequently upon abrupt cessation, particularly after prolonged administration of more than 3 days. Where reported, re-introduction and tapering of the infusion has been beneficial. The use of Remifentanyl in mechanically ventilated intensive care patients is not recommended for duration of treatment greater than 3 days.

Muscle rigidity - prevention and management

At the doses recommended muscle rigidity may occur. As with other opioids, the incidence of muscle rigidity is related to the dose and rate of administration. Therefore, bolus injections should be administered over not less than 30 seconds.

Muscle rigidity induced by remifentanyl must be treated in the context of the patient's clinical condition with appropriate supporting measures including ventilatory support. Excessive muscle rigidity occurring during the induction of anaesthesia should be treated by the administration of a neuromuscular blocking agent and/or additional hypnotic agents. Muscle rigidity seen during the use of remifentanyl as an analgesic may be treated by stopping or decreasing the rate of administration of remifentanyl. Resolution of muscle rigidity after discontinuing the infusion of remifentanyl occurs within minutes. Alternatively an opioid antagonist may be administered; however this may reverse or attenuate the analgesic effect of remifentanyl.

Respiratory depression – preventive measures and treatment

As with all potent opioids, profound analgesia is accompanied by marked respiratory depression. Therefore, remifentanyl should only be used in areas where facilities for monitoring and dealing with respiratory depression are available. Special care should be taken in patients with impaired lung function and with severe hepatic impairment. These patients may be slightly more sensitive to the respiratory depressant effects of remifentanyl. These patients should be closely monitored and the dose of remifentanyl titrated to individual patient need.

The appearance of respiratory depression should be managed appropriately, including decreasing the rate of infusion by 50 %, or by a temporary discontinuation of the infusion. Unlike other fentanyl analogues, remifentanyl has not been shown to cause recurrent respiratory depression even after prolonged administration. However in the presence of confounding factors (e.g. inadvertent administration of bolus doses (see section below) and administration of concomitant longer acting opioids), respiratory depression occurring up to 50 minutes after discontinuation of infusion has been reported. As many factors may affect post-operative recovery, it is important to ensure that full consciousness and adequate spontaneous ventilation are achieved before the patient is discharged from the recovery area.

Cardiovascular effects

Hypotension and bradycardia can give rise to asystole and cardiac arrest (see sections 4.5 and 4.8) may be managed by reducing the rate of infusion of remifentanyl or the dose of concurrent anaesthetics or by using IV fluids, vasopressor or anticholinergic agents as appropriate.

Debilited, hypovolaemic, hypotensive and elderly patients may be more sensitive to the cardiovascular effects of remifentanyl.

Neonates/infants

There is limited data available on use in neonates/infants under 1 year of age (see sections 4.2 and 5.1).

Drug abuse

As with other opioids remifentanyl may produce dependency.

This medicinal product contains less than 1 mmol sodium (23 mg) per ml.

4.5 Interaction with other medicinal products and other forms of interaction

Remifentanyl is not metabolised by plasmacholinesterase, therefore, interactions with medicinal products metabolized by this enzyme are not anticipated.

As with other opioids remifentanyl, whether given by manually-controlled infusion or TCI, decreases the amounts or

doses of inhaled and IV anaesthetics, and benzodiazepines required for anaesthesia (see section 4.2). If doses of concomitantly administered CNS depressant medicinal products are not reduced patients may experience an increased incidence of adverse effects associated with these agents.

Information of drug interactions with other opioids in relation to anaesthesia is very limited.

The cardiovascular effects of remifentanyl (hypotension and bradycardia), may exacerbate in patients receiving concomitant cardiac depressant drugs, such as beta-blockers and calcium channel blocking agents (see also sections 4.4 and 4.8).

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate and well-controlled studies in pregnant women should Remifentanyl be used during pregnancy only if the potential benefit justifies the potential risk to the foetus.

Breast-feeding

It is not known whether remifentanyl is excreted in human milk. However, because fentanyl analogues are excreted in human milk and remifentanyl-related material was found in rat milk after dosing with remifentanyl, nursing mothers should be advised to discontinue breast- feeding for 24 hours following administration of remifentanyl.

Labour and delivery

There are insufficient data to recommend remifentanyl for use during labour and caesarean section. It is known that remifentanyl crosses the placental barrier and fentanyl analogues can cause respiratory depression in the child.

4.7 Effects on ability to drive and use machines

Remifentanyl has major influence on the ability to drive and use machines.

If an early discharge is envisaged after application of remifentanyl, following treatment using anaesthetic agents, patients should be advised not to drive or operate machinery. It is advisable that the patient is accompanied when returning home and that alcoholic drink is avoided. The physician should decide when these activities may be resumed.

4.8 Undesirable effects

The most common adverse events associated with remifentanyl are direct extensions of μ -opioid agonist pharmacology. These adverse events resolve within minutes of discontinuing or decreasing the rate of remifentanyl administration.

The following terminologies have been used in order to classify the occurrence of undesirable effects:

Very common	$\geq 1/10$		
Common	$\geq 1/100$	to	$< 1/10$
Uncommon	$\geq 1/1,000$	to	$< 1/100$
Rare	$\geq 1/10,000$	to	$< 1/1,000$
Very rare	$< 1/10,000$		
Not known (cannot be estimated from the available data)			

Incidence is listed below within each body system:

Immune system disorders	Rare	hypersensitivity reactions including anaphylaxis have been reported in patients receiving remifentanyl in conjunction with one or more anaesthetic agents
Psychiatric disorders	Not known	dependence
Nervous system disorders	Very common	skeletal muscle rigidity
	Rare	sedation (during awakening after general anaesthesia)
	Not known	convulsions
Cardiac disorders	Common	bradycardia
	Rare	asystole/cardiac arrest with preceding bradycardia in patients treated with remifentanyl in combination with other anaesthetics
	Not known	atrioventricular block
Vascular disorders	Very common	hypotension
	Common	post-operatively occurring hypertension
Respiratory, thoracic and mediastinal disorders	Common	acute respiratory depression, apnoea
	Uncommon	hypoxia
Gastrointestinal disorders	Very common	nausea, vomiting
	Uncommon	constipation
Skin and subcutaneous tissue disorders	Common	Pruritus
General disorders and administration site conditions	Common	post-operative shivering
	Uncommon	post-operative pain
	Not known	drug tolerance

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

4.9 Overdose

As with all potent opioid analgesics, overdose would be manifested by an extension of the pharmacologically

predictable actions of remifentanil. Due to the very short duration of action of remifentanil, the potential for deleterious effects due to overdose is limited to the immediate time period following medicinal product administration. Response to discontinuation of the medicinal product is rapid, with return to baseline within ten minutes.

In the event of overdose, or suspected overdose, take the following actions: discontinue administration of Remifentanil , maintain a patent airway, initiate assisted or controlled ventilation with oxygen, and maintain adequate cardiovascular function. If depressed respiration is associated with muscle rigidity, a neuromuscular blocking agent may be required to facilitate assisted or controlled respiration. Intravenous fluids and vasopressor agents for the treatment of hypotension and other supportive measures may be employed.

Intravenous administration of an opioid antagonist such as naloxone may be given as a specific antidote in addition to ventilatory support to manage severe respiratory depression. The duration of respiratory depression following overdose with Remifentanil is unlikely to exceed the duration of action of the opioid antagonist.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Opioid anaesthetics, ATC code: N01AH06

Remifentanil is a selective μ -opioid agonist with a rapid onset and very short duration of action. The μ -opioid activity of remifentanil is antagonized by narcotic antagonists, such as naloxone.

Assays of histamine in patients and normal volunteers have shown no elevation in histamine levels after administration of remifentanil in bolus doses up to 30 micrograms/kg.

Neonates/infants (aged less than 1 year):

In a randomised (ratio of 2:1, remifentanil:halothane), open label, parallel group, multicentre study in 60 young infants and neonates $\leq$ 8 weeks of age (mean 5.5 weeks) with an ASA physical status of I-II who were undergoing pyloromyotomy, the efficacy and safety of remifentanil (given as a 0.4 micrograms/kg/min initial continuous infusion plus supplemental doses or infusion rate changes as needed) was compared with halothane (given at 0.4 % with supplemental increases as needed). Maintenance of anaesthesia was achieved by the additional administration of 70 % nitrous oxide (N2O) plus 30 % oxygen. Recovery times were superior in the remifentanil relative to the halothane groups (not significant).

Use for Total Intravenous anaesthesia (TIVA) – children aged 6 months to 16 years:

TIVA with remifentanil in paediatric surgery was compared to inhalation anaesthesia in three randomised, open-label studies. The results are summarised in the table below.

Surgical intervention	Age (y), (N)	Study condition (maintenance)	Extubation (min) (mean (SD))
Lower abdominal/urological surgery	0.5-16 (120)	TIVA: propofol (5-10 mg/kg/h) + remifentanil (0.125 – 1.0 micrograms/kg/min)	11.8 (4.2)
		Inhalation anaesthesia: sevoflurane (1.0 – 1.5 MAC) and remifentanil (0.125 – 1.0 micrograms/kg/min)	15.0(5.6) (p<0.05)
ENT-surgery	4-11 (50)	TIVA: propofol (3 mg/kg/h) + remifentanil (0.5 micrograms /kg/min)	11 (3.7)
		Inhalation anaesthesia:	9.4 (2.9)

		desflurane (1.3 MAC) and N ₂ O mixture	Not significant
General or ENT surgery	2-12 (153)	TIVA: remifentanyl (0.2 – 0.5 micrograms/kg/min) + propofol (100 – 200 micrograms/kg/min)	Comparable extubation times (based on limited data)
		Inhalation anaesthesia: sevoflurane (1-1.5 MAC) + N ₂ O mixture	

In the study in lower abdominal/urological surgery comparing remifentanyl/propofol with remifentanyl/sevoflurane, hypotension occurred significantly more often under remifentanyl/sevoflurane, and bradycardia occurred significantly more often under remifentanyl/propofol. In the study in ENT surgery comparing remifentanyl/propofol with desflurane/nitrous oxide, a significantly higher heart rate was seen in subjects receiving desflurane/nitrous oxide compared with remifentanyl/propofol and with baseline values.

5.2 Pharmacokinetic properties

Following administration of the recommended doses of remifentanyl, the effective biological half-life is 3-10 minutes. The average clearance of remifentanyl in young healthy adults is 40 ml/min/kg, the central volume of distribution is 100 ml/kg and the steady-state volume of distribution is 350 ml/kg. Blood concentrations of remifentanyl are proportional to the dose administered throughout the recommended dose range. For every 0.1 micrograms/kg/min increase in i.v. infusion rate, the blood concentration of remifentanyl will rise to 2.5 nanograms/ml. Remifentanyl is approximately 70 % bound to plasma proteins.

Metabolism

Remifentanyl is an esterase metabolised opioid that is susceptible to metabolism by non-specific blood and tissue esterases. The metabolism of remifentanyl results in the formation of an essentially inactive carboxylic acid metabolite (1/4600th as potent as remifentanyl). Studies in man indicate that all pharmacological activity is associated with the parent compound. The activity of this metabolite is therefore not of any clinical consequence. The half life of the metabolite in healthy adults is 2 hours. Approximately 95 % of remifentanyl as the carboxylic acid metabolite is recovered in the urine in patients with normal renal function. Remifentanyl is not a substrate for plasma cholinesterase.

Placental barrier and breast milk

In a human clinical trial, the average maternal remifentanyl concentrations were approximately twice those seen in the foetus. In some cases, however, foetal concentrations were similar to those in the mother. The umbilical arteriovenous ratio of remifentanyl concentrations was approximately 30 % suggesting metabolism of remifentanyl in the neonate. Remifentanyl related material is transferred to the milk of lactating rats.

Cardiac anaesthesia

The clearance of remifentanyl is reduced by approximately 20% during hypothermic (28 °C) cardiopulmonary bypass. A decrease in body temperature lowers elimination clearance by 3% per degree centigrade.

Renal impairment

The rapid recovery from remifentanyl-based sedation and analgesia is unaffected by renal status. The pharmacokinetics of remifentanyl are not significantly changed in patients with varying degrees of renal impairment even after administration for up to 3 days in the intensive care setting. There is no evidence that remifentanyl is extracted during renal replacement therapy. The clearance of the carboxylic acid metabolite is reduced in patients with renal impairment. In intensive care patients with moderate/severe renal impairment, the concentration of the carboxylic acid metabolite is expected to reach approximately 100-fold the level of remifentanyl at steady-state. Clinical data demonstrate that the accumulation of the

metabolite does not result in clinically relevant μ -opioid effects even after administration of remifentanyl infusions for up to 3 days in these patients.

Up to now, data on safety and pharmacokinetic activity of metabolites after infusion of remifentanyl for more than 3 days are lacking.

The carboxylic acid metabolite is extracted during haemodialysis by 25 – 35 %. In patients with anuria the half-life of the carboxylic acid metabolite is increased to 30 hours.

Hepatic impairment

The pharmacokinetic profile of remifentanyl is not changed in patients with severe hepatic impairment awaiting liver transplant, or during the anhepatic phase of liver transplant surgery. Patients with severe hepatic impairment may be slightly more sensitive to the respiratory depressant effects of remifentanyl. These patients should be closely monitored and the dose of remifentanyl should be titrated to the individual patient need.

Paediatric patients

The average clearance and steady state volume of distribution of remifentanyl are increased in younger children and decline to young healthy adult values by age 17. The elimination half-life of remifentanyl in neonates is not significantly different from that of young healthy adults. Changes in analgesic effect after changes in infusion rate of remifentanyl should be rapid and similar to those seen in young healthy adults. The pharmacokinetics of the carboxylic acid metabolite in paediatric patients 2-17 years of age are similar to those seen in adults after correcting for differences in body weight.

Elderly

The clearance of remifentanyl is slightly reduced (approximately 25 %) in elderly patients (over 65 years of age) compared to that in young patients. The pharmacodynamic activity of remifentanyl increases with increasing age. Elderly patients have a remifentanyl EC_{50} for formation of delta waves on the electroencephalogram that is 50 % lower than young patients; therefore, the initial dose of remifentanyl should be reduced by 50 % in elderly patients and then carefully titrated to meet the individual patient need.

5.3 Preclinical safety data

Acute toxicity

Expected signs of μ -opioid intoxication were observed in non-ventilated mice, rats, and dogs after large single bolus intravenous doses of remifentanyl. In these studies, the most sensitive species, the male rat, survived following administration of 5 mg/kg.

Intracranial bleedings in dogs caused by hypoxia declined within 14 days after stopping remifentanyl application.

Chronic toxicity

Bolus doses of remifentanyl administered to non-ventilated rats and dogs resulted in respiratory depression in all dose groups, and in reversible intracranial bleedings in dogs. Subsequent investigations showed that the microhaemorrhages resulted from hypoxia and were not specific to remifentanyl. Brain microhaemorrhages were not observed in infusion studies in non-ventilated rats and dogs because these studies were conducted at doses that did not cause severe respiratory depression. It is to be derived from preclinical studies that respiratory depression and associated sequelae are the most likely cause of potentially serious adverse events in humans.

Intrathecal administration to dogs of the glycine formulation alone (i.e. without remifentanyl) evoked agitation, pain and hind limb dysfunction and incoordination. These effects are believed to be secondary to the glycine excipient. Because of the better buffering properties of blood, the more rapid dilution, and the low glycine concentration of the remifentanyl formulation, this finding has no clinical relevance for intravenous administration of Remifentanyl.

Reproductive toxicity studies

Placental transfer studies in rats and rabbits showed that pups are exposed to remifentanyl and/or its metabolites during growth and development. Remifentanyl-related material is transferred to the milk of lactating rats.

Remifentanyl has been shown to reduce fertility in male rats when administered daily by intravenous injection for at least 70 days at a dose of 0.5 mg/kg, or approximately 250 times the maximum recommended human bolus dose of 2

microgram/kg. The fertility of female rats was not affected at doses up to 1 mg/kg when administered for at least 15 days prior to mating. No teratogenic effects have been observed with remifentanyl at doses up to 5 mg/kg in rats and 0.8 mg/kg in rabbits. Administration of remifentanyl to rats throughout late gestation and lactation at doses up to 5 mg/kg IV had no significant effect on the survival, development, or reproductive performance of the F1 generation.

Genotoxicity

Remifentanyl did not yield positive findings in a series of in vitro and in vivo genotoxicity tests, except in the in vitro mouse lymphoma tk assay, which gave a positive result with metabolic activation. Since the mouse lymphoma results could not be confirmed in further in vitro and in vivo tests, treatment with remifentanyl is not considered to pose a genotoxic hazard to patients.

Carcinogenicity

There were no long-term carcinogenicity studies performed.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Glycine
Hydrochloric acid 37 % (for pH adjustment)
Sodium hydroxide 17 % solution (for pH adjustment)

6.2 Incompatibilities

Remifentanyl must not be mixed with other medicinal products except those mentioned in section 6.6. It should not be admixed with Lactated Ringer's Injection or Lactated Ringer's and glucose 50 mg/ml (5 %) solution for injection. Remifentanyl should not be mixed with propofol in the same intravenous admixture solution. For compatibility when given into a running i.v. catheter, please see section 6.6.

Administration of Remifentanyl into the same intravenous line with blood/serum/plasma is not recommended as non-specific esterase in blood products may lead to the hydrolysis of remifentanyl to its inactive metabolite. Remifentanyl should not be mixed with other therapeutic agents prior to administration.

6.3 Shelf life

24 months

After reconstitution:

After reconstitution, chemical and physical in-use stability has been demonstrated for 24 hours at 25°C.

From a microbiological point of view, unless the method of reconstitution precludes the risk of microbial contamination, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

After dilution:

The diluted product should be used immediately.

6.4 Special precautions for storage

Unopened medicinal product

Do not store above 25 °C.

Do not refrigerate or freeze.

Reconstituted/diluted medicinal product. For storage conditions of the reconstituted and diluted medicinal product, see

section 6.3.

6.5 Nature and contents of container

8 ml vial of colourless glass (type 1) with bromobutyl rubber stopper and blue cap.

Pack sizes: 5 vials per pack

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Remifentanyl should be prepared for intravenous use by adding the appropriate volume (as stated in the table below) of one of the below listed diluents to give a reconstituted solution with a concentration of approximately 1mg/ml.

Presentation	Volume of diluent to be added	Concentration of the reconstituted solution
Remifentanyl 1 mg	1 ml	1 mg/ml
Remifentanyl 2 mg	2 ml	1 mg/ml
Remifentanyl 5 mg	5 ml	1 mg/ml

Following reconstitution, the product should be inspected visually (as far as supported by the vial) for solids, discoloration or damage to the vials. If such changes are detected, the solution must be discarded. The finished solution is for single use only. Unused solution must be discarded.

For manually-controlled infusion, Remifentanyl should be administered following further dilution to a concentration of 20 to 250 micrograms/ml (50 micrograms/ml is the recommended dilution for adults and 20 to 25 micrograms/ml for paediatric patients aged 1 year and over).

For target controlled infusion (TCI), Remifentanyl should be administered following further dilution to a concentration of 20 to 50 micrograms/ml.

Dilution should be adjusted to the technical equipment of the infusion system and the expected patient requirements.

The diluted solution is for single use only.

For dilution, one of the following IV fluids listed below should be used:

- Water for Injections
- Glucose 50 mg/ml (5%) solution for Injection
- Glucose 50 mg/ml (5%) and sodium chloride 9 mg/ml (0.9 %) solution for injection
- Sodium chloride 9 mg/ml (0.9 %) solution for injection
- Sodium chloride 4.5 mg/ml (0.45 %) solution for injection

Following dilution, the solution should be inspected visually to ensure that it is clear, colourless and virtually free from solids, and that there is no damage to the vials. If such changes are detected, the solution must be discarded.

Remifentanyl has been shown to be compatible with the following IV fluids when administered into a running IV catheter:

- Lactated Ringer's solution for injection
- Lactated Ringer's and Glucose 50 mg/ml (5%) solution for injection

Remifentanyl has been shown to be compatible with propofol when administered into a running IV catheter.

Any unused product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Hospira UK Limited
Horizon, Honey Lane,
Hurley,
Maidenhead,
SL6 6RJ
United Kingdom

8 MARKETING AUTHORISATION NUMBER

PA0437/068/003

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 28th October 2011

Date of last renewal: 31st January 2015

10 DATE OF REVISION OF THE TEXT

January 2018