

PACKAGE LEAFLET: INFORMATION FOR THE USER
FLAGYL®-S 200mg/5ml Oral Suspension
Metronidazole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If any side effects get serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist. See section 4.

What is in this leaflet:

1. What Flagyl-S Suspension is and what it is used for
2. What you need to know before you take Flagyl-S Suspension
3. How to take Flagyl-S Suspension
4. Possible side effects
5. How to store Flagyl-S Suspension
6. Contents of the pack and other information

1. WHAT FLAGYL-S SUSPENSION IS AND WHAT IT IS USED FOR

Flagyl-S Suspension belongs to a group of medicines called anti-microbials. Flagyl-S Suspension is used in the treatment and prevention of severe infections that are caused by anaerobic bacteria, particularly species of *Bacteroides*, *anaerobic Streptococci*, *Fusobacteria*, *Clostridia*, etc.

It works by killing bacteria and parasites that cause infections in your body.

It can be used to:

- Treat infections of the blood, brain, lung, bones, genital tract, pelvic area, stomach and intestines.
- Treat gum ulcers and other dental infections.
- Treat infected leg ulcers and pressure sores.
- Prevent infections after surgery.
- In the prevention and treatment of infections due to anaerobic bacteria (bacteria that are able to survive in the absence of oxygen)

If you need any further information on your illness, speak to your doctor.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE FLAGYL-S SUSPENSION

Do not take Flagyl-S Suspension:

- If you are allergic (hypersensitive) to metronidazole or any of the other ingredients of this medicine (listed in Section 6). Signs of an allergic reaction include: rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue.
- If you are pregnant or breast feeding or think you might be pregnant.
- If you are suffering or have ever suffered from any liver or kidney disease.
- If you are suffering or have ever suffered from any diseases of the nervous system.

Warnings and precautions

Cases of severe liver toxicity/acute liver failure, including cases with a fatal outcome, in patients with Cockayne syndrome have been reported with product containing metronidazole. If you are affected by

Cockayne syndrome, your doctor should also monitor your liver function frequently while you are being treated with metronidazole and afterwards.

Tell your doctor immediately and stop taking metronidazole if you develop stomach pain, anorexia, nausea, vomiting, fever, malaise, vertigo (spinning sensation), fatigue, jaundice, dark urine, putty or mastic-coloured stools or itching.

There have been rare reports of potentially life-threatening skin rashes (Stevens-Johnson Syndrome, Toxic Epidermal Necrolysis, Acute Generalised Exanthematous Pustulosis) with the use of Flagyl-S Suspension. Symptoms of these may include: flu-like symptoms, a painful red or purplish rash that spreads and blisters or a rash of pustules which may be accompanied by a fever. You may also have ulcers in your mouth, throat, nose or genitals. If you develop any of these you should stop taking your medicine and inform your doctor straight away (see section 4).

If you experience flu-like symptoms and a rash with a fever and, enlarged lymph nodes and increase in a type of white blood cell (eosinophilia). Other abnormal blood test results may include (but are not limited to) increased levels of liver enzymes. If you develop any of the reactions listed above, stop treatment immediately and contact your doctor or healthcare professional (see Section 4).

Other medicines and Flagyl-S Suspension

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines obtained without a prescription, including herbal medicines. This is because Flagyl-S Suspension can affect the way some medicines work. Also, some other medicines can affect the way Flagyl-S Suspension works.

In particular tell your doctor or pharmacist if you are taking any of the following medicines:

- Medicines used to thin the blood, such as warfarin
- Lithium (used to treat depression)
- Disulfiram (used to treat alcoholism)
- Phenobarbital or Phenytoin (used to treat epilepsy)
- Cyclosporin (used to treat arthritis and some skin conditions)
- 5-flourouracil (used in the treatment of cancer).
- Busulfan (used in the treatment of leukaemia).
- Any medicine that may cause heart rhythm disturbances (so called QT prolongation which can be seen on ECG) such as certain anti-arrhythmics (medicines for heart rhythm disorders), certain antibiotics and medication primarily used to manage psychosis (including delusions, hallucinations, paranoia or disordered thought).

Flagyl-S Suspension with alcohol

Do not drink any alcohol while you are taking Flagyl-S Suspension and for 48 hours after finishing your course. Drinking alcohol while using Flagyl-S Suspension might cause unpleasant side effects, such as feeling sick (nausea), being sick (vomiting), stomach pain, hot flushes, very fast or uneven heart-beat (palpitations) and headache.

Pregnancy and breast-feeding

Tell your doctor before using Flagyl-S Suspension if:

- You are pregnant, might become pregnant or think you may be pregnant. Flagyl-S Suspension should not be taken during pregnancy unless considered absolutely necessary.
- If you are breast-feeding. It is better not to use Flagyl-S Suspension if you are breast-feeding. This is because small amounts may pass into the mother's milk.

Driving and using machines

Whilst taking Flagyl-S Suspension you may feel sleepy, dizzy or experience vertigo (spinning sensation), be confused, see or hear things that are not there (hallucinations), have fits (convulsions) or temporary eyesight problems (such as blurred or double vision). If this happens, do not drive or use any machinery or tools.

Tests

Your doctor may wish to carry out some tests if you have been using this medicine for more than 10 days. If you are going to have a blood test, tell the doctor or nurse performing the test that you are using Flagyl-S Suspension. Flagyl-S Suspension can affect the results of some blood tests.

Flagyl-S Suspension contains

- **Sucrose:** Each dose contains up to 3.01g sucrose. This should be taken into account in patients with diabetes mellitus.
- **Alcohol:** This medicine contains 31.4 mg of alcohol (ethanol) in each 5 ml of suspension which is equivalent to 6.28 mg per ml (0.628% w/v). The amount in 5 ml of this medicine is equivalent to less than 1 ml beer or 1 ml wine.
The small amount of alcohol in this medicine will not have any noticeable effects.
- **Sodium:** This medicine contains less than 1mmol sodium (23mg) per 5ml, that is to say essentially 'sodium free'.

3. HOW TO TAKE FLAGYL-S SUSPENSION

Taking this medicine

Always take this medicine exactly as your doctor has told you. Shake the bottle gently before each use. It is important to finish a full course of treatment. The length of a course will depend on your needs and the illness being treated.

Check with your doctor or pharmacist if you are not sure.

- Take Flagyl-S Suspension by mouth.
- The dose of Flagyl-S Suspension will depend on your needs and the illness being treated. The length of your treatment will depend on the type of illness you have and how bad it is.
- Your doctor or pharmacist may dilute your Flagyl-S Suspension. In this case the pharmacist's label on the packaging will tell you exactly how much to take and how often.

The usual dose for adults and children is given below:

To treat certain bacterial infection

Adults and children over 10 years

- Take 10ml (400mg) of Flagyl-S Suspension once every 8 hours

Children and infants

- Your doctor will work out how much your child should take depending on their weight.
- Repeat the dose every 8 hours.
- If your child is a baby under 8 weeks of age, your doctor will give them one daily dose or two separate doses 12 hourly.

To prevent infections from happening after surgery

Children

Give your child Flagyl-S Suspension 2-3 hours before their operation. Your doctor will work out how much your child should take depending on their weight.

Other types of infections

For treatment of other infections caused by parasites and some bacteria your doctor will decide how much Flagyl-S Suspension you need to take and how often. This will depend on your illness and how

bad it is. The pharmacist's label on the packaging will tell you exactly how much Flagyl-S Suspension to take and how often.

People having kidney dialysis

Kidney dialysis removes Flagyl-S Suspension from your blood. If you are having kidney dialysis you must take this medicine after your dialysis treatment.

People with liver problems

Your doctor may tell you to take a lower dose or to use the medicine less often.

If you take more Flagyl-S Suspension than you should

If you take more Flagyl-S Suspension than you should, tell your doctor or go to your nearest hospital casualty department straight away. Take the pack with you. This is so the doctor knows what you have taken.

If you forget to take Flagyl-S Suspension

If you forget to take Flagyl-S Suspension, take it as soon as you remember. However, if it is almost time for your next dose, skip the missed dose. **Do not** use a double dose to make up for a forgotten dose.

If you stop taking Flagyl-S Suspension

You should only stop taking Flagyl-S Suspension if your doctor tells you to. It is essential to take the medicine and not to stop and start as this could cause unwanted side effects.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor or pharmacist straight away if you notice any of the following side effects:

- You get swelling of the hands, feet, ankles, face, lips or throat which may cause difficulty in swallowing or breathing. You could also notice an itchy, lumpy rash (hives) or nettle rash (urticaria). This may mean you are having an allergic reaction to Flagyl-S Suspension.
- You experience blistering or bleeding of the skin around the lips, eyes, mouth, nose and genitals. Also, flu-like symptoms and fever. This may be something called 'Stevens-Johnson syndrome'.
- You have a severe blistering rash where layers of the skin may peel off to leave large areas of raw exposed skin over the body. Also, a feeling of being generally unwell, fever, chills and aching muscles. This may be something called 'Toxic epidermal necrolysis'.
- Your skin is red and swollen and you have a skin rash of small pustules, you may also have a fever and swelling of the face. This may be something called 'Acute Generalised Exanthematous Pustulosis'.
- Flu-like symptoms, rash with a fever, enlarged lymph nodes, other abnormal blood test results like increase in a type of white blood cell (eosinophilia) or increased levels of liver enzymes. These could be signs/symptoms of a serious condition called DRESS (Drug Reaction with Eosinophilia and Systemic Symptoms). Contact a healthcare professional straight away if you experience any of the above signs/symptoms (applicable only for systemic formulations).
- A serious but very rare side effect is a brain disease (encephalopathy). Symptoms vary but you might get a fever, stiff neck, headache, see or hear things that aren't there. You might also have problems using your arms and legs, problems with speaking or feel confused.
- You develop symptoms such as abnormal sensitivity to light, abdominal pain, fever these could be symptoms of meningitis.
- Yellowing of the skin and eyes. This could be due to a liver problem (jaundice).
- Unexpected infections, mouth ulcers, bleeding gums, severe tiredness or weakness. This could be caused by a blood problem.
- Severe stomach pain which may reach through to your back (pancreatitis).

Tell your doctor or pharmacist if you notice any of the following side effects:

- Fits (convulsions)
- Mental problems such as feeling confused or seeing or hearing things that are not there (hallucinations)
- Problems with your eyesight such as blurred or double vision, changes in colour vision
- Tingling or burning pain in your eye
- Hearing problems, ringing in ears (tinnitus)
- Skin rash, urticaria, itching
- Headache
- Darkening of the urine
- Diarrhoea
- Feeling sleepy or dizzy
- Vertigo (spinning sensation)
- Pains in the muscles or joints
- Numbness, tingling, pain or a feeling of weakness in the arms or legs
- Fever
- Unpleasant taste in the mouth
- Furred tongue, dry mouth
- Feeling sick (nausea), being sick (vomiting), upset stomach, stomach pain or diarrhoea
- Loss of appetite, anorexia
- Depression
- Heart rhythm disturbances (so called QT prolongation which can be seen on ECG), particularly when Flagyl-S Suspension was used with other medicines that may cause heart rhythm disturbances
- Acute liver failure in patients with Cockayne Syndrome (see section 2 “Warnings and precautions”)
- PRES – A brain condition with symptoms such as headache, confusion, visual disturbance and seizures.

All medicines may have unwanted effects which are not mentioned in the package leaflet. If you notice any other changes in your health whilst taking this medicine, tell your doctor immediately.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRC Pharmacovigilance Website: www.hpra.ie.

5. HOW TO STORE FLAGYL-S SUSPENSION

Keep your medicine in a safe place, out of the sight and reach of children.

Do not use Flagyl-S Suspension after the expiry date which is stated on the bottle label and carton after 'EXP'. The expiry date refers to the last day of that month.

Do not store above 25°C. Store in the original container in order to protect the product from the light. After opening discard any unused suspension after 28 days.

Flagyl-S Suspension may be diluted, if necessary, with Syrup BP. Discard any unused diluted suspension after 14 days.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Flagyl-S 200mg/5ml Oral Suspension contains

The active substance in Flagyl-S Suspension is Metronidazole. Each 5ml of suspension contains Metronidazole Benzoate equivalent to 200mg of Metronidazole.

The oral suspension also contains the following ingredients: Sucrose, Sodium Dihydrogen Phosphate Dihydrate (E339), Aluminium Magnesium Silicate, Methyl Parahydroxybenzoate (E218), Propyl Parahydroxybenzoate (E216), Ethanol (96% v/v), 0.8%, Lemon Flavour (contains propylene glycol), Orange Oil Terpeneless and Purified Water.

What Flagyl-S 200mg/5ml Oral Suspension looks like and contents of the pack

Flagyl-S Suspension is a white to cream coloured oral suspension and is available in bottles of 100ml suspension packed in an outer carton.

Marketing Authorisation Holder and Manufacturer:

Marketing Authorisation Holder

Fidia farmaceutici S.p.A.
Via Ponte della Fabbrica, 3/A
35031 Abano Terme (PD) – Italy
Email: medinfo@fidiapharma.ie

Manufacturer

Unither Liquid Manufacturing,
1-3 Allee de la Neste, Z.I. en Sigal,
31770 Colomiers, France.

This leaflet was revised in September 2025.