

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Tildiem 60 mg Prolonged Release Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 60 mg diltiazem hydrochloride.

Excipients: Each tablet contains 125.5mg lactose monohydrate

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Prolonged released tablet.

Off-white biconvex tablets, engraved 'Tildiem 60' on one face.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Angina pectoris and hypertension.

4.2 Posology and method of administration

Adults: The usual dose is one tablet (60mg) three times daily. However, patient responses may vary and dosage requirements can differ significantly between individual patients. If necessary the dosage may be increased to 360mg/day. Higher doses up to 480mg/day have been used with benefit in some patients especially in unstable angina. There is no evidence of any decrease in efficacy at these high doses.

Elderly and patients with impaired hepatic or renal function: The recommended starting dose is one tablet (60mg) twice daily. The heart rate should be measured regularly in these groups of patients and the dose should not be increased if the heart rate falls below 40 beats per minute.

Tildiem 60mg Prolonged Release Tablets should be used with caution in patients with renal or hepatic impairment (see section 4.4 Special Warnings and Special Precautions For Use)

Children: Safety and efficacy in children have not been established. Therefore diltiazem is not recommended for use in children.

4.3 Contraindications

Pregnancy, women of childbearing potential (*see section 4.6 Pregnancy and Lactation*). Sick sinus syndrome except in the presence of a functioning ventricular pacemaker

Second or third degree AV block except in the presence of a functioning ventricular pacemaker

Severe bradycardia (below 40 beats per minute).

Left ventricular failure with pulmonary congestion.

Concomitant use with dantrolene infusion (see section 4.5 Interactions With Other Medicinal Products and Other Forms of Interaction).

Hypersensitivity to diltiazem or to any of the excipients.

4.4 Special warnings and precautions for use

Close observation is necessary in patients with reduced left ventricular function, bradycardia (risk of exacerbation) or with a first degree AV block detected on the electrocardiogram (risk of exacerbation and rarely of complete block).

Prior to general anaesthesia, the anaesthetist must be informed of ongoing diltiazem treatment. Depression of cardiac contractility, conductivity and automaticity, as well as the vascular dilatation associated with anaesthetics may be potentiated by calcium channel blockers.

Increase of plasma concentrations of diltiazem may be observed in the elderly and in patients with renal or hepatic insufficiency. The contraindications and precautions should be carefully observed and close monitoring, particularly of the heart rate, should be carried out at the beginning of treatment.

Calcium channel blocking agents, such as diltiazem, may be associated with mood changes, including depression.

Like other calcium channel antagonists, diltiazem has an inhibitory effect on intestinal motility. Therefore it should be used with caution in patients at risk to develop an intestinal obstruction. Tablet residues from slow release formulations of the product may pass into the patient's stools; however, this finding has no clinical relevance.

Careful monitoring is necessary in patients with latent or manifest diabetes mellitus due to a possible increase in blood glucose.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Concomitant use contraindicated:

Dantrolene (infusion)

Lethal ventricular fibrillation is regularly observed in animals when intravenous verapamil and dantrolene are administered concomitantly. The combination of calcium antagonist and dantrolene is therefore potentially dangerous (*see section 4.3 Contraindications*).

Concomitant use requiring caution:

Lithium:

Risk of increase in lithium-induced neurotoxicity.

Nitrate derivatives:

Increased hypotensive effects and faintness (additive vasodilating effects). In all patients treated with calcium antagonists, the prescription of nitrate derivatives should only be carried out at gradually increasing doses.

Theophylline:

Increase in circulating theophylline levels.

Alpha-antagonists:

Increased anti-hypertensive effects.

Concomitant treatment with alpha-antagonists may produce or aggravate hypotension. The combination of diltiazem with an alpha antagonist should be considered only with strict monitoring of blood pressure.

Amiodarone, Digoxin:

Increased risk of bradycardia; caution is required when these are combined with diltiazem, particularly in elderly subjects and when high doses are used.

Beta-blockers:

Possibility of rhythm disturbances (pronounced bradycardia, sinus arrest), sino-atrial and atrio-ventricular conduction disturbances and heart failure (synergistic effect).

Such a combination must only be used under close clinical and ECG monitoring, particularly at the beginning of treatment.

Other Antiarrhythmic agents:

Since diltiazem has antiarrhythmic properties, its concomitant prescription with other antiarrhythmic agents is not recommended (additive risk of increased cardiac adverse effects) This combination should only be used under close clinical and ECG monitoring.

Carbamazepine:

Increase in circulating carbamazepine levels. It is recommended that the plasma carbamazepine concentrations be assayed and that the dose should be adjusted if necessary.

Ciclosporin:

Increase in circulating ciclosporin levels. It is recommended that the ciclosporin dose be reduced, renal function be monitored, circulating ciclosporin levels be assayed and that the dose should be adjusted during combined therapy and after its discontinuation.

Anti-H₂ agents (cimetidine and ranitidine):

Increase in plasma diltiazem concentrations. Patients currently receiving diltiazem therapy should be carefully monitored when initiating or discontinuing therapy with anti-H₂ agents. An adjustment in diltiazem daily dose may be necessary.

Rifampicin:

Risk of decrease of diltiazem plasma levels after initiating therapy with rifampicin. The patient should be carefully monitored when initiating or discontinuing rifampicin treatment.

General information to be taken into account:

Due to the potential for additive effects, caution and careful titration are necessary in patients receiving diltiazem concomitantly with other agents known to affect cardiac contractility and/or conduction.

Grapefruit juice may increase diltiazem exposure (1.2 fold). Patients who consume grapefruit juice should be monitored for increased adverse effects of diltiazem. Grapefruit juice should be avoided if an interaction is suspected.

Diltiazem is metabolized by CYP3A4. A moderate (less than 2-fold) increase of diltiazem plasma concentration in cases of co-administration with a stronger CYP3A4 inhibitor has been documented. Diltiazem is also a CYP3A4 isoform inhibitor. Co-administration with other CYP3A4 substrates may result in an increase in plasma concentration of either co-administered drug. Co-administration of diltiazem with a CYP3A4 inducer may result in a decrease of diltiazem plasma concentrations.

BENZODIAZEPINES (midazolam, triazolam): Diltiazem significantly increases plasma concentrations of midazolam and triazolam and prolongs their half-life. Special care should be taken when prescribing short-acting benzodiazepines metabolized by the CYP3A4 pathway in patients using diltiazem.

CORTICOSTEROIDS (methylprednisolone): Inhibition of methylprednisolone metabolism (CYP3A4) and inhibition of P-glycoprotein. The patient should be monitored when initiating methylprednisolone treatment. An adjustment in the dose of methylprednisolone may be necessary.

STATINS: Diltiazem is an inhibitor of CYP3A4 and has been shown to significantly increase the AUC of some statins. The risk of myopathy and rhabdomyolysis due to statins metabolised by CYP3A4 may be increased with concomitant use of diltiazem. When possible, a non CYP3A4-metabolised statin should be used together with diltiazem, otherwise close monitoring for signs and symptoms of a potential statin toxicity is required.

4.6 Fertility, pregnancy and lactation

There are very limited data from the use of diltiazem in pregnant patients. Diltiazem has been shown to have reproductive toxicity in certain animal species (rat, mice, rabbit). Diltiazem is therefore not recommended during pregnancy, as well as in women of child-bearing potential not using effective contraception.

Diltiazem is excreted in breast milk at low concentrations. Breast-feeding while taking this drug should be avoided. If use of Diltiazem is considered medically essential, an alternative method of infant feeding should be instituted.

4.7 Effects on ability to drive and use machines

On the basis of reported adverse drug reactions, i.e. dizziness (common), malaise (common), the ability to drive and use machines could be altered. However, no studies have been performed.

4.8 Undesirable effects

The following CIOMS frequency rating is used, when applicable: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $\leq 1/100$); rare ($\geq 1/10,000$ to $\leq 1/1,000$); very rare ($\leq 1/10,000$); not known (cannot be estimated from the available data).

Within each frequency grouping, adverse events are presented in order of decreasing seriousness.

	Very Common	Common	Uncommon	Rare	Not known
<i>Blood and lymphatic system disorders</i>					Thrombocytopenia
<i>Psychiatric disorders</i>			Nervousness, insomnia		Mood changes (including depression)
<i>Nervous system disorders</i>		Headache, dizziness			Extrapyramidal syndrome
<i>Cardiac disorders</i>		Atrioventricular block (may be of first, second or third degree; bundle branch block may occur), palpitations	Bradycardia		Sinoatrial block, congestive heart failure
<i>Vascular disorders</i>		Flushing	Orthostatic hypotension		Vasculitis (including leukocytoclastic vasculitis)
<i>Gastrointestinal disorders</i>		Constipation, dyspepsia, gastric pain, nausea	Vomiting, Diarrhoea	Dry mouth	Gingival hyperplasia
<i>Metabolism and nutrition disorders</i>					Hyperglycemia
<i>Hepatobiliary disorders</i>			Hepatic enzymes increase (AST, ALT, LDH,		Hepatitis

<i>Skin and subcutaneous tissue disorders</i>		Erythema	ALP increase)	Urticaria	Photosensitivity (including lichenoid keratosis at sun exposed skin areas), angioneurotic oedema, erythema multiforme (including Steven-Johnson's syndrome and toxic epidermal necrolysis), sweating, exfoliative dermatitis, acute generalized exanthematous pustulosis, occasionally desquamative erythema with or without fever
<i>Reproductive system and breast disorders</i>					Gynecomastia
<i>General disorders and administration site conditions</i>	Peripheral oedema	Malaise			

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517; Website: www.hpra.ie; e-mail: medsafety@hpra.ie.

4.9 Overdose

The clinical effects of acute overdose can involve pronounced hypotension leading to collapse, sinus bradycardia with or without isorhythmic dissociation, and atrioventricular conduction disturbances.

Treatment, in a hospital setting, will include gastric lavage and/or osmotic diuresis. Conduction disturbances may be managed by temporary cardiac pacing.

Proposed corrective treatments: atropine, vasopressors, inotropic agents, glucagon and calcium gluconate infusion.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Calcium channel blockers, CO8 DB01.

Properties

Calcium antagonist, anti-anginal agent, antihypertensive agent.

Diltiazem selectively restricts calcium entry into the slow calcium channel of vascular smooth muscle and myocardial muscle fibres in a voltage-dependent manner. By this mechanism, diltiazem reduces the concentration of intracellular calcium in the vicinity of contractile proteins.

In animals:

Anti-anginal properties:

Diltiazem increases coronary blood flow without inducing any coronary steal phenomena. It acts both on small, large and collateral arteries. This vasodilator effect, which is moderate on peripheral systemic arterial territories, can be seen at doses that are not negatively inotropic. The two major active circulating metabolites, i.e. desacetyl diltiazem and N-monodesmethyl diltiazem, induce coronary vasodilation corresponding to 10 and 20% respectively of that of the parent compound.

Anti-hypertensive properties:

Diltiazem reduces arterial smooth muscle tone by reducing calcium influx in vascular smooth muscle cells, and causes vasodilation which produces a decrease in total peripheral resistance. Diltiazem reduces blood pressure without producing reflex tachycardia in various animal models of hypertension, particularly in the spontaneously hypertensive rat. It does not modify cardiac output and maintains renal blood flow. Furthermore, it preferentially inhibits the vasoconstrictor effects of noradrenaline and angiotensin II. Diltiazem increases diuresis and reduces cardiac hypertrophy in the spontaneously hypertensive rat.

High doses of diltiazem lessen the development of arterial calcinosis arterialis in the rat. The two major active circulating metabolites (desacetyl diltiazem and N-monodesmethyl diltiazem) possess pharmacological activity which is approximately 50% that of diltiazem.

In humans:

Anti-anginal properties:

Diltiazem increases coronary blood flow by reducing coronary resistance. Due to its moderate bradycardia-inducing activity and the reduction in systemic arterial resistance, diltiazem reduces cardiac workload.

Electrophysiologically, diltiazem causes moderate bradycardia in normal subjects, marginally prolongs intranodal conduction and has no effect on hisian and infrahisian conduction.

Antihypertensive properties:

At a vascular level, the calcium antagonist effect of diltiazem produces moderate arterial vasodilation and improves large artery compliance. This well-balanced vasodilation leads to a decrease in blood pressure on the hypertensive subject, due to lowered peripheral resistance, without producing reflex tachycardia. On the contrary, slight slowing of the heart rate is observed. Visceral blood flow rates, in particular renal and coronary, are maintained or increased.

A slight natriuretic effect is observed following acute administration. Diltiazem does not stimulate the renin-angiotensin-aldosterone system during long-term therapy, and does not cause water and sodium retention, as evidenced by the absence of body weight variation and a lack of change in the water and electrolyte balance of the plasma. Diltiazem acts as a coronary vasodilator on the heart, reducing left ventricular hypertrophy in the hypertensive subject. It has little effect on cardiac output.

Diltiazem reduces cardiac work by its moderate bradycardiac effect coupled with the lowering of systemic arterial resistance.

A negative inotropic effect has not been observed in a healthy myocardium. Diltiazem slows heart rate to a moderate extent and may exert a depressant effect on a diseased sinus node. It slows atrioventricular conduction and there is thus a risk of AV block.

Diltiazem does not affect conduction at the bundle of His or at an infrahisian level.

Diltiazem does not affect glycoregulation. It does not adversely affect plasma lipoproteins or lipid metabolism.

5.2 Pharmacokinetic properties

Diltiazem is well absorbed (90%) in healthy volunteers following oral administration.

Peak plasma concentrations occur 3 to 4 hours after dosing.

Due to a first pass effect, the bioavailability of the 60 mg tablet is about 40%.

The mean apparent plasma half-life is 4-8 hours.

Diltiazem is 80 to 85% bound to plasma proteins.

It is extensively metabolised by the liver.

The major circulating metabolite, N-monodesmethyl diltiazem accounts for approximately 35% of the circulating diltiazem.

Less than 5% of diltiazem is excreted unchanged in the urine.

There is a linear relationship between dose and plasma concentration. During long term administration to any one patient, plasma concentrations of diltiazem remain constant.

Mean plasma concentrations in elderly subjects and patients with renal and hepatic insufficiency are higher than in young subjects.

Diltiazem and its metabolites are poorly dialysed.

5.3 Preclinical safety data

In subacute and chronic dog and rat studies designed to produce toxicity, high doses of Tildiem 60mg Prolonged Release Tablets were associated with hepatic damage. In special subacute hepatic studies, oral doses of 125 mg/kg and higher in rats were associated with histological changes in the liver which were reversible when the drug was discontinued.

In dogs, doses of 20mg/kg were also associated with hepatic changes; however, these changes were reversible with continued dosing.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Lactose monohydrate
Castor oil, hydrogenated
Macrogol 6000
Magnesium stearate.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

This medicinal product does not require any special temperature storage conditions. Store in the original package.

This product is sensitive to moisture.

6.5 Nature and contents of container

90 or 100 tablets in PVC/foil strips.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Sanofi-aventis Ireland Ltd. T/A SANOFI
Citywest Business Campus
Dublin 24
Ireland

8 MARKETING AUTHORISATION NUMBER

PA0540/161/003

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 7th May 1999

Date of last renewal: 7th May 2009

10 DATE OF REVISION OF THE TEXT

November 2014