

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Innopril 10 mg Tablets.

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 10 mg Enalapril maleate.
For excipients, see 6.1.

3 PHARMACEUTICAL FORM

Tablet

Innopril 10mg Tablets are rusty red, arc triangle shaped, biconvex tablets, debossed with "10" over "G" on one side and scoreline on the other side. Approximately 9mm long, 9mm wide and 4mm thick.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

- Treatment of Hypertension.
- Treatment of Symptomatic Heart Failure.
- Prevention of Symptomatic Heart Failure in patients with Asymptomatic Left ventricular Dysfunction (ejection fraction <35%).

(See Section 5.1 Pharmacodynamic properties)

4.2 Posology and method of administration

The absorption of Enalapril is not affected by food.

The dose should be individualised according to patient profile (see 4.4 Special warnings and precautions for use) and blood pressure response.

Hypertension:

The initial dose is 5 to maximally 20 mg, depending on the degree of hypertension and the condition of the patient (see below). Enalapril is given once daily. In mild hypertension, the recommended initial dose is 5 to 10 mg. Patients with a strongly activated renin-angiotensin-aldosterone system (e.g., renovascular hypertension, salt and/or volume depletion, cardiac decompensation, or severe hypertension) may experience an excessive blood pressure fall following the initial dose. A starting dose of 5 mg or lower is recommended in such patients and the initiation of treatment should take place under medicinal supervision.

Prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with Enalapril. A starting dose of 5 mg or lower is recommended in such patients. If possible, diuretic therapy should be discontinued for 2-3 days prior to initiation of therapy with Enalapril. Renal function and serum potassium should be monitored.

The usual maintenance dose is 20 mg daily. The maximum maintenance dose is 40 mg daily.

Heart failure/asymptomatic left ventricular dysfunction:

In the management of symptomatic heart failure, Enalapril is used in addition to diuretics and, where appropriate, digitalis or beta-blockers. The initial dose of Enalapril in patients with symptomatic heart failure or asymptomatic left

ventricular dysfunction is 2.5mg, and it should be administered under close medical supervision to determine the initial effect on the blood pressure. In the absence of, or after effective management of, symptomatic hypotension following initiation of therapy with Enalapril in heart failure, the dose should be increased gradually to the usual maintenance dose of 20 mg, given in a single dose or two divided doses, as tolerated by the patient. This dose titration is recommended to be performed over a 2 to 4 week period. The maximum dose is 40 mg daily given in two divided doses.

Suggested Dosage Titration of Enalapril in Patients with heart failure/asymptomatic left ventricular dysfunction

Week	Dose mg/day
Week 1	Days 1 to 3: 2.5 mg/day* in a single dose Days 4 to 7: 5 mg/day in two divided doses
Week 2	10 mg /day in a single dose or in two divided doses
Week 3 and 4	20 mg/day in a single dose or in two divided doses

- Special precautions should be followed in patients with impaired renal function or taking diuretics (See 4.4 Special warnings and precautions for use).

Blood pressure and renal function should be monitored closely both before and after starting treatment with Enalapril (See 4.4 Special warnings and precautions for use) because hypotension and (more rarely) consequent renal failure have been reported. In patients treated with diuretics, the dose should be reduced if possible before beginning treatment with Enalapril. The appearance of hypotension after the initial dose of Enalapril does not imply that hypotension will recur during chronic therapy with Enalapril and does not preclude continued use of the drug. Serum potassium and renal function also should be monitored.

Dosage in Renal Insufficiency:

Generally, the intervals between the administrations of Enalapril should be prolonged and/or the dosage reduced.

Creatinine Clearance (CrCL) mL/min	Initial Dose mg/day
30<CRCL<80 ml/min	5 – 10 mg
10<CRCL≤30 ml/min	2.5 mg
CrCL≤10 ml/min	2.5 mg on dialysis days*

* See 4.4 Special warnings and precautions for use – *Hemodialysis Patients*. Enalaprilat is dialysable. Dosage on non-dialysis days should be adjusted depending on the blood pressure response.

Use in Elderly

The dose should be in line with the renal function of the elderly patient (see 4.4 Special warnings and precautions for use, Renal Function Impairment).

Use in paediatrics

There is limited clinical trial experience of the use of Enalapril in hypertensive paediatric patients. (see 4.4 Special warnings and precautions for use, 5.1 Pharmacodynamic properties and 5.2 Pharmacokinetic properties).

For patients who can swallow tablets, the dose should be individualized according to patient profile and blood pressure response. The recommended initial dose is 2.5 mg in patients 20 to < 50 kg and 5 mg in patients \geq 50 kg. Enalapril is given once daily. The dosage should be adjusted according to the needs of the patient to a maximum of 20 mg daily in patients 20 to <50 kg and 40 mg in patients \geq 50 kg. (See 4.4 Special warnings and precautions for use).

Enalapril is not recommended in neonates and in paediatric patients with glomerular filtration rate <30 ml/min/1.73m², as no data are available.

4.3 Contraindications

- Hypersensitivity to Enalapril, to any of the excipients or any other ACE inhibitor.
- History of angioedema associated with previous ACE inhibitor therapy
- Hereditary or idiopathic angioedema
- Second and third trimesters of pregnancy (see sections 4.4 and 4.6)

4.4 Special warnings and precautions for use

Symptomatic Hypotension

Symptomatic hypotension is rarely seen in uncomplicated hypertensive patients. In hypertensive patients receiving Enalapril, symptomatic hypotension is more likely to occur if the patient has been volume-depleted, e.g., by diuretic therapy, dietary salt restriction, dialysis, diarrhoea or vomiting (See 4.5 Interaction with other medicaments and other forms of interaction and 4.8 Undesirable Effects). In patients with heart failure, with or without associated renal insufficiency, symptomatic hypotension has been observed.

This is most likely to occur in those patients with more severe degrees of heart failure, as reflected by the use of high doses of loop diuretics, hyponatremia or functional renal impairment. In these patients, therapy should be started under medical supervision and the patients should be followed closely whenever the dose of Enalapril and/or diuretic is adjusted. Similar considerations may apply to patients with ischemic heart or cerebrovascular disease in whom an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident.

If hypotension occurs, the patient should be placed in the supine position and, if necessary, should receive an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further doses, which can be given usually without difficulty once the blood pressure has increased after volume expansion.

In some patients with heart failure who have normal or low blood pressure, additional lowering of systemic blood pressure may occur with Enalapril. This effect is anticipated, and usually is not a reason to discontinue treatment. If hypotension becomes symptomatic, a reduction of dose and/or discontinuation of the diuretic and/or Enalapril may be necessary.

Aortic or Mitral Valve Stenosis/ Hypertrophic Cardiomyopathy

As with all vasodilators, ACE inhibitors should be given with caution in patients with left ventricular valvular and outflow tract obstruction and avoided in cases of cardiogenic shock and haemodynamically significant obstruction.

Renal Function Impairment

In cases of renal impairment (creatinine clearance <80 ml/min) the initial Enalapril dosage should be adjusted according to the patient's creatinine clearance (See 4.2 Posology and method of administration) and then as a function of the patient's response to treatment. Routine monitoring of potassium and creatinine are part of normal medical practice for these patients.

Renal failure has been reported in association with Enalapril and has been mainly in patients with severe heart failure or underlying renal disease, including renal artery stenosis. If recognized promptly and treated appropriately, renal failure when associated with therapy with Enalapril is usually reversible.

Some hypertensive patients, with no apparent pre-existing renal disease have developed increases in blood urea and creatinine when Enalapril has been given concurrently with a diuretic. Dosage reduction of Enalapril and/or discontinuation of the diuretic may be required. This situation should raise the possibility of underlying renal artery stenosis (See 4.4 Special warnings and precautions for use, *Renovascular hypertension*).

Renovascular hypertension

There is an increased risk of hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with ACE inhibitors. Loss of renal function may occur with only mild changes in serum creatinine. In these patients, therapy should be initiated under close medical supervision with low doses, careful titration, and monitoring of renal function.

Kidney transplantation

There is no experience regarding the administration of Enalapril in patients with recent kidney transplantation. Treatment with Enalapril is therefore not recommended.

Hepatic failure

Rarely, ACE inhibitors have been associated with a syndrome that starts with cholestatic jaundice or hepatitis and progresses to fulminant hepatic necrosis and (sometimes) death. The mechanism of this syndrome is not understood. Patients receiving ACE inhibitors who develop jaundice or marked elevations of hepatic enzymes should discontinue the ACE inhibitor and receive appropriate medical follow-up.

Neutropenia/Agranulocytosis

Neutropenia/agranulocytosis, thrombocytopenia and anaemia have been reported in patients receiving ACE inhibitors. In patients with normal renal function and no other complicating factors, neutropenia occurs rarely. Enalapril should be used with extreme caution in patients with collagen vascular disease, immunosuppressant therapy, treatment with allopurinol or procainamide, or a combination of these complicating factors, especially if there is pre-existing impaired renal function. Some of these patients developed serious infections, which in a few instances did not respond to intensive antibiotic therapy.

If Enalapril is used in such patients, periodic monitoring of white blood cell counts is advised and patients should be instructed to report any sign of infection.

Hypersensitivity/Angioneurotic oedema

Angioneurotic oedema of the face, extremities, lips, tongue, glottis and/or larynx has been reported in patients treated with angiotensin-converting enzyme inhibitors, including Enalapril. This may occur at any time during treatment. In such cases, Enalapril should be discontinued promptly and appropriate monitoring should be instituted to ensure complete resolution of symptoms prior to dismissing the patient.

Even in those instances where swelling of only the tongue is involved, without respiratory distress, patients may require prolonged observation since treatment with antihistamines and corticosteroids may not be sufficient.

Very rarely, fatalities have been reported due to angioedema associated with laryngeal oedema or tongue oedema. Patients with involvement of the tongue, glottis or larynx are likely to experience airway obstruction, especially those with a history of airway surgery. Where there is involvement of the tongue, glottis or larynx, likely to cause airway obstruction, appropriate therapy, which may include subcutaneous epinephrine solution 1:1000 (0.3 ml to 0.5 ml) and/or measures to ensure a patent airway, should be administered promptly.

Black patients receiving ACE inhibitors have been reported to have a higher incidence of angioedema compared to non-blacks.

Patients with a history of angioedema unrelated to ACEinhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor. (Also see 4.3 Contraindications).

Anaphylactoid reactions during hymenoptera desensitisation

Rarely, patients receiving ACE inhibitors during desensitisation with hymenoptera venom have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding ACE-inhibitor therapy prior to each desensitisation.

Anaphylactoid reactions during LDL Apheresis

Rarely, patients receiving ACE inhibitors during low density lipoprotein [LDL] apheresis with dextran sulphate have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding ACE-inhibitor therapy prior to each apheresis.

Haemodialysis patients

Anaphylactoid reactions have been reported in patients dialysed with high-flux membranes (e.g. AN 69[®]) and treated concomitantly with an ACE inhibitor. In these patients, consideration should be given to using a different type of dialysis membrane or a different class of antihypertensive agent.

Hypoglycaemia

In diabetic patients treated with oral antidiabetic agents or insulin, glycaemic control should be closely monitored especially during the first month of treatment with an ACE inhibitor. (See 4.5 Interaction with other medicinal products and other forms of interaction, *Antidiabetics*.)

Cough

Cough has been reported with the use of ACE inhibitors. Characteristically, the cough is non-productive, persistent and resolves after discontinuation of therapy. ACE inhibitor-induced cough should be considered as part of the differential diagnosis of cough.

Surgery/Anaesthesia

In patients undergoing major surgery or during anaesthesia with agents that produce hypotension, Enalapril blocks angiotensin II formation secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it can be corrected by volume expansion.

Hyperkalaemia

Elevations in serum potassium have been observed in some patients treated with ACE inhibitors, including enalapril. Risk factors for the development of hyperkalaemia include those with renal insufficiency, worsening of renal function, age (>70 years) diabetes mellitus, inter-current events in particular dehydration, acute cardiac decompensation, metabolic acidosis and concomitant use of potassium-sparing diuretics (e.g., spironolactone, eplerenone, triamterene, or amiloride), potassium supplements or potassium-containing salt substitutes; or those patients taking other drugs associated with increases in serum potassium (e.g. heparin). The use of potassium supplements, potassium-sparing diuretics, or potassium-containing salt substitutes particularly in patients with impaired renal function may lead to a significant increase in serum potassium. Hyperkalaemia can cause serious, sometimes fatal arrhythmias. If concomitant use of enalapril and any of the above-mentioned agents is deemed appropriate, they should be used with caution and with frequent monitoring of serum potassium. (See 4.5 'Interaction with other medicinal products and other forms of interaction'.)

Lithium

The combination of lithium and Enalapril is generally not recommended (see 4.5 Interaction with other medicinal products and other forms of interaction).

Lactose

Innopril 10mg Tablets contains less than 200 mg of lactose per tablet. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Paediatric use

There is limited efficacy and safety experience in hypertensive children >6 years old, but no experience in other indications. Limited pharmacokinetic data are available in children above 2 months of age. (Also see 4.2 Posology and method of administration, 5.1 Pharmacodynamic properties, and 5.2 Pharmacokinetic properties.)

Enalapril is not recommended in children in other indications than hypertension.

Enalapril is not recommended in neonates and in paediatric patients with glomerular filtration rate <30 ml/min/1.73 m², as no data are available. (See 4.2 Posology and method of administration.)

Pregnancy and lactation

ACE inhibitors should not be initiated during pregnancy. Unless continued ACE inhibitor therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

Use of Enalapril is not recommended during breast feeding (see sections 4.6 and 5.2)

Ethnic differences

As with other angiotensin-converting enzyme inhibitors, Enalapril is apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of a higher prevalence of low-renin states in the black hypertensive population.

4.5 Interaction with other medicinal products and other forms of interaction

Potassium-sparing diuretics or potassium supplements

ACE inhibitors attenuate diuretic-induced potassium loss. Potassium-sparing diuretics (e.g. spironolactone, triamterene or amiloride), potassium supplements, or potassium-containing salt substitutes may lead to significant increases in serum potassium. If concomitant use is indicated because of demonstrated hypokalaemia they should be used with caution and with frequent monitoring of serum potassium (see 4.4 Special warnings and precautions for use).

Diuretics (thiazide or loop diuretics)

Prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with Enalapril (see 4.4 Special warnings and precautions for use). The hypotensive effects can be reduced by discontinuation of the diuretic, by increasing volume or salt intake or by initiating therapy with a low dose of Enalapril.

Other antihypertensive agents

Concomitant use of these agents may increase the hypotensive effects of Enalapril. Concomitant use with nitroglycerine and other nitrates, or other vasodilators, may further reduce blood pressure.

Lithium

Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with ACE inhibitors.

Concomitant use of thiazide diuretics may further increase lithium levels and enhance the risk of lithium toxicity with ACE inhibitors. Use of Enalapril with lithium is not recommended, but if the combination proves necessary, careful monitoring of serum lithium levels should be performed (see 4.4 Special warnings and precautions for use).

Tricyclic antidepressants/Antipsychotics/Anaesthetics/Narcotics

Concomitant use of certain anaesthetic medicinal products, tricyclic antidepressants and antipsychotics with ACE inhibitors may result in further reduction of blood pressure (see 4.4 Special warnings and precautions for use).

Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)

Chronic administration of NSAIDs may reduce the antihypertensive effect of an ACE inhibitor. NSAIDs (including

COX-2 inhibitors) and ACE inhibitors exert an additive effect on the increase in serum potassium, and may result in a deterioration of renal function. These effects are usually reversible. Rarely, acute renal failure may occur, especially in patients with compromised renal function (such as the elderly or patients who are volume-depleted, including those on diuretic therapy). Patients should be adequately hydrated and consideration should be given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.

Gold

Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including enalapril.

Sympathomimetics

Sympathomimetics may reduce the antihypertensive effects of ACE inhibitors.

Antidiabetics

Epidemiological studies have suggested that concomitant administration of ACE inhibitors and antidiabetic medicines (insulins, oral hypoglycaemic agents) may cause an increased blood-glucose-lowering effect with risk of hypoglycaemia. This phenomenon appeared to be more likely to occur during the first weeks of combined treatment and in patients with renal impairment. (See 4.4 Special warnings and precautions for use; Hypoglycaemia and 4.8 Undesirable effects; Metabolism and nutrition disorders).

Alcohol

Alcohol enhances the hypotensive effect of ACE inhibitors.

Acetyl salicylic acid, thrombolytics and β -blockers.

Enalapril can be safely administered concomitantly with acetyl salicylic acid (at cardiologic doses), thrombolytics and β -blockers.

4.6 Fertility, pregnancy and lactation

Pregnancy

The use of ACE inhibitors is not recommended during the first trimester of pregnancy (see section 4.4). The use of ACE inhibitors is contra-indicated during the second and third trimester of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Unless continued ACE inhibitors therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to ACE inhibitor therapy during the second and third trimesters is known to induce human foetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). (See section 5.3).

Should exposure to ACE inhibitors have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.

Infants whose mothers have taken ACE inhibitors should be closely observed for hypotension (see sections 4.3 and 4.4).

Lactation

Limited pharmacokinetic data demonstrate very low concentrations in breast milk (see section 5.2). Although these concentrations seem to be clinically irrelevant the use of Enalapril Maleate in breast-feeding is not recommended for preterm infants and for the first few weeks after delivery, because of the hypothetical risk of cardiovascular and renal effects and because there is not enough clinical experience. In case of an older infant the use of Enalapril Maleate in breast-feeding mother may be considered if this treatment is necessary for the mother and the child is observed for any adverse effect.

4.7 Effects on ability to drive and use machines

When driving vehicles or operating machines it should be taken into account that occasionally dizziness or weariness may occur.

4.8 Undesirable effects

Undesirable effects reported for Enalapril include:

Very common (>1/10)

Common (>1/100, <1/10)

Uncommon (>1/1,000, <1/100)

Rare (>1/10,000, <1/1,000)

Very rare (<1/10,000), including isolated reports.

Blood and lymphatic system disorders:

Uncommon: anaemia (including aplastic and haemolytic).

Rare: neutropenia, decreases in haemoglobin, decreases in haematocrit, thrombocytopenia, agranulocytosis, bone marrow depression, pancytopenia, lymphadenopathy, autoimmune diseases.

Endocrine disorders:

not known: syndrome of inappropriate antidiuretic hormone secretion (SIADH)

Metabolism and nutrition disorders:

Uncommon: hypoglycaemia (see 4.4 Special warnings and precautions for use, *Hypoglycemia*).

Nervous system and psychiatric disorders:

Common: headache, depression.

Uncommon: confusion, somnolence, insomnia, nervousness, paresthesia, vertigo.

Rare: dream abnormality, sleep disorders.

Eye disorders:

Very common: blurred vision.

Cardiac and vascular disorders:

Very common: dizziness

Common: hypotension (including orthostatic hypotension), syncope, chest pain, rhythm disturbances, angina pectoris, tachycardia

Uncommon: orthostatic hypotension, palpitations, myocardial infarction or cerebrovascular accident*, possibly secondary to excessive hypotension in high risk patients (see 4.4 Special warnings and precautions for use)

* Incidence rates were comparable to those in the placebo and active control groups in the clinical trials

Rare: Raynaud's phenomenon

Respiratory, thoracic and mediastinal disorders:

Very common: cough.

Common: dyspnea.

Uncommon: rhinorrhea, sore throat and hoarseness, bronchospasm/asthma.

Rare: pulmonary infiltrates, rhinitis, allergic alveolitis/eosinophilic pneumonia.

Gastro-intestinal disorders:

Very common: nausea.

Common: diarrhoea, abdominal pain, taste alteration.

Uncommon: ileus, pancreatitis, vomiting, dyspepsia, constipation, anorexia, gastric irritations, dry mouth, peptic ulcer.

Rare: stomatitis/aphthous ulcerations, glossitis.

Very rare: intestinal angioedema

Hepatobiliary disorders:

Rare: hepatic failure, hepatitis – either hepatocellular or cholestatic, hepatitis including necrosis, cholestasis (including jaundice).

Skin and subcutaneous tissue disorders:

Common: rash, hypersensitivity/angioneurotic oedema: angioneurotic oedema of the face, extremities, lips, tongue, glottis and/or larynx has been reported (see 4.4 Special warnings and precautions for use).

Uncommon: diaphoresis, pruritus, urticaria, alopecia.

Rare: erythema multiforme, Stevens-Johnson syndrome, exfoliative dermatitis, toxic epidermal necrolysis, pemphigus, erythroderma.

A symptom complex has been reported which may include some or all of the following: fever, serositis, vasculitis, myalgia/myositis, arthralgia/arthritis, a positive ANA, elevated ESR, eosinophilia, and leukocytosis. Rash, photosensitivity or other dermatologic manifestations may occur.

Renal and urinary disorders:

Uncommon: renal dysfunction, renal failure, proteinuria.

Rare: oliguria.

Reproductive system and breast disorders:

Uncommon: impotence.

Rare: gynecomastia.

General disorders and administration site conditions:

Very common: asthenia.

Common: fatigue.

Uncommon: muscle cramps, flushing, tinnitus, malaise, fever.

Investigations:

Common: hyperkalaemia, increases in serum creatinine.

Uncommon: increases in blood urea, hyponatraemia.

Rare: elevations of liver enzymes, elevations of serum bilirubin.

4.9 Overdose

Limited data are available in humans. The most prominent symptoms reported to date are severe hypotension, shock, bradycardia, electrolyte disturbances and renal failure, with onset approximately 6 hours after ingestion, concurrent with blocking of the renin-angiotensin system and stupor. Serum Innopril levels 100-200 – fold higher than usually seen after therapeutic doses have been reported after ingestion of 300 mg and 440 mg of Innopril, respectively. The recommended treatment of overdose is intravenous infusion of normal saline solution.

After ingestion of an overdose, the patient should be kept under close supervision, preferably in an intensive care unit. Serum electrolytes and creatinine should be monitored frequently. Measurements to prevent absorption (gastric lavage, activated charcoal) should be applied at the physician's own discretion.

If hypotension occurs, the patient should be placed in shock position and intravenously salt and volume supplementation should be given rapidly. Treatment with angiotensin II should be considered. Bradycardia should be treated by administering atropin. The use of a pacemaker may be considered. ACE inhibitors may be removed from the

circulation by haemodialysis. The use of high-flux polyacrylonitrile membranes should be avoided.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Angiotensin-converting enzyme inhibitors, ATC Code: C09A A02

Enalapril, in its maleate salt, is a derivative of two amino-acids, L-alanine and L-proline. Angiotensin converting enzyme (ACE) is a peptidyl dipeptidase which catalyses the conversion of angiotensin I to the pressor substance angiotensin II. After absorption, Enalapril is hydrolysed to Enalaprilat, which inhibits ACE. Inhibition of ACE results in decreased plasma angiotensin II, which leads to increased plasma renin activity (due to removal of negative feedback of renin release), and decreased aldosterone secretion.

ACE is identical to kininase II. Thus Enalapril may also block the degradation of bradykinin, a potent vasodepressor peptide. However, the role that this plays in the therapeutic effects of Enalapril remains to be elucidated.

While the mechanism through which Enalapril lowers blood pressure is believed to be primarily suppression of the renin-angiotensin-aldosterone system, Enalapril is antihypertensive even in patients with low-renin hypertension.

Administration of Enalapril to patients with hypertension results in a reduction of both supine and standing blood pressure without a significant increase in heart rate.

Symptomatic postural hypotension is infrequent. In some patients the development of optimal blood pressure reduction may require several weeks of therapy. Abrupt withdrawal of Enalapril has not been associated with rapid increase in blood pressure.

Effective inhibition of ACE activity usually occurs 2 to 4 hours after oral administration of an individual dose of Enalapril. Onset of antihypertensive activity was usually seen at one hour, with peak reduction of blood pressure achieved by 4 to 6 hours after administration. The duration of effect is dose-related. However, at recommended doses, antihypertensive and haemodynamic effects have been shown to be maintained for at least 24 hours.

In haemodynamic studies in patients with essential hypertension, blood pressure reduction was accompanied by a reduction in peripheral arterial resistance with an increase in cardiac output and little or no change in heart rate. Following administration of Enalapril there was an increase in renal blood flow; glomerular filtration rate was unchanged. There was no evidence of sodium or water retention. However, in patients with low pre-treatment glomerular filtration rates, the rates were usually increased.

In short-term clinical studies in diabetic and non-diabetic patients with renal disease, decreases in albuminuria and urinary excretion of IgG and total urinary protein were seen after the administration of Enalapril.

When given together with thiazide-type diuretics, the blood pressure-lowering effects of Enalapril are at least additive. Enalapril may reduce or prevent the development of thiazide-induced hypokalaemia.

In patients with heart failure on therapy with digitalis and diuretics, treatment with oral or injected Enalapril was associated with decreases in peripheral resistance and blood pressure. Cardiac output increased, while heart rate (usually elevated in patients with heart failure) decreased. Pulmonary capillary wedge pressure was also reduced. Exercise tolerance and severity of heart failure, as measured by New York Heart Association criteria, improved. These actions continued during chronic therapy.

In patients with mild to moderate heart failure, Enalapril retarded progressive cardiac dilatation/enlargement and failure, as evidenced by reduced left ventricular end diastolic and systolic volumes and improved ejection fraction.

A multicentre, randomised, double-blind, placebo-controlled trial (SOLVD Prevention trial) examined a population with asymptomatic left ventricular dysfunction (LVEF<35%). 4228 patients were randomised to receive either placebo (n=2117) or Enalapril (n=2111). In the placebo group, 818 patients had heart failure or died (38.6%) as compared with

630 in the Enalapril group (29.8%) (risk reduction: 29%; 95% CI; 21 - 36%; $p < 0.001$). 518 patients in the placebo group (24.5%) and 434 in the Enalapril group (20.6%) died or were hospitalised for new or worsening heart failure (risk reduction 20%; 95% CI; 9-30%; $p < 0.001$).

A multicentre, randomised, double-blind, placebo-controlled trial (SOLVD treatment trial) examined a population with symptomatic congestive heart failure due to systolic dysfunction (ejection fraction $< 35\%$). 2569 patients receiving conventional treatment for heart failure were randomly assigned to receive either placebo ($n=1284$) or Enalapril ($n=1285$). There were 510 deaths in the placebo group (39.7%) as compared with 452 in the Enalapril group (35.2%) (reduction in risk, 16%; 95% CI, 5 - 26%; $p=0.0036$). There were 461 cardiovascular deaths in the placebo group as compared with 399 in the Enalapril group (risk reduction 18%, 95% CI, 6 - 28%, $p < 0.002$), mainly due to a decrease of deaths due to progressive heart failure (251 in the placebo group vs 209 in the Enalapril group, risk reduction 22%, 95% CI, 6 - 35%). Fewer patients died or were hospitalised for worsening heart failure (736 in the placebo group and 613 in the Enalapril group; risk reduction, 26%; 95% CI, 18 - 34%; $p < 0.0001$).

Overall in SOLVD study, in patients with left ventricular dysfunction, Enalapril reduced the risk of myocardial infarction by 23% (95% CI, 11 – 34%; $p < 0.001$) and reduced the risk of hospitalisation for unstable angina pectoris by 20% (95% CI, 9 – 29%; $p < 0.001$).

There is limited experience of the use in hypertensive paediatric patients > 6 years. In a clinical study involving 110 hypertensive paediatric patients 6 to 16 years of age with a body weight ≥ 20 kg and a glomerular filtration rate > 30 ml/min/1.73 m², patients who weighed < 50 kg received either 0.625, 2.5 or 20 mg of Enalapril daily and patients who weighed ≥ 50 kg received either 1.25, 5 or 40 mg of Enalapril daily. Enalapril administration once daily lowered trough blood pressure in a dose-dependent manner. The dose-dependent antihypertensive efficacy of Enalapril was consistent across all subgroups (age, Tanner stage, gender, race). However, the lowest doses studied, 0.625 mg and 1.25 mg, corresponding to an average of 0.02 mg/kg once daily, did not appear to offer consistent antihypertensive efficacy. The maximum dose studied was 0.58 mg/kg (up to 40 mg) once daily. The adverse experience profile for paediatric patients is not different from that seen in adult patients.

5.2 Pharmacokinetic properties

Absorption

Oral Enalapril is rapidly absorbed, with peak serum concentrations of Enalapril occurring within one hour. Based on urinary recovery, the extent of absorption of Enalapril from oral Enalapril tablet is approximately 60%. The absorption of oral Enalapril is not influenced by the presence of food in the gastro-intestinal tract.

Following absorption, oral Enalapril is rapidly and extensively hydrolysed to Enalaprilat, a potent angiotensin-converting enzyme inhibitor. Peak serum concentrations of Enalaprilat occur about 4 hours after an oral dose of Enalapril tablet. The effective half-life for accumulation of Enalaprilat following multiple doses of oral Enalapril is 11 hours. In subjects with normal renal function, steady-state serum concentrations of Enalaprilat were reached after 4 days of treatment.

Distribution

Over the range of concentrations, which are therapeutically relevant, Enalaprilat binding to human plasma proteins does not exceed 60%.

Biotransformation

Except for conversion of Enalaprilat, there is no evidence for significant metabolism of Enalapril.

Elimination

Excretion of Enalaprilat is primarily renal. The principal components in the urine are Enalaprilat, accounting for about 40% of the dose, and intact Enalapril (about 20%).

Renal impairment

The exposure of Enalapril and Enalaprilat is increased in patients with renal insufficiency. In patients with mild to moderate renal insufficiency (creatinine clearance 40-60 ml/min) steady state AUC of Enalaprilat was approximately two-fold higher than in patients with normal renal function after administration of 5 mg once daily. In severe renal

impairment (creatinine clearance ≤ 30 ml/min), AUC was increased approximately 8-fold. The effective half-life of Enalaprilat following multiple doses of Enalapril Maleate is prolonged at this level of renal insufficiency and time to steady state is delayed. (See 4.2 Posology and method of administration). Enalaprilat may be removed from the general circulation by haemodialysis. The dialysis clearance is 62 ml/min.

Children and adolescents

A multiple dose pharmacokinetic study was conducted in 40 hypertensive male and female paediatric patients aged 2 months to ≤ 16 years following daily oral administration of 0.07 to 0.14 mg/kg Enalapril Maleate. There were no major differences in the pharmacokinetics of Enalaprilat in children compared with historic data in adults. The data indicate an increase in AUC (normalised to dose per body weight) with increased age; however, an increase in AUC is not observed when data are normalised by body surface area. At steady state, the mean effective half-life for accumulation of Enalaprilat was 14 hours.

Lactation

After a single 20 mg oral dose in 5 postpartum women the average peak Enalapril milk level was 1.7 $\mu\text{g/L}$ (range 0.54 to 5.9 $\mu\text{g/L}$) at 4 to 6 hours after the dose. The average peak Enalaprilat level was 1.7 $\mu\text{g/L}$ (range 1.2 to 2.3 $\mu\text{g/L}$); peaks occurred at various times over the 24-hour period. Using the peak milk level data, the estimated maximum intake of an exclusively breastfed infant would be about 0.16% of the maternal weight-adjusted dosage.

A woman who had been taking oral Enalapril 10 mg daily for 11 months had peak Enalapril milk levels of 2 $\mu\text{g/L}$ 4 hours after a dose and peak Enalaprilat levels of 0.75 $\mu\text{g/L}$ about 9 hours after the dose. The total amount of Enalapril and Enalaprilat measured in milk during the 24 hour period was 1.44 $\mu\text{g/L}$ and 0.63 $\mu\text{g/L}$ of milk respectively.

Enalaprilat milk levels were undetectable (<0.2 $\mu\text{g/L}$) 4 hours after a single dose of Enalapril 5 mg in 1 mother and 10 mg in 2 mothers; Enalapril levels were not determined.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity and carcinogenic potential. Reproductive toxicity studies suggest that Enalapril has no effects on fertility and reproductive performance in rats, and is not teratogenic. In a study in which female rats were dosed prior to mating through gestation, an increased incidence of rat pup deaths occurred during lactation. The compound has been shown to cross the placenta and is secreted in milk. Angiotensin-converting enzyme inhibitors, as a class, have been shown to be foetotoxic (causing injury and/or death to the foetus) when given in the second or third trimester.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium hydrogen carbonate
 Pregelatinized Starch
 Maize Starch
 Lactose Monohydrate
 Magnesium Stearate
 Iron oxide (E172)(10mg and 20mg tablet only)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

2 years

6.4 Special precautions for storage

Do not store above 25°C.
Store in the original container.
Keep the container tightly closed.

6.5 Nature and contents of container

- (i) High density polyethylene tablet container with desiccant and tamper-evident polypropylene cap.
-pack sizes 10, 11, 14, 20, 28, 30, 50, 56, 60, 84, 90, 100, 250 and 500
- (ii) aluminium foil/aluminium foil blister packs in cardboard outer carton.
- pack sizes 10, 11, 14, 20, 28, 30, 49, (49 x 1 blister), 50, 56, 60, 84, 90, 98,100, 250 and 500

Not all pack sizes may be marketed.

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

7 MARKETING AUTHORISATION HOLDER

McDermott Laboratories Limited
trading as Gerard Laboratories
35-36 Baldoyle Industrial Estate
Grange Road
Dublin 13
Ireland

8 MARKETING AUTHORISATION NUMBER

PA 0577/035/003

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 27 October 2000

Date of last renewal: 22 January 2009

10 DATE OF REVISION OF THE TEXT

September 2011