

**PACKAGE LEAFLET**

## **Package leaflet: Information for the patient**

### **Montelukast Viatris 10 mg film-coated tablets** montelukast

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Montelukast Viatris is and what it is used for
2. What you need to know before you take Montelukast Viatris
3. How to take Montelukast Viatris
4. Possible side effects
5. How to store Montelukast Viatris
6. Contents of the pack and other information

#### **1. What Montelukast Viatris is and what it is used for**

Montelukast Viatris contains montelukast, which is a leukotriene receptor antagonist that blocks substances called leukotrienes.

Leukotrienes cause narrowing and swelling of airways in the lungs and also cause allergy symptoms. By blocking leukotrienes, Montelukast Viatris improves asthma symptoms, helps control asthma and improves seasonal allergy symptoms (also known as hay fever or seasonal allergic rhinitis).

Your doctor has prescribed Montelukast Viatris to treat asthma, preventing your asthma symptoms during the day and night.

- Montelukast Viatris is used for the treatment of adults and adolescents 15 years of age and older who are not adequately controlled on their medication and need additional therapy.
- Montelukast Viatris also helps prevent the narrowing of airways triggered by exercise.
- In those asthmatic patients in whom Montelukast Viatris is indicated in asthma, it can also provide symptomatic relief of seasonal allergic rhinitis.

Your doctor will determine how Montelukast Viatris should be used depending on the symptoms and severity of your asthma.

Montelukast Viatris 10 mg Tablets are used in adults and adolescents 15 years of age and over only.

#### **What is asthma?**

Asthma is a long-term disease.

Asthma includes:

- difficulty breathing because of narrowed airways. This narrowing of airways worsens and improves in response to various conditions.

- Sensitive airways that react to many things, such as cigarette smoke, pollen, cold air, or exercise.
- Swelling (inflammation) in the lining of the airways.

Symptoms of asthma include: Coughing, wheezing, and chest tightness.

### **What are seasonal allergies?**

Seasonal allergies (also known as hay fever or seasonal allergic rhinitis) are an allergic response often caused by airborne pollens from trees, grasses and weeds. The symptoms of seasonal allergies typically may include: stuffy, runny, itchy nose; sneezing; watery, swollen, red, itchy eyes.

## **2. What you need to know before you take Montelukast Viatris**

Tell your doctor about any medical problems or allergies you have now or have had.

### **Do not take Montelukast Viatris:**

- if you are allergic to montelukast or any of the other ingredients of this medicine (listed in section 6).

### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Montelukast.

- If your asthma or breathing gets worse, tell your doctor immediately.
- Montelukast Viatris is not meant to treat acute asthma attacks. If an attack occurs, follow the instructions your doctor has given you. Always have your inhaled rescue medicine for asthma attacks with you. Talk to your doctor if you need more inhaled rescue medicine than usual for the acute asthma attacks.
- It is important that you take all asthma medications prescribed by your doctor. Montelukast Viatris should not be substituted for other asthma medications your doctor has prescribed for you.
- Any patient on anti-asthma medicines should be aware that if you develop a combination of symptoms such as a flu-like illness, pins and needles or numbness of arms or legs, worsening of lung (pulmonary) symptoms, and/or rash, you should consult your doctor.
- You should not take acetyl-salicylic acid (aspirin) or anti-inflammatory medicines (also known as non-steroidal anti-inflammatory drugs or NSAIDs) if they make your asthma worse.

**Various neuropsychiatric events (for example behaviour and mood-related changes, depression and suicidality) have been reported in patients of all ages treated with montelukast (see section 4). If you develop such symptoms while taking montelukast, you should contact your doctor.**

### **Children and adolescents**

Do not give this medicine to children less than 15 years of age.

There are different form(s) of this medicine available for paediatric patients under 18 years of age based on age range.

### **Other medicines and Montelukast Viatris**

Some medicines may affect how Montelukast Viatris works, or Montelukast Viatris may affect how other medicines work.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including those obtained without a prescription.

Tell your doctor if you are taking the following medicines before starting Montelukast Viatris:

- phenobarbital (used for treatment of epilepsy)
- phenytoin (used for treatment of epilepsy)
- rifampicin (used to treat tuberculosis and some other infections)
- gemfibrozil (used for treatment of high lipid levels in plasma)

### **Pregnancy, breast-feeding and fertility**

#### *Pregnancy*

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. Your doctor will assess whether you can take Montelukast Viatris during this time.

#### *Breast-feeding*

It is not known if Montelukast Viatris appears in breast milk. You should talk to your doctor before taking Montelukast Viatris if you are breast-feeding or intend to breast-feed.

### **Driving and using machines**

Montelukast Viatris is not expected to affect your ability to drive a car or operate machinery. However, individual responses to medication may vary. Certain side effects (such as dizziness and drowsiness) that have been reported with Montelukast Viatris may affect some patient's ability to drive or operate machinery.

### **Montelukast Viatris contains sunset yellow and sodium**

Sunset yellow (E110), may cause allergic reactions.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

## **3. How to take Montelukast Viatris**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- You should only take one tablet of Montelukast Viatris once a day as prescribed by your doctor.
- It should be taken even when you have no symptoms or have an acute asthma attack.

### **For adults (including the elderly) and adolescents of 15 years of age and older:**

The recommended dose is:

- One 10 mg tablet to be taken daily in the evening.

### **Use in children and adolescents**

Montelukast Viatris is not appropriate for children and adolescents under 15 years of age.

If you are taking Montelukast Viatris, be sure that you do not take any other products that contain the same active ingredient, montelukast.

This medicine is for oral use.

You can take Montelukast with or without food.

**If you take more Montelukast Viatris than you should**

Contact your doctor immediately for advice.

There were no side effects reported in the majority of over dosage reports. The most frequently occurring symptoms reported with over dosage included abdominal pain, sleepiness, thirst, headache, vomiting, and hyperactivity.

**If you forget to take Montelukast Viatris**

Try to take Montelukast Viatris as prescribed. However, if you miss a dose, just resume the usual schedule of one tablet once daily.

Do not take a double dose to make up for a forgotten dose.

**If you stop taking Montelukast Viatris**

Montelukast Viatris can treat your asthma only if you continue to take it.

It is important to continue taking Montelukast Viatris for as long as your doctor prescribes. It will help control your asthma.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

In clinical studies with montelukast 10 mg film-coated tablets, the most commonly reported side effects (may affect up to 1 in 10 people) thought to be related to montelukast were:

- abdominal pain
- headache

These were usually mild and occurred at a greater frequency in patients treated with montelukast than placebo (a pill containing no medication).

**Serious side effects**

**Talk with your doctor immediately** if you notice any of the following side effects, which may be serious, and for which you may need urgent medical treatment.

Uncommon: the following may affect up to 1 in 100 people

- allergic reactions including swelling of the face, lips, tongue, and/or throat which may cause difficulty in breathing or swallowing
- behaviour and mood related changes: agitation including aggressive behaviour or hostility, depression
- seizure

Rare: the following may affect up to 1 in 1,000 people

- increased bleeding tendency
- tremor
- palpitations

Very rare: the following may affect up to 1 in 10,000 people

- combination of symptoms such as flu-like illness, pins and needles or numbness of arms and legs, worsening of pulmonary symptoms and/or rash (Churg-Strauss syndrome) (see section 2)
- low blood platelet count
- behaviour and mood related changes: hallucinations, disorientation, suicidal thoughts and actions
- swelling (inflammation) of the lungs
- severe skin reactions (erythema multiforme) that may occur without warning
- inflammation of the liver (hepatitis)

#### **Other side effects while the medicine has been on the market**

Very common: the following may affect more than 1 in 10 people

- upper respiratory infection

Common: the following may affect up to 1 in 10 people

- diarrhoea, nausea, vomiting
- rash
- fever
- elevated liver enzymes

Uncommon: the following may affect up to 1 in 100 people

- behaviour and mood related changes: dream abnormalities, including nightmares, trouble sleeping, sleepwalking, irritability, feeling anxious, restlessness
- dizziness, drowsiness, pins and needles/numbness
- nosebleed
- dry mouth, indigestion
- bruising, itching, hives
- joint or muscle pain, muscle cramps
- bedwetting in children
- weakness/tiredness, feeling unwell, swelling

Rare: the following may affect up to 1 in 1,000 people

- behaviour and mood related changes: disturbance in attention, memory impairment, uncontrolled muscle movements

Very rare: the following may affect up to 1 in 10,000 people

- tender red lumps under the skin, most commonly on your shins (erythema nodosum)
- behaviour and mood related changes: obsessive-compulsive symptoms, stuttering

#### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA

Pharmacovigilance. Website: [www.hpra.ie](http://www.hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

### **5. How to store Montelukast Viatris**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label, carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Montelukast Viatris contains**

- The active substance is montelukast.
- Each tablet contains 10 mg of montelukast as montelukast sodium.
- The other ingredients are:  
**Core:** microcrystalline cellulose, mannitol, croscarmellose sodium, magnesium stearate, sodium laurilsulfate, colloidal anhydrous silica.

**Film-coating:** polydextrose, titanium dioxide (E171), hypromellose, Triacetin, macrogol, indigo carmine aluminium lake (E132), sunset yellow aluminium lake (E110) (see section 2 “**Montelukast contains sunset yellow(E110)**”).

### **What Montelukast Viatris looks like and contents of the pack**

Blue, round, film-coated, tablets with sides that curve outwards and sloping edges marked with ‘MO’ over ‘10’ on one side and ‘M’ on the other side.

Montelukast Viatris 10 mg tablets are available in blisters of: 10, 14, 20, 28, 30, 50, 56, 60, 84, 90 and 100 tablets or in perforated unit dose blisters of 28 x 1 or in bottles with a silica gel desiccant containing 28, 30, 56, 60, 84, 90, 100, 112, 120, 180 and 500 tablets.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Viatris Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland

### **Manufacturer(s)**

McDermott Laboratories Ltd, t/a Gerard Laboratories, 35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13, Ireland.

Mylan Hungary Kft, H-2900 Komárom, Mylan utca 1, Hungary.

### **This medicinal product is authorised in the Member States of the EEA under the following names:**

Czech Republic: Montelukast Viatris

Denmark: Montelukast Viatris

France: MONTELUKAST Viatris 10 mg, comprimé pelliculé

Germany: Montelukast dura 10 mg Filmtabletten

Ireland: Montelukast Viatris 10 mg film-coated tablets

Italy: Montelukast Mylan Generics

Portugal: Montelucaste Mylan

Slovakia: Montelukast Viatris 10 mg

Spain: Montelukast Viatris 10 mg comprimidos recubiertos con película EFG

Sweden: Montelukast Viatris

The Netherlands: Montelukast Viatris 10 mg, filmomhulde tabletten

United Kingdom (Northern Ireland): Montelukast 10 mg film-coated tablets

**This leaflet was last revised in 01/2025.**