

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Efavirenz Mylan 600 mg Film-coated Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 600 mg of efavirenz.

Excipient with known effect:

Each film-coated tablet contains 128 mg of lactose monohydrate.

For the full list of excipients, see section 6.1

3 PHARMACEUTICAL FORM

Film-coated tablet

Yellow capsule shaped, biconvex, bevelled-edge film-coated tablet debossed with "M" on one side and "EV6" on the other.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Efavirenz Mylan is indicated in antiviral combination treatment of human immunodeficiency virus-1 (HIV-1) infected adults, adolescents and children 3 months of age and older and weighing at least 3.5 kg.

Efavirenz has not been adequately studied in patients with advanced HIV disease, namely in patients with CD4 counts < 50 cells/mm³, or after failure of protease inhibitor (PI) containing regimens. Although cross-resistance of efavirenz with PIs has not been documented, there are at present insufficient data on the efficacy of subsequent use of PI based combination therapy after failure of regimens containing efavirenz.

For a summary of clinical and pharmacodynamic information, see section 5.1.

4.2 Posology and method of administration

Therapy should be initiated by a physician experienced in the management of HIV infection.

Posology

Efavirenz must be given in combination with other antiretroviral medicines (see section 4.5).

In order to improve the tolerability of nervous system adverse reactions, bedtime dosing is recommended (see section 4.8).

Adults and adolescents over 40 kg: The recommended dose of efavirenz in combination with nucleoside analogue reverse transcriptase inhibitors (NRTIs) with or without a PI (see section 4.5) is 600 mg orally, once daily.

Efavirenz film-coated tablets are not suitable for children weighing less than 40 kg. Other efavirenz formulations are available in the market for these patients.

Dose adjustment: If efavirenz is co-administered with voriconazole, the voriconazole maintenance dose must be increased to 400 mg every 12 hours and the efavirenz dose must be reduced by 50%, i.e., to 300 mg once daily using other efavirenz formulations available in the market. When treatment with voriconazole is stopped, the initial dose of efavirenz should be restored (see section 4.5).

If Efavirenz is co-administered with rifampicin to patients weighing 50 kg or more, an increase in the dose of efavirenz to 800 mg/day may be considered using available efavirenz formulations (see section 4.5).

Special populations

Renal impairment: The pharmacokinetics of efavirenz have not been studied in patients with renal insufficiency; however, less than 1% of an efavirenz dose is excreted unchanged in the urine, so the impact of renal impairment on efavirenz elimination should be minimal (see section 4.4).

Hepatic impairment: Patients with mild liver disease may be treated with their normally recommended dose of efavirenz. Patients should be monitored carefully for dose-related adverse reactions, especially nervous system symptoms (see sections 4.3 and 4.4).

Paediatric population

The safety and efficacy of efavirenz in children below the age of 3 months or weighing less than 3.5 kg have not been established. No data are available.

Method of administration

It is recommended that Efavirenz Mylan be taken on an empty stomach. The increased efavirenz concentrations observed following administration with food may lead to an increase in frequency of adverse reactions (see sections 4.4 and 5.2).

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Patients with severe hepatic impairment (Child Pugh Grade C) (see section 5.2).

Co-administration with terfenadine, astemizole, cisapride, midazolam, triazolam, pimozide, bepridil, or ergot alkaloids (for example, ergotamine, dihydroergotamine, ergonovine, and methylergonovine) because competition for CYP3A4 by efavirenz could result in inhibition of metabolism and create the potential for serious and/or life-threatening adverse reactions (for example, cardiac arrhythmias, prolonged sedation or respiratory depression) (see section 4.5).

Herbal preparations containing St. John's wort (*Hypericum perforatum*) due to the risk of decreased plasma concentrations and reduced clinical effects of efavirenz (see section 4.5).

4.4 Special warnings and precautions for use

Efavirenz must not be used as a single agent to treat HIV or added on as a sole agent to a failing regimen. Resistant virus emerges rapidly when efavirenz is administered as monotherapy. The choice of new antiretroviral agent(s) to be used in combination with efavirenz should take into consideration the potential for viral cross-resistance (see section 5.1).

Co-administration of efavirenz with the fixed combination tablet containing efavirenz, emtricitabine, and tenofovir disoproxil fumarate is not recommended unless needed for dose adjustment (for example, with rifampicin).

Concomitant use of Ginkgo biloba extracts is not recommended (see section 4.5).

When prescribing medicinal products concomitantly with Efavirenz, physicians should refer to the corresponding Summary of Product Characteristics.

While effective viral suppression with antiretroviral therapy has been proven to substantially reduce the risk of sexual transmission, a residual risk cannot be excluded. Precautions to prevent transmission should be taken in accordance with national guidelines.

If any antiretroviral medicinal product in a combination regimen is interrupted because of suspected intolerance, serious consideration should be given to simultaneous discontinuation of all antiretroviral medicinal products. The antiretroviral medicinal products should be restarted at the same time upon resolution of the intolerance symptoms. Intermittent monotherapy and sequential reintroduction of antiretroviral agents is not advisable because of the increased potential for selection of resistant virus.

Rash: mild to moderate rash has been reported in clinical studies with efavirenz and usually resolves with continued therapy. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. Severe rash associated with blistering, moist desquamation or ulceration has been reported in less than 1% of patients treated with efavirenz. The incidence of erythema multiforme or Stevens-Johnson syndrome was approximately 0.1%. Efavirenz must be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement or fever. If therapy with efavirenz is discontinued, consideration should also be given to interrupting therapy with other antiretroviral agents to avoid development of resistant virus (see section 4.8).

Experience with efavirenz in patients who discontinued other antiretroviral agents of the NNRTI class is limited (see section 4.8). Efavirenz is not recommended for patients who have had a life-threatening cutaneous reaction (e.g., Stevens-Johnson syndrome) while taking another NNRTI.

Psychiatric symptoms: psychiatric adverse reactions have been reported in patients treated with efavirenz. Patients with a prior history of psychiatric disorders appear to be at greater risk of these serious psychiatric adverse reactions. In particular, severe depression was more common in those with a history of depression. There have also been post-marketing reports of severe depression, death by suicide, delusions and psychosis-like behaviour. Patients should be advised that if they experience symptoms such as severe depression, psychosis or suicidal ideation, they should contact their doctor immediately to assess the possibility that the symptoms may be related to the use of efavirenz, and if so, to determine whether the risks of continued therapy outweigh the benefits (see section 4.8).

Nervous system symptoms: symptoms including, but not limited to, dizziness, insomnia, somnolence, impaired concentration and abnormal dreaming are frequently reported adverse reactions in patients receiving efavirenz 600 mg daily in clinical studies (see section 4.8). Nervous system symptoms usually begin during the first one or two days of therapy and generally resolve after the first 2 - 4 weeks. Patients should be informed that if they do occur, these common symptoms are likely to improve with continued therapy and are not predictive of subsequent onset of any of the less frequent psychiatric symptoms.

Seizures: convulsions have been observed in adult and paediatric patients receiving efavirenz, generally in the presence of known medical history of seizures. Patients who are receiving concomitant anticonvulsant medicinal products primarily metabolised by the liver, such as phenytoin, carbamazepine and phenobarbital, may require periodic monitoring of plasma levels. In a drug interaction study, carbamazepine plasma concentrations were decreased when carbamazepine was co-administered with efavirenz (see section 4.5). Caution must be taken in any patient with a history of seizures.

Hepatic events: a few of the postmarketing reports of hepatic failure occurred in patients with no pre-existing hepatic disease or other identifiable risk factors (see section 4.8). Liver enzyme monitoring should be considered for patients without pre-existing hepatic dysfunction or other risk factors.

Effect of food: the administration of Efavirenz with food may increase efavirenz exposure (see section 5.2) and may lead to an increase in the frequency of adverse reactions (see section 4.8). It is recommended that Efavirenz be taken on an empty stomach, preferably at bedtime.

Immune Reactivation Syndrome: in HIV infected patients with severe immune deficiency at the time of institution of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic pathogens may arise and cause serious clinical conditions, or aggravation of symptoms. Typically, such reactions have been observed within the first few weeks or months of initiation of CART. Relevant examples are cytomegalovirus retinitis, generalised and/or focal mycobacterial infections, and pneumonia caused by *Pneumocystis jiroveci* (formerly known as *Pneumocystis carinii*). Any inflammatory symptoms should be evaluated and treatment instituted when necessary. Autoimmune disorders (such as Graves' disease) have also been reported to occur in the setting of immune reactivation; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment.

Weight and metabolic parameters: an increase in weight and in levels of blood lipids and glucose may occur during antiretroviral therapy. Such changes may in part be linked to disease control and lifestyle. For lipids, there is in some cases evidence for a treatment effect, while for weight gain there is no strong evidence relating this to any particular treatment. For monitoring of blood lipids and glucose reference is made to established HIV treatment guidelines. Lipid disorders should be managed as clinically appropriate.

Osteonecrosis: although the aetiology is considered to be multifactorial (including corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported particularly in patients with advanced HIV-disease and/or long-term exposure to combination antiretroviral

therapy (CART). Patients should be advised to seek medical advice if they experience joint aches and pain, joint stiffness or difficulty in movement.

Special populations:

Liver disease: Efavirenz is contraindicated in patients with severe hepatic impairment (see sections 4.3 and 5.2) and not recommended in patients with moderate hepatic impairment because of insufficient data to determine whether dose adjustment is necessary. Because of the extensive cytochrome P450-mediated metabolism of efavirenz and limited clinical experience in patients with chronic liver disease, caution must be exercised in administering efavirenz to patients with mild hepatic impairment. Patients should be monitored carefully for dose related adverse reactions, especially nervous system symptoms. Laboratory tests should be performed to evaluate their liver disease at periodic intervals (see section 4.2).

The safety and efficacy of efavirenz has not been established in patients with significant underlying liver disorders. Patients with chronic hepatitis B or C and treated with combination antiretroviral therapy are at increased risk for severe and potentially fatal hepatic adverse reactions. Patients with pre-existing liver dysfunction including chronic active hepatitis have an increased frequency of liver function abnormalities during combination antiretroviral therapy and should be monitored according to standard practice. If there is evidence of worsening liver disease or persistent elevations of serum transaminases to greater than 5 times the upper limit of the normal range, the benefit of continued therapy with efavirenz needs to be weighed against the potential risks of significant liver toxicity. In such patients, interruption or discontinuation of treatment must be considered (see section 4.8).

In patients treated with other medicinal products associated with liver toxicity, monitoring of liver enzymes is also recommended. In case of concomitant antiviral therapy for hepatitis B or C, please refer also to the relevant product information for these medicinal products.

Renal insufficiency: the pharmacokinetics of efavirenz have not been studied in patients with renal insufficiency; however, less than 1% of an efavirenz dose is excreted unchanged in the urine, so the impact of renal impairment on efavirenz elimination should be minimal (see section 4.2). There is no experience in patients with severe renal failure and close safety monitoring is recommended in this population.

Elderly patients: insufficient numbers of elderly patients have been evaluated in clinical studies to determine whether they respond differently than younger patients.

Paediatric population:

Efavirenz has not been evaluated in children below 3 months of age or who weigh less than 3.5 kg. Therefore, efavirenz should not be given to children less than 3 months of age. Efavirenz film-coated tablets are not suitable for children weighing less than 40 kg.

Rash was reported in 59 of 182 children (32%) treated with efavirenz and was severe in six patients. Prophylaxis with appropriate antihistamines prior to initiating therapy with efavirenz in children may be considered.

Lactose: patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. Individuals with these conditions may take efavirenz oral solution, which is free from lactose.

4.5 Interaction with other medicinal products and other forms of interactions

Efavirenz is an *in vivo* inducer of CYP3A4, CYP2B6 and UGT1A1. Compounds that are substrates of these enzymes may have decreased plasma concentrations when co-administered with efavirenz. *In vitro* efavirenz is also an inhibitor of CYP3A4. Theoretically, efavirenz may therefore initially increase the exposure to CYP3A4 substrates and caution is warranted for CYP3A4 substrates with narrow therapeutic index (see section 4.4). Efavirenz may be an inducer of CYP2C19 and CYP2C9; however, inhibition has also been observed *in vitro* and the net effect is of co-administration with substrates of these enzymes is not clear (see section 5.2).

Efavirenz exposure may be increased when given with medicinal products (for example, ritonavir) or food (for example, grapefruit juice), which inhibit CYP3A4 or CYP2B6 activity. Compounds or herbal preparations (for example Ginkgo biloba extracts and St. John's wort) which induce these enzymes may give rise to decreased plasma concentrations of efavirenz. Concomitant use of St. John's wort is contraindicated (see section 4.3). Concomitant use of Ginkgo biloba extracts is not recommended (see section 4.4).

Paediatric population

Interaction studies have only been performed in adults.

Contraindications of concomitant use

Efavirenz must not be administered concurrently with terfenadine, astemizole, cisapride, midazolam, triazolam, pimozide, bepridil, or ergot alkaloids (for example, ergotamine, dihydroergotamine, ergonovine, and methylergonovine), since inhibition of their metabolism may lead to serious, life-threatening events (see section 4.3).

St. John's wort (Hypericum perforatum): Co-administration of efavirenz and St. John's wort or herbal preparations containing St. John's wort is contraindicated. Plasma levels of efavirenz can be reduced by concomitant use of St. John's wort due to induction of drug metabolising enzymes and/or transport proteins by St. John's wort. If a patient is already taking St. John's wort, stop St. John's wort, check viral levels and if possible efavirenz levels. Efavirenz levels may increase on stopping St. John's wort and the dose of efavirenz may need adjusting. The inducing effect of St. John's wort may persist for at least 2 weeks after cessation of treatment (see section 4.3).

Other interactions

Interactions between efavirenz and protease inhibitors, antiretroviral agents other than protease inhibitors and other non-antiretroviral medicinal products are listed in Table 1 below (increase is indicated as "↑", decrease as "↓", no change as "↔", and once every 8 or 12 hours as "q8h" or "q12h"). If available, 90% or 95% confidence intervals are shown in parentheses. Studies were conducted in healthy subjects unless otherwise noted.

Table 1: Interactions between efavirenz and other medicinal products in adults

| Medicinal product by therapeutic areas (dose) | Effects on drug levels Mean percent change in AUC, C_{max} , C_{min} with confidence intervals if available ^a (mechanism) | Recommendation concerning co-administration with efavirenz |
|--|---|---|
| ANTI-INFECTIVES | | |
| HIV antivirals | | |
| Protease inhibitors (PI) | | |
| Atazanavir/ritonavir/Efavirenz (400 mg once daily/100 mg once daily/600 mg once daily, all administered with food) | Atazanavir (pm): AUC: ↔* (↓9 to ↑10) C_{max} : ↑17%* (↑8 to ↑27) C_{min} : ↓42%* (↓31 to ↓51) | Co-administration of efavirenz with atazanavir/ritonavir is not recommended. If the co-administration of atazanavir with an NNRTI is required, an increase in the dose of both atazanavir and ritonavir to 400 mg and 200 mg, respectively, in combination with efavirenz could be considered with close clinical monitoring. |
| Atazanavir/ritonavir/Efavirenz (400 mg once daily/200 mg once daily/600 mg once daily, all administered with food) | Atazanavir (pm): AUC: ↔*/** (↓ 10 to ↑26) C_{max} : ↔*/** (↓ 5 to ↑26) C_{min} : ↑12%*/** (↓ 16 to ↑49) (CYP3A4 induction). * When compared to atazanavir 300 mg/ritonavir 100 mg once daily in the evening without efavirenz. This decrease in atazanavir C_{min} might negatively impact the efficacy of atazanavir. | |

| | | |
|---|--|---|
| | ** based on historical comparison | |
| Darunavir/ritonavir/Efavirenz (300 mg twice daily*/100 mg twice daily/600 mg once daily) *lower than recommended doses; similar findings are expected with recommended doses. | Darunavir: AUC: ↓13% C _{min} : ↓31% C _{max} : ↓15% (CYP3A4 induction) Efavirenz: AUC: ↑ 21% C _{min} : ↑ 17% C _{max} : ↑ 15% (CYP3A4 inhibition) | Efavirenz in combination with darunavir/ritonavir 800/100 mg once daily may result in suboptimal darunavir C _{min} . If Efavirenz is to be used in combination with darunavir/ritonavir, the darunavir/ritonavir 600/100 mg twice daily regimen should be used. This combination should be used with caution. See also ritonavir row below. |
| Fosamprenavir/ritonavir/ Efavirenz (700 mg twice daily/100 mg twice daily/600 mg once daily) | No clinically significant pharmacokinetic interaction | No dose adjustment is necessary for any of these medicinal products. See also ritonavir row below. |
| Fosamprenavir/Nelfinavir/ Efavirenz | Interaction not studied | No dose adjustment is necessary for any of these medicinal products. |
| Fosamprenavir/Saquinavir/ Efavirenz | Interaction not studied | Not recommended as the exposure to both PIs is expected to be significantly decreased. |
| Indinavir/Efavirenz (800 mg q8h/200 mg once daily) | Indinavir: AUC : ↓31% (↓ 8 to ↓ 47) C _{min} : ↓ 40% A similar reduction in indinavir exposures was observed when indinavir 1000 mg q8h was given with efavirenz 600 mg daily. (CYP3A4 induction) Efavirenz: No clinically significant pharmacokinetic interaction | While the clinical significance of decreased indinavir concentrations has not been established, the magnitude of the observed pharmacokinetic interaction should be taken into consideration when choosing a regimen containing both efavirenz and indinavir. No dose adjustment is necessary for efavirenz when given with indinavir or indinavir/ritonavir. See also ritonavir row below. |
| Indinavir/ritonavir/Efavirenz (800 mg twice daily/100 mg twice daily/600 mg once daily) | Indinavir: AUC: ↓ 25% (↓ 16 to ↓32) _b C _{max} : ↓17% (↓ 6 to ↓26) _b C _{min} : ↓50% (↓ 40 to ↓59) _b Efavirenz: No clinically significant pharmacokinetic interaction The geometric mean C _{min} for indinavir (0.33 | |

| | | |
|---|---|---|
| | <p>mg/l) when given with ritonavir and efavirenz was higher than the mean historical C_{min} (0.15 mg/l) when indinavir was given alone at 800 mg q8h. In HIV-1 infected patients (n = 6), the pharmacokinetics of indinavir and efavirenz were generally comparable to these uninfected volunteer data.</p> | |
| <p>Lopinavir/ritonavir soft capsules or oral solution/Efavirenz</p> <p>Lopinavir/ritonavir tablets/ Efavirenz (400/100 mg twice daily/ 600 mg once daily) (500/125 mg twice daily/ 600 mg once daily)</p> | <p>Substantial decrease in lopinavir exposure.</p> <p>Lopinavir concentrations: ↓30-40% Lopinavir concentrations: similar to lopinavir/ritonavir 400/100 mg twice daily without efavirenz</p> | <p>With efavirenz, an increase of the lopinavir/ritonavir soft capsule or oral solution doses by 33% should be considered (4 capsules/~6.5 ml twice daily instead of 3 capsules/5 ml twice daily). Caution is warranted since this dose adjustment might be insufficient in some patients. The dose of lopinavir/ritonavir tablets should be increased to 500/125 mg twice daily when co-administered with efavirenz 600 mg once daily. See also ritonavir row below.</p> |
| <p>Nelfinavir/Efavirenz (750 mg q8h/600 mg once daily)</p> | <p>Nelfinavir: AUC: ↑ 20% (↑ 8 to ↑ 34) C_{max}: ↑ 21% (↑ 10 to ↑ 33) The combination was generally well tolerated.</p> | <p>No dose adjustment is necessary for either medicinal product.</p> |
| <p>Ritonavir/Efavirenz (500 mg twice daily/600 mg once daily)</p> | <p>Ritonavir: Morning AUC: ↑ 18% (↑ 6 to ↑ 33) Evening AUC: ↔ Morning C_{max}: ↑ 24% (↑ 12 to ↑ 38) Evening C_{max}: ↔ Morning C_{min}: ↑ 42% (↑ 9 to ↑ 86)^b Evening C_{min}: ↑ 24% (↑ 3 to ↑ 50)^b Efavirenz: AUC: ↑ 21% (↑ 10 to ↑ 34) C_{max}: ↑ 14% (↑ 4 to ↑ 26) C_{min}: ↑ 25% (↑ 7 to ↑ 46) ^b (inhibition of</p> | <p>When using efavirenz with low-dose ritonavir, the possibility of an increase in the incidence of efavirenz-associated adverse events should be considered, due to possible pharmacodynamic interaction.</p> |

| | | |
|---|--|---|
| | CYP-mediated oxidative metabolism) When efavirenz was given with ritonavir 500 mg or 600 mg twice daily, the combination was not well tolerated (for example, dizziness, nausea, paraesthesia and elevated liver enzymes occurred). Sufficient data on the tolerability of efavirenz with low-dose ritonavir (100 mg, once or twice daily) are not available. | |
| Saquinavir/ritonavir/Efavirenz | Interaction not studied. | No data are available to make a dose recommendation. See also ritonavir row above. Use of efavirenz in combination with saquinavir as the sole protease inhibitor is not recommended. |
| CCR5 antagonist | | |
| Maraviroc/Efavirenz (100 mg twice daily/600 mg once daily) | Maraviroc: AUC ₁₂ : ↓45% (↓ 38 to ↓51) C _{max} : ↓51% (↓ 37 to ↓62) Efavirenz concentrations not measured, no effect is expected. | Refer to the Summary of Product Characteristics for the medicinal product containing maraviroc. |
| Integrase strand transfer inhibitor | | |
| Raltegravir/Efavirenz (400 mg single dose/ -) | Raltegravir: AUC: ↓36% C ₁₂ : ↓21% C _{max} : ↓36% (UGT1A1 induction) | No dose adjustment is necessary for raltegravir. |
| NRTIs and NNRTIs | | |
| NRTIs/Efavirenz | Specific interaction studies have not been performed with efavirenz and NRTIs other than lamivudine, zidovudine, and tenofovir disoproxil fumarate. Clinically significant interactions are not expected since the NRTIs are metabolised via a | No dose adjustment is necessary for either medicinal product. |

| | | |
|--|---|--|
| | different route than efavirenz and would be unlikely to compete for the same metabolic enzymes and elimination pathways. | |
| NNRTIs/Efavirenz | Interaction not studied. | Since use of two NNRTIs proved not beneficial in terms of efficacy and safety, co-administration of efavirenz and another NNRTI is not recommended. |
| Hepatitis C antivirals | | |
| Boceprevir/Efavirenz (800 mg 3 times daily/600 mg once daily) | Boceprevir: AUC: ↔ 19%* C _{max} : ↔ 8% C _{min} : ↓44% Efavirenz: AUC: ↔ 20%* C _{max} : ↔ 11% CYP3A induction – effect on boceprevir *0 – 8 hours No effect (↔) equals a decrease in mean ratio estimate of ≤ 20% or increase in mean ratio estimate of ≤ 25% | Plasma trough concentrations of boceprevir were decreased when administered with efavirenz. The clinical outcome of this observed reduction of boceprevir trough concentrations has not been directly assessed. |
| Telaprevir/Efavirenz (1,125 mg q8h/600 mg once daily) | Telaprevir (relative to 750 mg q8h): AUC: ↓ 18% (↓8 to ↓27) C _{max} : ↓14% (↓3 to ↓24) C _{min} : ↓25% (↓14 to ↓34) Efavirenz: AUC: ↓18% (↓10 to ↓26) C _{max} : ↓24% (↓15 to ↓32) C _{min} : ↓10% (↓1 to ↓19) (CYP3A induction by efavirenz) | If efavirenz and telaprevir are co-administered, telaprevir 1,125 mg every 8 hours should be used. |
| Simeprevir/Efavirenz (150 mg once daily /600 mg once daily) | Simeprevir: AUC: ↓71% (↓67 to ↓74) C _{max} : ↓51% (↓46 to ↓56) C _{min} : ↓91% (↓88 to ↓92) Efavirenz: AUC: ↔ | Concomitant administration of simeprevir with efavirenz resulted in significantly decreased plasma concentrations of simeprevir due to CYP3A induction by efavirenz, which may result in loss of therapeutic effect of simeprevir. Co-administration of simeprevir with efavirenz is not |

| | | |
|--|--|---|
| | <p>C_{max}: ↔ C_{min}: ↔ No effect (↔) equals a decrease in mean ratio estimate of ≤20% or increase in mean ratio estimate of ≤25% (CYP3A4 enzyme induction)</p> | recommended |
| Antibiotics | | |
| Azithromycin/Efavirenz (600 mg single dose/400 mg once daily) | No clinically significant pharmacokinetic interaction. | No dose adjustment is necessary for either medicinal product. |
| Clarithromycin/Efavirenz (500 mg q12h/400 mg once daily) | <p>Clarithromycin: AUC: ↓39% (↓ 30 to ↓46) C_{max}: ↓26% (↓ 15 to ↓35) Clarithromycin 14-hydroxymetabolite: AUC: ↑ 34% (↑ 18 to ↑ 53) C_{max}: ↑ 49% (↑ 32 to ↑ 69) Efavirenz: AUC: ↔ C_{max}: ↑ 11% (↑ 3 to ↑ 19) (CYP3A4 induction) Rash developed in 46% of uninfected volunteers receiving efavirenz and clarithromycin.</p> | The clinical significance of these changes in clarithromycin plasma levels is not known. Alternatives to clarithromycin (e.g. azithromycin) may be considered. No dose adjustment is necessary for efavirenz. |
| Other macrolide antibiotics (e.g., erythromycin)/Efavirenz | Interaction not studied. | No data are available to make a dose recommendation. |
| Antimycobacterials | | |
| Rifabutin/Efavirenz (300 mg once daily/600 mg once daily) | <p>Rifabutin: AUC: ↓38% (↓ 28 to ↓47) C_{max}: ↓32% (↓ 15 to ↓46) C_{min}: ↓45% (↓ 31 to ↓56) Efavirenz: AUC: ↔ C_{max}: ↔ C_{min}: ↓12% (↓ 24 to ↑ 1) (CYP3A4 induction)</p> | The daily dose of rifabutin should be increased by 50% when administered with efavirenz. Consider doubling the rifabutin dose in regimens where rifabutin is given 2 or 3 times a week in combination with efavirenz. The clinical effect of this dose adjustment has not been adequately evaluated. Individual tolerability and virological response should be considered when making the dose adjustment (see section 5.2). |
| Rifampicin/Efavirenz (600 mg once daily/600 mg once daily) | <p>Efavirenz: AUC: ↓26% (↓ 15 to ↓36)</p> | When taken with rifampicin in patients weighing 50 kg or greater, increasing efavirenz daily dose to 800 mg may |

| | | |
|---|---|--|
| | <p>C_{max} : ↓20% (↓ 11 to ↓28) C_{min} : ↓32% (↓ 15 to ↓46) (CYP3A4 and CYP2B6 induction)</p> | <p>provide exposure similar to a daily dose of 600 mg when taken without rifampicin. The clinical effect of this dose adjustment has not been adequately evaluated. Individual tolerability and virological response should be considered when making the dose adjustment (see section 5.2). No dose adjustment is necessary for rifampicin.</p> |
| Antifungals | | |
| <p>Itraconazole/Efavirenz (200 mg q12h/600 mg once daily)</p> | <p>Itraconazole: AUC: ↓39% (↓ 21 to ↓53) C_{max} : ↓37% (↓ 20 to ↓51) C_{min} : ↓44% (↓ 27 to ↓58) (decrease in itraconazole concentrations: CYP3A4 induction) Hydroxyitraconazole: AUC: ↓37% (↓ 14 to ↓55) C_{max} : ↓35% (↓ 12 to ↓52) C_{min} : ↓43% (↓ 18 to ↓60) Efavirenz: No clinically significant pharmacokinetic change.</p> | <p>Since no dose recommendation for itraconazole can be made, alternative antifungal treatment should be considered.</p> |
| <p>Posaconazole/Efavirenz --/400 mg once daily</p> | <p>Posaconazole: AUC: ↓50% C_{max} : ↓45% (UDP-G induction)</p> | <p>Concomitant use of posaconazole and efavirenz should be avoided unless the benefit to the patient outweighs the risk.</p> |
| <p>Voriconazole/Efavirenz (200 mg twice daily/400 mg once daily)</p> <p>Voriconazole/Efavirenz (400 mg twice daily/300 mg once daily)</p> | <p>Voriconazole: AUC: ↑77% C_{max} : ↓61% Efavirenz: AUC: ↑ 44% C_{max} : ↑ 38% Voriconazole: AUC: ↓7% (↓ 23 to ↑ 13) * C_{max} : ↑ 23% (↓ 1 to ↑ 53) * Efavirenz: AUC: ↑ 17% (↑ 6 to ↑ 29) **</p> | <p>When efavirenz is co-administered with voriconazole, the voriconazole maintenance dose must be increased to 400 mg twice daily and the efavirenz dose must be reduced by 50%, i.e., to 300 mg once daily. When treatment with voriconazole is stopped, the initial dose of efavirenz should be restored</p> |

| | | |
|---|--|--|
| | <p>C_{max}: ↔** *compared to 200 mg twice daily alone ** compared to 600 mg once daily alone (competitive inhibition of oxidative metabolism)</p> | |
| Fluconazole/Efavirenz (200 mg once daily/400 mg once daily) | No clinically significant pharmacokinetic interaction | No dose adjustment is necessary for either medicinal product. |
| Ketoconazole and other imidazole antifungals | Interaction not studied | No data are available to make a dose recommendation. |
| ANTIMALERIAL | | |
| Artemether/lumefantrine/ Efavirenz (20/120 mg tablet, 6 doses of 4 tablets each over 3 days/600mg once daily) | <p>Artemether: AUC: ↓ 51% C_{max}: ↓ 21% Dihydroartemisinin: AUC: ↓ 46% C_{max}: ↓ 38% Lumefantrine: AUC: ↓ 21% C_{max}: ↔ Efavirenz: AUC: ↓ 17% C_{max}: ↔ (CYP3A4 induction)</p> | Since decreased concentrations of artemether, dihydroartemisinin, or lumefantrine may result in a decrease of antimalarial efficacy, caution is recommended when efavirenz and artemether/lumefantrine tablets are coadministered. |
| Atovaquone and proguanil hydrochlorothiazide/Efavirenz (250/100 mg single dose/600 mg once daily) | Atovaquone: AUC: ↓ 75% (↓ 62 to ↓ 84) | Concomitant administration of atovaquone/proguanil with efavirenz should be avoided whenever possible. |
| | C_{max} : ↓ 44% (↓ 20 to ↓ 61) Proguanil: AUC: ↓ 43% (↓ 7 to ↓ 65) C_{max} : ↔ | |
| ACID REDUCING AGENTS | | |
| Aluminium hydroxide-magnesium hydroxide-simethicone antacid/Efavirenz (30 ml single dose/400 mg single dose) Famotidine/Efavirenz (40 mg single dose/400 mg single dose) | Neither aluminium/magnesium hydroxide antacids nor famotidine altered the absorption of efavirenz. | Co-administration of efavirenz with medicinal products that alter gastric pH would not be expected to affect efavirenz absorption. |
| ANTI-ANXIETY AGENTS | | |
| Lorazepam/Efavirenz (2 mg single dose/600 mg once daily) | <p>Lorazepam: AUC: ↑ 7% (↑ 1 to ↑ 14) C_{max}: ↑ 16% (↑ 2 to ↑ 32) These changes are not</p> | No dose adjustment is necessary for either medicinal product. |

| | | |
|--|---|--|
| | considered clinically significant. | |
| ANTICOAGULANTS | | |
| Warfarin/Efavirenz Acenocoumarol/Efavirenz | Interaction not studied. Plasma concentrations and effects of warfarin or acenocoumarol are potentially increased or decreased by efavirenz. | Dose adjustment of warfarin or acenocoumarol may be required. |
| ANTICONVULSANTS | | |
| Carbamazepine/Efavirenz (400 mg once daily/600 mg once daily) | Carbamazepine: AUC: ↓27% (↓ 20 to ↓33) C_{max} : ↓20% (↓ 15 to ↓24) C_{min} : ↓35% (↓ 24 to ↓44) Efavirenz: AUC: ↓36% (↓ 32 to ↓40) C_{max} : ↓21% (↓ 15 to ↓26) C_{min} : ↓47% (↓ 41 to ↓53) (decrease in carbamazepine concentrations: CYP3A4 induction; decrease in efavirenz concentrations: CYP3A4 and CYP2B6 induction) The steady-state AUC, C_{max} and C_{min} of the active carbamazepine epoxide metabolite remained unchanged. Co-administration of higher doses of either efavirenz or carbamazepine has not been studied. | No dose recommendation can be made. An alternative anticonvulsant should be considered. Carbamazepine plasma levels should be monitored periodically. |
| Phenytoin, Phenobarbital, and other anticonvulsants that are substrates of CYP450 isoenzymes | Interaction not studied. There is a potential for reduction or increase in the plasma concentrations of phenytoin, phenobarbital and other anticonvulsants that are substrates of | When efavirenz is co-administered with an anticonvulsant that is a substrate of CYP450 isoenzymes, periodic monitoring of anticonvulsant levels should be conducted. |

| | | |
|---|---|---|
| | CYP450 isoenzymes when co-administered with efavirenz. | |
| Valproic acid/Efavirenz (250 mg twice daily/600 mg once daily) | No clinically significant effect on efavirenz pharmacokinetics. Limited data suggest there is no clinically significant effect on valproic acid pharmacokinetics. | No dose adjustment is necessary for efavirenz. Patients should be monitored for seizure control. |
| Vigabatrin/Efavirenz Gabapentin/Efavirenz | Interaction not studied. Clinically significant interactions are not expected since vigabatrin and gabapentin are exclusively eliminated unchanged in the urine and are unlikely to compete for the same metabolic enzymes and elimination pathways as efavirenz. | No dose adjustment is necessary for any of these medicinal products. |
| ANTIDEPRESSANTS | | |
| Selective Serotonin Reuptake Inhibitors (SSRIs) | | |
| Sertraline/Efavirenz (50 mg once daily/600 mg once daily) | Sertraline: AUC: ↓39% (↓ 27 to ↓50) C_{max} : ↓29% (↓ 15 to ↓40) C_{min} : ↓46% (↓ 31 to ↓58) Efavirenz: AUC: ↔ C_{max} : ↑ 11% (↑ 6 to ↑ 16) C_{min} : ↔ (CYP3A4 induction) | Sertraline dose increases should be guided by clinical response. No dose adjustment is necessary for efavirenz. |
| Paroxetine/Efavirenz (20 mg once daily/600 mg once daily) | No clinically significant pharmacokinetic interaction | No dose adjustment is necessary for either medicinal product. |
| Fluoxetine/Efavirenz | Interaction not studied. Since fluoxetine shares a similar metabolic profile with paroxetine, i.e. a strong CYP2D6 inhibitory effect, a similar lack of interaction would be expected for fluoxetine. | No dose adjustment is necessary for either medicinal product. |

| NOREPINEPHRINE AND DOPAMINE REUPTAKE INHIBITOR | | |
|---|--|---|
| Bupropion/Efavirenz [150 mg single dose (sustained release)/600 mg once daily] | Bupropion: AUC: ↓55% (↓ 48 to ↓62) C_{max} : ↓34% (↓ 21 to ↓47) Hydroxybupropion: AUC: ↔ C_{max} : ↑ 50% (↑ 20 to ↑ 80) (CYP2B6 induction) | Increases in bupropion dosage should be guided by clinical response, but the maximum recommended dose of bupropion should not be exceeded. No dose adjustment is necessary for efavirenz. |
| ANTIHISTAMINES | | |
| Cetirizine/Efavirenz (10 mg single dose/600 mg once daily) | Cetirizine: AUC: ↔ C_{max} : ↓24% (↓ 18 to ↓30) These changes are not considered clinically significant. Efavirenz: No clinically significant pharmacokinetic interaction | No dose adjustment is necessary for either medicinal product. |
| CARDIOVASCULAR AGENTS | | |
| Calcium Channel Blockers | | |
| Diltiazem/Efavirenz (240 mg once daily/600 mg once daily) | Diltiazem: AUC: ↓69% (↓55 to ↓79) C_{max} : ↓60% (↓50 to ↓68) C_{min} : ↓63% (↓44 to ↓75) Desacetyl diltiazem: AUC: ↓75% (↓59 to ↓84) C_{max} : ↓64% (↓57 to ↓69) C_{min} : ↓62% (↓44 to ↓75) N- monodesmethyl diltiazem: AUC: ↓37% (↓17 to ↓52) C_{max} : ↓28% (↓7 to ↓44) C_{min} : ↓37% (↓17 to ↓52) Efavirenz: AUC: ↑ 11% (↑ 5 to ↑ 18) C_{max} : ↑ 16% (↑ 6 to ↑ 26) C_{min} : ↑ 13% (↑ 1 to ↑ 26) (CYP3A4 induction) The increase in efavirenz pharmacokinetic parameters is not considered clinically significant. | Dose adjustments of diltiazem should be guided by clinical response (refer to the Summary of Product Characteristics for diltiazem). No dose adjustment is necessary for efavirenz. |

| | | |
|--|--|---|
| Verapamil, Felodipine, Nifedipine and Nicardipine | Interaction not studied. When efavirenz is co-administered with a calcium channel blocker that is a substrate of the CYP3A4 enzyme, there is a potential for reduction in the plasma concentrations of the calcium channel blocker. | Dose adjustments of calcium channel blockers should be guided by clinical response (refer to the Summary of Product Characteristics for the calcium channel blocker). |
| LIPID LOWERING MEDICINAL PRODUCTS | | |
| HMG Co-A Reductase Inhibitors | | |
| Atorvastatin/Efavirenz (10 mg once daily/600 mg once daily) | Atorvastatin: AUC: ↓43% (↓34 to ↓50) C_{max} : ↓12% (↓1 to ↓26) 2-hydroxy atorvastatin: AUC: ↓35% (↓13 to ↓40) C_{max} : ↓13% (↓0 to ↓23) 4-hydroxy atorvastatin: AUC: ↓4% (↓0 to ↓31) C_{max} : ↓47% (↓9 to ↓51) Total active HMG Co-A reductase inhibitors: AUC: ↓34% (↓21 to ↓41) C_{max} : ↓20% (↓2 to ↓26) | Cholesterol levels should be periodically monitored. Dose adjustment of atorvastatin may be required (refer to the Summary of Product Characteristics for atorvastatin). No dose adjustment is necessary for efavirenz. |
| Pravastatin/Efavirenz (40 mg once daily/600 mg once daily) | Pravastatin: AUC: ↓40% (↓26 to ↓57) C_{max} : ↓18% (↓9 to ↓27) | Cholesterol levels should be periodically monitored. Dose adjustment of pravastatin may be required (refer to the Summary of Product Characteristics for pravastatin). No dose adjustment is necessary for efavirenz. |
| Simvastatin/Efavirenz (40 mg once daily/600 mg once daily) | Simvastatin: AUC: ↓69% (↓62 to ↓73) C_{max} : ↓76% (↓63 to ↓79) Simvastatin acid: AUC: ↓58% (↓39 to ↓68) C_{max} : ↓51% (↓32 to ↓58) Total active HMG Co-A reductase inhibitors: AUC: ↓60% (↓52 to ↓68) C_{max} : ↓62% (↓55 to ↓78) (CYP3A4 induction) Co-administration of efavirenz with atorvastatin, pravastatin, or simvastatin did not affect efavirenz AUC or | Cholesterol levels should be periodically monitored. Dose adjustment of simvastatin may be required (refer to the Summary of Product Characteristics for simvastatin). No dose adjustment is necessary for efavirenz. |

| | | |
|--|---|---|
| | C_{max} values. | |
| Rosuvastatin/Efavirenz | Interaction not studied. Rosuvastatin is largely excreted unchanged via the faeces, therefore interaction with efavirenz is not expected. | No dose adjustment is necessary for either medicinal product. |
| HORMONAL CONTRACEPTIVES | | |
| Oral: Ethinylestradiol + Norgestimate/ Efavirenz (0.035 mg + 0.25 mg once daily/600 mg once daily) | Ethinylestradiol: AUC: ↔ C_{max} : ↔ C_{min} : ↓8% (↑ 14 to ↓25) Norelgestromin (active metabolite): AUC: ↓64% (↓62 to ↓67) C_{max} : ↓46% (↓39 to ↓52) C_{min} : ↓82% (↓79 to ↓85) Levonorgestrel (active metabolite): AUC: ↓83% (↓79 to ↓87) C_{max} : ↓80% (↓77 to ↓83) C_{min} : ↓86% (↓80 to ↓90) (induction of metabolism) Efavirenz: no clinically significant interaction. The clinical significance of these effects is not known. | A reliable method of barrier contraception must be used in addition to hormonal contraceptives (see section 4.6). |
| Injection: Depomedroxyprogesterone acetate (DMPA)/Efavirenz (150 mg IM single dose DMPA) | In a 3-month drug interaction study, no significant differences in MPA pharmacokinetic parameters were found between subjects receiving efavirenz-containing antiretroviral therapy and subjects receiving no antiretroviral therapy. Similar results were found by other investigators, although the MPA plasma levels were more variable in the second study. In both studies, plasma progesterone levels for subjects receiving | Because of the limited information available, a reliable method of barrier contraception must be used in addition to hormonal contraceptives (see section 4.6). |

| | | |
|--|---|---|
| | efavirenz and DMPA remained low consistent with suppression of ovulation. | |
| Implant: Etonogestrel/Efavirenz | Interaction not studied. Decreased exposure of etonogestrel may be expected (CYP3A4 induction). There have been occasional post-marketing reports of contraceptive failure with etonogestrel in efavirenz-exposed patients. | A reliable method of barrier contraception must be used in addition to hormonal contraceptives (see section 4.6). |
| IMMUNOSUPPRESSANTS | | |
| Immunosuppressants metabolized by CYP3A4 (eg, cyclosporine, tacrolimus, sirolimus)/Efavirenz | Interaction not studied. Decreased exposure of the immunosuppressant may be expected (CYP3A4 induction). These immunosuppressants are not anticipated to affect exposure of efavirenz. | Dose adjustments of the immunosuppressant may be required. Close monitoring of immunosuppressant concentrations for at least 2 weeks (until stable concentrations are reached) is recommended when starting or stopping treatment with efavirenz. |
| OPIOIDS | | |
| Methadone/Efavirenz (stable maintenance, 35-100 mg once daily/600 mg once daily) | Methadone: AUC: ↓52% (↓33 to ↓66) C_{max} : ↓45% (↓25 to ↓59) (CYP3A4 induction) In a study of HIV infected intravenous drug users, co-administration of efavirenz with methadone resulted in decreased plasma levels of methadone and signs of opiate withdrawal. The methadone dose was increased by a mean of 22% to alleviate withdrawal symptoms. | Patients should be monitored for signs of withdrawal and their methadone dose increased as required to alleviate withdrawal symptoms. |
| Buprenorphine/naloxone/Efavirenz | Buprenorphine: AUC: ↓50% Norbuprenorphine: AUC: ↓71% | Despite the decrease in buprenorphine exposure, no patients exhibited withdrawal symptoms. Dose adjustment of buprenorphine or |

| | | |
|--|--|--|
| | Efavirenz: No clinically significant pharmacokinetic interaction. | efavirenz may not be necessary when co-administered. |
|--|--|--|

a - 90% confidence intervals unless otherwise noted.

b - 95% confidence intervals.

Other interactions: efavirenz does not bind to cannabinoid receptors. False-positive urine cannabinoid test results have been reported with some screening assays in uninfected and HIV-infected subjects receiving efavirenz. Confirmatory testing by a more specific method such as gas chromatography/mass spectrometry is recommended in such cases.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential: see below and section 5.3. Efavirenz should not be used during pregnancy, unless the patient's clinical condition requires such treatment. Women of childbearing potential should undergo pregnancy testing before initiation of efavirenz.

Contraception in males and females

Barrier contraception should always be used in combination with other methods of contraception (for example, oral or other hormonal contraceptives, see section 4.5). Because of the long half-life of efavirenz, use of adequate contraceptive measures for 12 weeks after discontinuation of efavirenz is recommended.

Pregnancy

There have been seven retrospective reports of findings consistent with neural tube defects, including meningomyelocele, all in mothers exposed to efavirenz-containing regimens (excluding any efavirenz-containing fixed-dose combination tablets) in the first trimester. Two additional cases (1 prospective and 1 retrospective) including events consistent with neural tube defects have been reported with the fixed-dose combination tablet containing efavirenz, emtricitabine and tenofovir disoproxil fumarate. A causal relationship of these events to the use of efavirenz has not been established, and the denominator is unknown. As neural tube defects occur within the first 4 weeks of foetal development (at which time neural tubes are sealed), this potential risk would concern women exposed to efavirenz during the first trimester of pregnancy.

As of July 2013, the Antiretroviral Pregnancy Registry (APR) has received prospective reports of 904 pregnancies with first trimester exposure to efavirenz-containing regimens, resulting in 766 live births. One child was reported to have a neural tube defect, and the frequency and pattern of other birth defects were similar to those

seen in children exposed to non-efavirenz-containing regimens, as well as those in HIV negative controls. The incidence of neural tube defects in the general population ranges from 0.5-1 case per 1,000 live births.

Malformations have been observed in foetuses from efavirenz-treated monkeys (see section 5.3).

Breast-feeding

Efavirenz has been shown to be excreted in human milk. There is insufficient information on the effects of efavirenz in newborns/infants. Risk to the infant cannot be excluded. Breast-feeding should be discontinued during treatment with efavirenz. It is recommended that HIV infected women do not breast feed their infants under any circumstances in order to avoid transmission of HIV.

Fertility

The effect of efavirenz on male and female fertility in rats has only been evaluated at doses that achieved systemic drug exposures equivalent to or below those achieved in humans given recommended doses of efavirenz. In these studies, efavirenz did not impair mating or fertility of male or female rats (doses up to 100 mg/kg/bid), and did not affect sperm or offspring of treated male rats (doses up to 200 mg/bid). The reproductive performance of offspring born to female rats given efavirenz was not affected.

4.7 Effects on ability to drive and use machines

Efavirenz may cause dizziness, impaired concentration, and/or somnolence. Patients should be instructed that, if they experience these symptoms, they should avoid potentially hazardous tasks such as driving or operating machinery.

4.8 Undesirable effects

Summary of the safety profile

Efavirenz has been studied in over 9,000 patients. In a subset of 1,008 adult patients who received 600 mg efavirenz daily in combination with PIs and/or NRTIs in controlled clinical studies, the most frequently reported adverse reactions of at least moderate severity reported in at least 5% of patients were rash (11.6%), dizziness (8.5%), nausea (8.0%), headache (5.7%) and fatigue (5.5%). The most notable adverse reactions associated with efavirenz are rash and nervous system symptoms. Nervous system symptoms usually begin soon after therapy onset and generally resolve after the first 2 - 4 weeks. Severe skin reactions such as Stevens-Johnson syndrome and erythema multiforme; psychiatric adverse reactions including severe depression, death by suicide, and psychosis like behaviour; and seizures have been reported in patients treated with efavirenz. The administration of efavirenz with food may

increase efavirenz exposure and may lead to an increase in the frequency of adverse reactions (see section 4.4).

The long-term safety profile of efavirenz containing regimens was evaluated in a controlled trial (006) in which patients received efavirenz + zidovudine + lamivudine (n = 412, median duration 180 weeks), efavirenz + indinavir (n = 415, median duration 102 weeks), or indinavir + zidovudine + lamivudine (n = 401, median duration 76 weeks). Long-term use of efavirenz in this study was not associated with any new safety concerns.

Tabulated list of adverse reactions

Adverse reactions of moderate or greater severity with at least possible relationship to treatment regimen (based on investigator attribution) reported in clinical trials of efavirenz at the recommended dose in combination therapy (n = 1,008) are listed below. Also listed in italics are adverse reactions observed post-marketing in association with efavirenz-containing antiretroviral treatment regimens. Frequency is defined using the following convention: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$) or very rare ($< 1/10,000$).

| | |
|---|---|
| Immune system disorders | |
| Uncommon | hypersensitivity |
| Metabolism and nutrition disorders | |
| Common | hypertriglyceridaemia ¹ |
| Uncommon | hypercholesterolaemia ¹ |
| Psychiatric disorders | |
| Common | abnormal dreams, anxiety, depression, insomnia ¹ |
| Uncommon | affect lability, aggression, confusional state, euphoric mood, hallucination, mania, paranoia, <i>psychosis</i> ² , suicide attempt, suicide ideation ¹ |
| Rare | <i>delusion</i> ³ , <i>neurosis</i> ³ , <i>completed suicide</i> ^{1,3} |
| Nervous system disorders | |
| Common | <i>cerebellar coordination and balance disturbances</i> ² , disturbance in attention (3.6%), dizziness (8.5%), headache (5.7%), somnolence (2.0%) ¹ |
| Uncommon | agitation, amnesia, ataxia, coordination abnormal, convulsions, thinking abnormal ¹ , <i>tremor</i> ² |
| Eye disorders | |
| Uncommon | vision blurred |
| Ear and labyrinth disorders | |
| Uncommon | <i>tinnitus</i> ² , vertigo |
| Vascular disorders | |

| | |
|---|---|
| Uncommon | <i>flushing</i> ² |
| Gastrointestinal disorders | |
| Common | abdominal pain, diarrhoea, nausea, vomiting |
| Uncommon | pancreatitis |
| Hepatobiliary disorders | |
| Common | aspartate amino transferase (AST) increased ¹ , alanine aminotransferase (ALT) increased ¹ , gamma-glutamyltransferase (GGT) increased ¹ |
| Uncommon | hepatitis acute |
| Rare | <i>hepatic failure</i> ^{1,3} |
| Skin and subcutaneous tissue disorders | |
| Very common | rash (11.6%) ¹ |
| Common | pruritus |
| Uncommon | erythema multiforme, Stevens-Johnson syndrome ¹ |
| Rare | <i>photoallergic dermatitis</i> ² |
| Reproductive system and breast disorders | |
| Uncommon | gynaecomastia |

| | |
|---|--|
| General disorders and administration site conditions | |
|---|--|

| | |
|--------|---------|
| Common | fatigue |
|--------|---------|

1. See section below. Description of selected adverse reactions for more details.

Description of selected adverse reactions

Information regarding post-marketing surveillance

2. These adverse reactions were identified through post-marketing surveillance; however, the frequencies were determined using data from 16 clinical trials (n = 3,969).

3. These adverse reactions were identified through post-marketing surveillance but not reported as drug-related events for efavirenz-treated patients in 16 clinical trials. The frequency category of rare was defined per *A Guideline on Summary of Product Characteristics (SmPC) (rev. 2, Sept 2009)* on the basis of an estimated upper bound of the 95% confidence interval for 0 events given the number of patients treated with efavirenz in these clinical trials (n = 3,969).

Rash: in clinical studies, 26% of patients treated with 600 mg of efavirenz experienced skin rash compared with 17% of patients treated in control groups. Skin rash was

considered treatment related in 18% of patients treated with efavirenz. Severe rash occurred in less than 1% of patients treated with efavirenz, and 1.7% discontinued therapy because of rash. The incidence of erythema multiforme or Stevens Johnson syndrome was approximately 0.1%.

Rashes are usually mild to moderate maculopapular skin eruptions that occur within the first two weeks of initiating therapy with efavirenz. In most patients rash resolves with continuing therapy with efavirenz within one month. Efavirenz can be reinitiated in patients interrupting therapy because of rash. Use of appropriate antihistamines and/or corticosteroids is recommended when efavirenz is restarted.

Experience with efavirenz in patients who discontinued other antiretroviral agents of the NNRTI class is limited. Reported rates of recurrent rash following a switch from nevirapine to efavirenz therapy, primarily based on retrospective cohort data from published literature, range from 13 to 18%, comparable to the rate observed in patients treated with efavirenz in clinical studies. (See section 4.4.)

Psychiatric symptoms: serious psychiatric adverse reactions have been reported in patients treated with efavirenz. In controlled trials, the frequency of specific serious psychiatric events were:

| | Efavirenz regimen (n = 1,008) | Control regimen (n = 635) |
|------------------------------|----------------------------------|------------------------------|
| - severe depression | 1.6 % | 0.6 % |
| - suicidal ideation | 0.6 % | 0.3 % |
| - non-fatal suicide attempts | 0.4 % | 0 % |
| - aggressive behaviour | 0.4 % | 0.3 % |
| - paranoid reactions | 0.4 % | 0.3 % |
| - manic reactions | 0.1 % | 0% |

Patients with a history of psychiatric disorders appear to be at greater risk of these serious psychiatric adverse reactions with frequencies ranging from 0.3% for manic reactions to 2.0% for both severe depression and suicidal ideation. There have also been post-marketing reports of death by suicide, delusions and psychosis-like behaviour.

Nervous system symptoms: in clinical controlled trials, frequently reported adverse reactions included, but were not limited to dizziness, insomnia, somnolence, impaired concentration and abnormal dreaming. Nervous system symptoms of moderate-to-severe intensity were experienced by 19% (severe 2%) of patients compared to 9% (severe 1%) of patients receiving control regimens. In clinical studies 2% of patients treated with efavirenz discontinued therapy due to such symptoms.

Nervous system symptoms usually begin during the first one or two days of therapy and generally resolve after the first 2 - 4 weeks. In a study of uninfected volunteers, a representative nervous system symptom had a median time to onset of 1 hour post-dose and a median duration of 3 hours. Nervous system symptoms may occur more frequently when efavirenz is taken concomitantly with meals possibly due to increased efavirenz plasma levels (see section 5.2). Dosing at bedtime seems to improve the tolerability of these symptoms and can be recommended during the first weeks of therapy and in patients who continue to experience these symptoms (see section 4.2). Dose reduction or splitting the daily dose has not been shown to provide benefit.

Analysis of long-term data showed that, beyond 24 weeks of therapy, the incidences of new-onset nervous system symptoms among efavirenz-treated patients were generally similar to those in the control arm.

Hepatic failure: A few of the post-marketing reports of hepatic failure, including cases in patients with no pre-existing hepatic disease or other identifiable risk factors, were characterised by a fulminant course, progressing in some cases to transplantation or death.

Immune Reactivation Syndrome: In HIV-infected patients with severe immune deficiency at the time of initiation of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic infections may arise. Autoimmune disorders (such as Graves' disease) have also been reported; however, the reported time to onset is more variable and these events can also occur many months after initiation of treatment (see section 4.4).

Osteonecrosis: cases of osteonecrosis have been reported, particularly in patients with generally acknowledged risk factors, advanced HIV disease or long-term exposure to combination antiretroviral therapy (CART). The frequency of this is unknown (see section 4.4).

Laboratory test abnormalities:

Liver enzymes: elevations of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) to greater than five times the upper limit of the normal range (ULN) were seen in 3% of 1,008 patients treated with 600 mg of efavirenz (5% - 8% after long-term treatment in study 006). Similar elevations were seen in patients treated with control regimens (5% after long-term treatment). Elevations of gamma-glutamyltransferase (GGT) to greater than five times ULN were observed in 4% of all patients treated with 600 mg of efavirenz and 1.5 - 2% of patients treated with control regimens (7% of efavirenz-treated patients and 3% of control-treated patients after long-term treatment). Isolated elevations of GGT in patients receiving

efavirenz may reflect enzyme induction. In the long-term study (006), 1% of patients in each treatment arm discontinued because of liver or biliary system disorders.

Amylase: in the clinical trial subset of 1,008 patients, asymptomatic increases in serum amylase levels greater than 1.5 times the upper limit of normal were seen in 10% of patients treated with efavirenz and 6% of patients treated with control regimens. The clinical significance of asymptomatic increases in serum amylase is unknown.

Metabolic parameters: weight and levels of blood lipids and glucose may increase during antiretroviral therapy (see section 4.4).

Paediatric population

Undesirable effects in children were generally similar to those of adult patients. Rash was reported more frequently in children (59 of 182 (32%) treated with efavirenz) and was more often of higher grade than in adults (severe rash was reported in 6 of 182 (3.3%) of children). Prophylaxis with appropriate antihistamines prior to initiating therapy with efavirenz in children may be considered.

Other special populations

Liver enzymes in hepatitis B or C co-infected patients: in the long-term data set from study 006, 137 patients treated with efavirenz-containing regimens (median duration of therapy, 68 weeks) and 84 treated with a control regimen (median duration 56 weeks) were seropositive at a screening for hepatitis B (surface antigen positive) and/or C (hepatitis C antibody positive). Among co-infected patients in study 006, elevations in AST to greater than five times ULN developed in 13% of efavirenz-treated patients and in 7% of control, and elevations in ALT to greater than five times ULN developed in 20% and 7% respectively. Among co-infected patients, 3% of those treated with efavirenz and 2% in the control arm discontinued because of liver disorders (see section 4.4).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

4.9 Overdose

Some patients accidentally taking 600 mg twice daily have reported increased nervous system symptoms. One patient experienced involuntary muscle contractions.

Treatment of overdose with efavirenz should consist of general supportive measures, including monitoring of vital signs and observation of the patient's clinical status. Administration of activated charcoal may be used to aid removal of unabsorbed efavirenz. There is no specific antidote for overdose with efavirenz. Since efavirenz is highly protein bound, dialysis is unlikely to remove significant quantities of it from blood.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antivirals for systemic use, non-nucleoside reverse transcriptase inhibitors.

ATC code: J05AG03

Mechanism of action:

Efavirenz is a NNRTI of HIV-1. Efavirenz is a non-competitive inhibitor of HIV-1 reverse transcriptase (RT) and does not significantly inhibit HIV-2 RT or cellular DNA polymerases (α , β , γ or δ).

Antiviral activity

The free concentration of efavirenz required for 90 to 95% inhibition of wild type or zidovudine-resistant laboratory and clinical isolates *in vitro* ranged from 0.46 to 6.8 nM in lymphoblastoid cell lines, peripheral blood mononuclear cells (PBMCs) and macrophage/monocyte cultures.

Resistance:

The potency of efavirenz in cell culture against viral variants with amino acid substitutions at positions 48, 108, 179, 181 or 236 in RT or variants with amino acid substitutions in the protease was similar to that observed against wild type viral strains. The single substitutions which led to the highest resistance to efavirenz in cell culture correspond to a leucine-to-isoleucine change at position 100 (L100I, 17 to 22-fold resistance) and a lysine-to-asparagine at position 103 (K103N, 18 to 33-fold resistance). Greater than 100-fold loss of susceptibility was observed against HIV variants expressing K103N in addition to other amino acid substitutions in RT.

K103N was the most frequently observed RT substitution in viral isolates from patients who experienced a significant rebound in viral load during clinical studies of efavirenz in combination with indinavir or zidovudine + lamivudine. This mutation was observed in 90% of patients receiving efavirenz with virological failure. Substitutions at RT positions 98, 100, 101, 108, 138, 188, 190 or 225 were also observed, but at lower frequencies, and often only in combination with K103N. The pattern of amino acid substitutions in RT associated with resistance to efavirenz was independent of the other antiviral medicines used in combination with efavirenz.

Cross resistance

Cross resistance profiles for efavirenz, nevirapine and delavirdine in cell culture demonstrated that the K103N substitution confers loss of susceptibility to all three NNRTIs. Two of three delavirdine-resistant clinical isolates examined were cross-resistant to efavirenz and contained the K103N substitution. A third isolate which carried a substitution at position 236 of RT was not cross-resistant to efavirenz.

Viral isolates recovered from PBMCs of patients enrolled in efavirenz clinical studies who showed evidence of treatment failure (viral load rebound) were assessed for susceptibility to NNRTIs. Thirteen isolates previously characterised as efavirenz-resistant were also resistant to nevirapine and delavirdine. Five of these NNRTI-resistant isolates were found to have K103N or a valine-to-isoleucine substitution at position 108 (V108I) in RT. Three of the efavirenz treatment failure isolates tested remained sensitive to efavirenz in cell culture and were also sensitive to nevirapine and delavirdine.

The potential for cross resistance between efavirenz and PIs is low because of the different enzyme targets involved. The potential for cross-resistance between efavirenz and NRTIs is low because of the different binding sites on the target and mechanism of action.

Clinical efficacy

Efavirenz has not been studied in controlled studies in patients with advanced HIV disease, namely with CD4 counts < 50 cells/mm³, or in PI or NNRTI experienced patients. Clinical experience in controlled studies with combinations including didanosine or zalcitabine is limited.

Two controlled studies (006 and ACTG 364) of approximately one year duration with efavirenz in combination with NRTIs and/or PIs, have demonstrated reduction of viral load below the limit of quantification of the assay and increased CD4 lymphocytes in antiretroviral therapy-naïve and NRTI-experienced HIV-infected patients. Study 020 showed similar activity in NRTI-experienced patients over 24 weeks. In these studies the dose of efavirenz was 600 mg once daily; the dose of indinavir was 1,000 mg every 8 hours when used with efavirenz and 800 mg every 8 hours when used without efavirenz. The dose of nelfinavir was 750 mg given three times a day. The standard doses of NRTIs given every 12 hours were used in each of these studies.

Study 006, a randomised, open-label trial, compared efavirenz + zidovudine + lamivudine or efavirenz + indinavir with indinavir + zidovudine + lamivudine in 1,266 patients who were required to be efavirenz-, lamivudine-, NNRTI-, and PI-naïve at study entry. The mean baseline CD4 cell count was 341 cells/mm³ and the mean baseline HIV-RNA level was 60,250 copies/ml. Efficacy results for study 006 on a subset of 614 patients who had been enrolled for at least 48 weeks are found in Table 2. In the analysis of responder rates (the non-completer equals failure analysis [NC = F]), patients who terminated the study early for any reason, or who had a

missing HIV-RNA measurement that was either preceded or followed by a measurement above the limit of assay quantification were considered to have HIV-RNA above 50 or above 400 copies/ml at the missing time points.

Table 2: Efficacy results for study 006

| | | Responder rates (NC = F ^a) Plasma HIV-RNA | | Mean change from baseline-CD4 cell count cells/mm ³ (S.E.M. ^c) |
|--------------------------------|-----|--|--|--|
| | | < 400 copies/ml (95% C.I. ^b) | < 50 copies/ml (95% C.I. ^b) | |
| Treatment Regimen ^d | n | 48 weeks | 48 weeks | 48 weeks |
| EFV + ZDV + 3TC | 202 | 67% (60%, 73%) | 62% (55%, 69%) | 187 (11.8) |
| EFV + IDV | 206 | 54% (47%, 61%) | 48% (41%, 55%) | 177 (11.3) |
| IDV + ZDV + 3TC | 206 | 45% (38%, 52%) | 40% (34%, 47%) | 153 (12.3) |

a NC = F, noncompleter = failure.

b C.I., confidence interval.

c S.E.M., standard error of the mean

d EFV, efavirenz; ZDV, zidovudine; 3TC, lamivudine; IDV, indinavir

Long-term results at 168 weeks of study 006 (160 patients completed study on treatment with EFV+IDV, 196 patients with EFV+ZDV+3TC and 127 patients with IDV+ZDV+3TC, respectively), suggest durability of response in terms of proportions of patients with HIV RNA < 400 copies/ml, HIV RNA < 50 copies/ml and in terms of mean change from baseline CD4 cell count.

Efficacy results for studies ACTG 364 and 020 are found in Table 3. Study ACTG 364 enrolled 196 patients who had been treated with NRTIs but not with PIs or NNRTIs. Study 020 enrolled 327 patients who had been treated with NRTIs but not with PIs or NNRTIs. Physicians were allowed to change their patient's NRTI regimen upon entry into the study. Responder rates were highest in patients who switched NRTIs.

Table 3: Efficacy results for studies ACTG 364 and 020

| Study Number/ Treatment Regimens ^b | n | Responder rates (NC = F ^a) Plasma HIV-RNA | | | | Mean change from baseline-CD4 cell count | |
|--|---|--|--------------------------|---|------------|--|------------------------|
| | | % | (95% C.I. ^c) | % | (95% C.I.) | cells/mm ³ | (S.E.M. ^d) |

| | | | | | | | |
|----------------------------|-----|--------------------|----------|-------------------|----------|-----|--------|
| Study ACTG 364 48 weeks | | < 500 copies/ml | | < 50 copies/ml | | | |
| EFV + NFV + NRTIs | 65 | 70 | (59,82) | --- | --- | 107 | (17.9) |
| EFV + NRTIs | 65 | 58 | (46,70) | --- | --- | 114 | (21.0) |
| NFV + NRTIs | 66 | 30 | (19,42) | --- | --- | 94 | (13.6) |
| | | | | | | | |
| Study 020 24 weeks | | < 400 copies/ml | | < 50 copies/ml | | | |
| EFV + IDV + NRTIs | 157 | 60 | (52, 68) | 49 | (41, 58) | 104 | (9.1) |
| IDV + NRTIs | 170 | 51 | (43, 59) | 38 | (30, 45) | 77 | (9.9) |
| | | | | | | | |

- a NC = F, noncompleter = failure.
- b EFV, efavirenz; ZDV, zidovudine; 3TC, lamivudine; IDV, indinavir; NRTI, nucleoside reverse transcriptase inhibitor; NFV, nelfinavir.
- c C.I., confidence interval for proportion of patients in response.
- d S.E.M., standard error of the mean
- not performed.

Paediatric population

Study AI266922 was an open-label study to evaluate the pharmacokinetics, safety, tolerability, and antiviral activity of efavirenz in combination with didanosine and emtricitabine in antiretroviral-naïve and -experienced paediatric patients.

Thirty-seven patients 3 months to 6 years of age (median 0.7 years) were treated with efavirenz. At baseline, median plasma HIV-1 RNA was 5.88 log₁₀ copies/mL, median CD4+ cell count was 1144 cells/mm³, and median CD4+ percentage was 25%. The median time on study therapy was 132 weeks; 27% of patients discontinued before Week 48. Using an ITT analysis, the overall proportions of patients with HIV RNA <400 copies/mL and <50 copies/mL at Week 48 were 57% (21/37) and 46% (17/37), respectively. The median increase from baseline in CD4+ count at 48 weeks was 215 cells/mm³ and the median increase in CD4+ percentage was 6%.

Study PACTG 1021 was an open-label study to evaluate the pharmacokinetics, safety, tolerability, and antiviral activity of efavirenz in combination with didanosine and emtricitabine in paediatric patients who were antiretroviral therapy naïve. Forty-three patients 3 months to 21 years of age (median 9.6 years) were dosed with efavirenz. At baseline, median plasma HIV-1 RNA was 4.8 log₁₀ copies/mL, median CD4+ cell count was 367 cells/mm³, and median CD4+ percentage was 18%. The median time on study therapy was 181 weeks; 16% of patients discontinued before Week 48. Using an ITT analysis, the overall proportions of patients with HIV RNA <400 copies/mL and <50 copies/mL at Week 48 were 77% (33/43) and 70% (30/43), respectively. The median increase from baseline in CD4+ count at 48 weeks of therapy was 238 cells/mm³ and the median increase in CD4+ percentage was 13%.

Study PACTG 382 was an open-label study to evaluate the pharmacokinetics, safety, tolerability, and antiviral activity of efavirenz in combination with nelfinavir and an NRTI in antiretroviral-naïve and NRTI-experienced paediatric patients. One hundred two patients 3 months to 16 years of age (median 5.7 years) were treated with efavirenz. Eighty-seven percent of patients had received prior antiretroviral therapy. At baseline, median plasma HIV-1 RNA was $4.57 \log_{10}$ copies/mL, median CD4+ cell count was 755 cells/mm³, and median CD4+ percentage was 30%. The median time on study therapy was 118 weeks; 25% of patients discontinued before Week 48. Using an ITT analysis, the overall proportion of patients with HIV RNA <400 copies/mL and <50 copies/mL at Week 48 were 57% (58/102) and 43% (44/102), respectively. The median increase from baseline in CD4+ count at 48 weeks of therapy was 128 cells/mm³ and the median increase in CD4+ percentage was 5%.

5.2 Pharmacokinetic properties

Absorption

Peak efavirenz plasma concentrations of 1.6 - 9.1 µM were attained by 5 hours following single oral doses of 100 mg to 1,600 mg administered to uninfected volunteers. Dose related increases in C_{max} and AUC were seen for doses up to 1,600 mg; the increases were less than proportional suggesting diminished absorption at higher doses. Time to peak plasma concentrations (3 - 5 hours) did not change following multiple dosing and steady-state plasma concentrations were reached in 6 - 7 days.

In HIV infected patients at steady state, mean C_{max} , mean C_{min} , and mean AUC were linear with 200 mg, 400 mg, and 600 mg daily doses. In 35 patients receiving efavirenz 600 mg once daily, steady state C_{max} was 12.9 ± 3.7 µM (29%) [mean \pm S.D. (% C.V.)], steady state C_{min} was 5.6 ± 3.2 µM (57%), and AUC was 184 ± 73 µM·h (40%).

Effect of food

The AUC and C_{max} of a single 600 mg dose of efavirenz film-coated tablets in uninfected volunteers was increased by 28% (90% CI : 22 – 33%) and 79% (90% CI : 58 – 102%), respectively, when given with a high-fat meal, relative to when given under fasted conditions (see section 4.4).

Distribution

Efavirenz is highly bound (approximately 99.5 - 99.75%) to human plasma proteins, predominantly albumin. In HIV-1 infected patients (n = 9) who received efavirenz 200 to 600 mg once daily for at least one month, cerebrospinal fluid concentrations ranged from 0.26 to 1.19% (mean 0.69%) of the corresponding plasma concentration. This proportion is approximately 3-fold higher than the non-protein-bound (free) fraction of efavirenz in plasma.

Biotransformation

Studies in humans and *in vitro* studies using human liver microsomes have demonstrated that efavirenz is principally metabolised by the cytochrome P450 system to hydroxylated metabolites with subsequent glucuronidation of these hydroxylated metabolites. These metabolites are essentially inactive against HIV-1. The *in vitro* studies suggest that CYP3A4 and CYP2B6 are the major isozymes responsible for efavirenz metabolism and that it inhibited P450 isozymes 2C9, 2C19, and 3A4. In *in vitro* studies efavirenz did not inhibit CYP2E1 and inhibited CYP2D6 and CYP1A2 only at concentrations well above those achieved clinically. Efavirenz plasma exposure may be increased in patients with the homozygous G516T genetic variant of the CYP2B6 isoenzyme. The clinical implications of such an association are unknown; however, the potential for an increased frequency and severity of efavirenz-associated adverse events cannot be excluded. Efavirenz has been shown to induce CYP3A4 and CYP2B6, resulting in the induction of its own metabolism, which may be clinically relevant in some patients. In uninfected volunteers, multiple doses of 200 - 400 mg per day for 10 days resulted in a lower than predicted extent of accumulation (22 - 42% lower) and a shorter terminal half-life compared with single dose administration (see below). Efavirenz has also been shown to induce UGT1A1. Exposures of raltegravir (a UGT1A1 substrate) are reduced in the presence of efavirenz (see section 4.5, table 1). Although *in vitro* data suggest that efavirenz inhibits CYP2C9 and CYP2C19, there have been contradictory reports of both increased and decreased exposures to substrates of these enzymes when coadministered with efavirenz *in vivo*. The net effect of coadministration is not clear.

Elimination

Efavirenz has a relatively long terminal half-life of at least 52 hours after single doses and 40 - 55 hours after multiple doses. Approximately 14 - 34% of a radiolabelled dose of efavirenz was recovered in the urine and less than 1% of the dose was excreted in urine as unchanged efavirenz.

Hepatic impairment

In a single-dose study, half life was doubled in the single patient with severe hepatic impairment (Child Pugh Class C), indicating a potential for a much greater degree of accumulation. A multiple-dose study showed no significant effect on efavirenz pharmacokinetics in patients with mild hepatic impairment (Child-Pugh Class A) compared with controls. There were insufficient data to determine whether moderate or severe hepatic impairment (Child-Pugh Class B or C) affects efavirenz pharmacokinetics.

Gender, race, elderly

Although limited data suggest that females as well as Asian and Pacific Island patients may have higher exposure to efavirenz, they do not appear to be less tolerant of efavirenz. Pharmacokinetic studies have not been performed in the elderly.

Paediatric population

The pharmacokinetic parameters for efavirenz at steady state in paediatric patients were predicted by a population pharmacokinetic model and are summarised in Table 4 by weight ranges that correspond to the recommended doses.

Table 4: Predicted steady-state pharmacokinetics of efavirenz (capsules/capsule sprinkles) in HIV-infected paediatric patients

| Body Weight | Dose | Mean AUC ₍₀₋₂₄₎ μM·h | Mean C _{max} μg/mL | Mean C _{min} μg/mL |
|-------------|--------|------------------------------------|--------------------------------|--------------------------------|
| 3.5-5 kg | 100 mg | 220.52 | 5.81 | 2.43 |
| 5-7.5 kg | 150 mg | 262.62 | 7.07 | 2.71 |
| 7.5-10 kg | 200 mg | 284.28 | 7.75 | 2.87 |
| 10-15 kg | 200 mg | 238.14 | 6.54 | 2.32 |
| 15-20 kg | 250 mg | 233.98 | 6.47 | 2.3 |
| 20-25 kg | 300 mg | 257.56 | 7.04 | 2.55 |
| 25-32.5 kg | 350 mg | 262.37 | 7.12 | 2.68 |
| 32.5-40 kg | 400 mg | 259.79 | 6.96 | 2.69 |
| >40 kg | 600 mg | 254.78 | 6.57 | 2.82 |

5.3 Preclinical safety data

Efavirenz was not mutagenic or clastogenic in conventional genotoxicity assays. Efavirenz induced foetal resorptions in rats. Malformations were observed in 3 of 20 fetuses/newborns from efavirenz-treated cynomolgus monkeys given doses resulting in plasma efavirenz concentrations similar to those seen in humans. Anencephaly and unilateral anophthalmia with secondary enlargement of the tongue were observed in one foetus, microphthalmia was observed in another foetus, and cleft palate was observed in a third foetus. No malformations were observed in fetuses from efavirenz-treated rats and rabbits.

Biliary hyperplasia was observed in cynomolgus monkeys given efavirenz for ≥ 1 year at a dose resulting in mean AUC values approximately 2-fold greater than those in humans given the recommended dose. The biliary hyperplasia regressed upon cessation of dosing. Biliary fibrosis has been observed in rats. Non-sustained convulsions were observed in some monkeys receiving efavirenz for ≥ 1 year, at doses yielding plasma AUC values 4 to 13-fold greater than those in humans given the recommended dose (see sections 4.4 and 4.8).

Carcinogenicity studies showed an increased incidence of hepatic and pulmonary tumours in female mice, but not in male mice. The mechanism of tumour formation and the potential relevance for humans are not known.

Carcinogenicity studies in male mice, male and female rats were negative. While the carcinogenic potential in humans is unknown, these data suggest that the clinical benefit of efavirenz outweighs the potential carcinogenic risk to humans.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Efavirenz 600 mg film-coated tablets contain:

Core

Cellulose, microcrystalline
Croscarmellose sodium
Hydroxypropyl cellulose
Lactose monohydrate
Silica, colloidal anhydrous
Magnesium stearate

Film-coating

Hypromellose
Macrogol 400
Titanium dioxide (E171)
Iron oxide yellow (E172)
Iron oxide red (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

The in-use shelf life of the product when stored in HDPE bottles is 100 days.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

PVC / PVDC-Aluminium blister pack of 30, 60, 90 and 100 film-coated tablets.
PVC / PVDC-Aluminium perforated unit dose blister pack of 30 x 1 film-coated tablets.

HDPE bottles with PP screw cap of 30 and 500 film-coated tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused product or waste should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Mc Dermott Laboratories Ltd.
T/A Gerard Laboratories
35/36 Baldoyle Industrial Estate,
Grange Road,
Dublin 13

8 MARKETING AUTHORISATION NUMBER

PA0577/156/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 12th October 2012

Date of last renewal: 13th August 2017

10 DATE OF REVISION OF THE TEXT

November 2018