

Package leaflet: Information for the patient

Agomelatine Viatris 25 mg film-coated tablets

agomelatine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Agomelatine Viatris is and what it is used for
2. What you need to know before you take Agomelatine Viatris
3. How to take Agomelatine Viatris
4. Possible side effects
5. How to store Agomelatine Viatris
6. Contents of the pack and other information

1. What Agomelatine Viatris is and what it is used for

Agomelatine Viatris contains the active ingredient agomelatine. It belongs to a group of medicines called antidepressants and you have been given this medicine to treat your depression.

Agomelatine Viatris is used in adults.

Depression is a continuing disturbance of mood that interferes with everyday life. The symptoms of depression vary from one person to another, but often include deep sadness, feelings of worthlessness, loss of interest in favourite activities, sleep disturbances, feeling of being slowed down, feelings of anxiety, changes in weight.

The expected benefits of Agomelatine Viatris are to reduce and gradually remove the symptoms related to your depression.

2. What you need to know before you take Agomelatine Viatris

Do not take Agomelatine Viatris

- if you are allergic to agomelatine or any of the other ingredients of this medicine (listed in section 6).
- **if your liver does not work properly (hepatic impairment).**
- if you are taking fluvoxamine (another medicine used in the treatment of depression) or ciprofloxacin (an antibiotic).

Warnings and precautions

There could be some reasons why Agomelatine Viatris may not be suitable for you:

- If you are taking medicines known to affect the liver. Ask your doctor for advice on which medicine that is.
- If you are obese or overweight, ask your doctor for advice.
- If you are diabetic, ask your doctor for advice.
- If you have increased levels of liver enzymes before treatment, your doctor will decide if Agomelatine Viatris is right for you.
- If you have bipolar disorder, have experienced or if you develop manic symptoms (a period of abnormally high excitability and emotions) talk to your doctor before you start taking this medicine or before you continue with this medicine (see also under “Possible side effects” in section 4).
- If you are suffering from dementia, your doctor will make an individual evaluation of whether it is right for you to take Agomelatine Viatris.

During your treatment with Agomelatine Viatris:

What to do to avoid potential serious liver problems

- Your doctor should have checked that your liver is working properly **before starting the treatment**. Some patients may get increased levels of liver enzymes in their blood during treatment with agomelatine. Therefore, follow-up tests should take place at the following time points:

	before initiation or	around 3 weeks	around 6 weeks	around 12 weeks	around 24 weeks
Blood tests	ü	ü	ü	ü	ü

Based on the evaluation of these tests your doctor will decide whether you should receive or continue using Agomelatine Viatriis (see also under “How to take Agomelatine Viatriis” in section 3).

Be vigilant about signs and symptoms that your liver may not be working properly

- **If you observe** any of these signs and symptoms of liver problems: **unusual darkening of the urine, light coloured stools, yellow skin/eyes, pain in the upper right belly, unusual fatigue (especially associated with other symptoms listed above), seek urgent advice from a doctor who may advise you to stop taking Agomelatine Viatriis.**

Effect of agomelatine is not documented in patients aged 75 years and older. Agomelatine Viatriis should therefore not be used in these patients.

Thoughts of suicide and worsening of your depression

If you are depressed, you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- if you have previously had thoughts about killing or harming yourself.
- if you are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in young adults (aged less than 25 years) with psychiatric conditions who were being treated with an antidepressant.

If you have thoughts of harming or killing yourself at any time, contact your doctor or go to a hospital straight away.

You may find it helpful to tell a relative or close friend that you are depressed and ask them to read this leaflet. You might ask them to tell you if they think your depression is getting worse, or if they are worried about changes in your behaviour.

Children and adolescents

Agomelatine Viatris is not recommended in children below 7 years due to lack of information. No data is available. Agomelatine Viatris should not be used in children and adolescents aged 7 to 17 years because safety and efficacy have not been established.

Other medicines and Agomelatine Viatris

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

You should not take Agomelatine Viatris together with certain medicines (see also under “Do not take Agomelatine Viatris” in section 2): fluvoxamine (another medicine used in the treatment of depression), ciprofloxacin (an antibiotic) can modify the expected dose of agomelatine in your blood.

Make sure to tell your doctor if you are taking any of the following medicines: propranolol (a beta-blocker used in the treatment of hypertension), enoxacin (antibiotic) and if you are smoking more than 15 cigarettes/day.

Agomelatine Viatris with alcohol

It is not advisable to drink alcohol while you are being treated with agomelatine.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Breastfeeding should be discontinued if you take Agomelatine Viatris.

Driving and using machines

You might experience dizziness or sleepiness which could affect your ability to drive or operate machinery. Make sure that your reactions are normal before driving or operating machines.

Agomelatine Viatris contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Agomelatine Viatris

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose of Agomelatine Viatris is one tablet (25 mg) at bedtime. In some cases, your doctor may prescribe a higher dose (50 mg), i.e. two tablets to be taken together at bedtime.

Agomelatine Viatris starts to act on symptoms of depression in most depressed people within two weeks of starting treatment.

Your depression should be treated for a sufficient period of at least 6 months to ensure that you are free of symptoms.

Your doctor may continue to give you Agomelatine Viatris when you are feeling better to prevent your depression from returning.

Agomelatine Viatris is for oral use. You should swallow your tablet with a drink of water. This medicine can be taken with or without food.

How to switch from an antidepressant medicine (SSRI/SNRI) to agomelatine?

If your doctor changes your previous antidepressant medicine from an SSRI or SNRI to agomelatine, he/she will advise you on how you should discontinue your previous medicine when starting Agomelatine Viatris.

You may experience discontinuation symptoms related to stopping of your previous medicine for a few weeks, even if the dose of your previous antidepressant medicine is decreased gradually.

Discontinuation symptoms include: dizziness, numbness, sleep disturbances, agitation or anxiety, headaches, feeling sick, being sick and shaking. These effects are usually mild to moderate and disappear spontaneously within a few days.

If Agomelatine Viatris is started while reducing the dosage of the previous medicine, possible discontinuation symptoms should not be confounded with a lack of early effect of Agomelatine Viatris.

You should discuss with your doctor on the best way of stopping your previous antidepressant medicine when starting Agomelatine Viatris.

Surveillance of the liver function (see also section 2):

Your doctor will run laboratory tests to check that your liver is working properly before starting treatment and then periodically during treatment, usually after 3 weeks, 6 weeks, 12 weeks and 24 weeks.

If your doctor increases the dose to 50 mg, laboratory tests should be performed at this initiation and then periodically during treatment, usually after 3 weeks, 6 weeks, 12 weeks and 24 weeks. Thereafter tests will be taken if the doctor finds it necessary.

You must not use Agomelatine Viatris if your liver does not work properly.

If you have trouble with your kidneys, your doctor will make an individual evaluation of whether it is safe for you to take Agomelatine Viatris.

If you take more Agomelatine Viatris than you should

If you have taken more Agomelatine Viatris than you should, or if for example a child has taken medicine by accident, contact your doctor immediately.

The experience of overdoses with agomelatine is limited but reported symptoms include pain in the upper part of the stomach, somnolence, fatigue, agitation, anxiety, tension, dizziness, cyanosis or malaise.

If you forget to take Agomelatine Viatris

Do not take a double dose to make up for a forgotten dose. Just carry on with the next dose at the usual time.

If you stop taking Agomelatine Viatris

Do not stop taking your medicine without the advice of your doctor even if you feel better.

If you think that the effect of Agomelatine Viatris is too strong or too weak, talk to your doctor or pharmacist. If you have any further questions on the use of this product, please ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Most side effects are mild or moderate. They usually occur within the first two weeks of the treatment and are usually temporary.

These side effects include:

- Very common (may affect more than to 1 in 10 people) :

headache

- Common (may affect up to 1 in 10 people):

dizziness, sleepiness (somnolence), difficulty in sleeping (insomnia), feeling sick (nausea), diarrhoea, constipation, abdominal pain,

back pain, tiredness, anxiety, abnormal dreams, increased levels of liver enzymes in your blood, vomiting, weight increased.

- Uncommon (may affect up to 1 in 100 people):

migraine, pins and needles in the fingers and toes (paraesthesia), blurred vision, restless legs syndrome (a disorder that is characterized by an

uncontrollable urge to move the legs), ringing in the ears, excessive sweating (hyperhidrosis), eczema, pruritus, urticaria (hives), agitation, irritability, restlessness, aggressive behaviour, nightmares, mania/hypomania (see also under “Warnings and precautions” in section 2), suicidal

thoughts or behaviour, confusion, weight decreased, muscle pain.

- Rare (may affect up to 1 in 1,000 people):

serious skin eruption (erythematous rash),

face oedema (swelling) and angioedema (swelling of the face, lips, tongue and/or throat that may

cause difficulty in breathing or swallowing), hepatitis, yellow coloration of the skin or the whites of

the eyes (jaundice), hepatic failure*, hallucinations, inability to remain still (due to physical and mental unrest), inability to completely empty the bladder.

* Few cases resulting in liver transplantation or death have been reported.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA



Pharmacovigilance Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Agomelatine Viatris

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month.

Store in the original package in order to protect from moisture. This medicine does not require any special temperature storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Agomelatine Viatris contains

- The active substance is agomelatine. Each film-coated tablet contains agomelatine-citric acid co-crystal equivalent to 25 mg of agomelatine. The other ingredients of the core are:

colloidal silicified dioxide, microcrystalline cellulose, mannitol, povidone, colloidal anhydrous silica, crospovidone, sodium stearyl fumarate, magnesium stearate, stearic acid

- The other ingredients of the coating are:

hypromellose, macrogol, titanium dioxide (E171), talc, yellow iron oxide (E172)

What Agomelatine Viatris looks like and contents of the pack

Agomelatine Viatris 25 mg film-coated tablets are yellow, oblong, biconvex (surface curved outwards on both sides) film-coated tablets.

Agomelatine Viatris 25 mg film-coated tablets are available in blister packs containing 7, 14, 28, 42, 56, 84, 98 or 100 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Viartis Limited, Damastown Industrial Park, Mulhuddart, Dublin 15, DUBLIN, Ireland.

Manufacturer

Medis International a.s., výrobní závod Bolatice, Prumyslova 961/16,
747 23 Bolatice, Czech Republic.

This medicinal product is authorised in the Member States of the EEA under the following names:

Czech Republic	Agomelatine Viatris
Germany	Agomelatin Mylan 25 mg Filmtabletten
Denmark	Agomelatin Viatris
Estonia	Agomelatine Viatris
Finland	Agomelatine Viatris
France	Agomelatine Viatris 25 mg, comprimé pelliculé
Hungary	Agomelatin Mylan 25 mg filmtablettá
Ireland	Agomelatine Viatris 25 mg film-coated tablets
Latvia	Agomelatine Viatris 25 mg apvalkotās tabletes
Lithuania	Agomelatine Viatris 25 mg plėvele dengtos tabletės
Netherlands	Agomelatine Viatris 25 mg, filmomhulde tabletten
Portugal	Agomelatina Mylan
Sweden	Agomelatine Viatris 25 mg filmdragerade tabletter
Slovakia	Agomelatine Viatris 25 mg

This leaflet was last revised 04/2025.