

Package leaflet: Information for the patient

Fental Matrix 25 micrograms/hour Transdermal patch
Fental Matrix 50 micrograms/hour Transdermal patch
Fental Matrix 75 micrograms/hour Transdermal patch
Fental Matrix 100 micrograms/hour Transdermal patch

fentanyl

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Fental Matrix is and what it is used for
2. What you need to know before you use Fental Matrix
3. How to use Fental Matrix
4. Possible side effects
5. How to store Fental Matrix
6. Contents of the pack and other information

1. What Fental Matrix is and what it is used for

The name of your medicine is Fental Matrix.

The patches help relieve pain that is very bad and long-lasting:

- in adults who need continuous pain treatment
- in children above 2 years of age who are already using opioid medicines and who need continuous pain treatment.

Fental Matrix contains the active substance called fentanyl. It belongs to a group of strong painkillers called opioids.

2. What you need to know before you use Fental Matrix

Do not use Fental Matrix if:

- You are allergic to fentanyl, colophonium resin (hydrogenated), soya, peanuts or any of the other ingredients of this medicine (listed in section 6)
- You have pain which lasts only for a short period, such as sudden pain or pain after having an operation
- You have breathing difficulties, with slow or shallow breathing.

Do not use this medicine if any of the above apply to you or your child. If you are not sure, talk to your doctor or pharmacist before using Fental Matrix.

Warnings and precautions

- Fental Matrix can have life-threatening side effects in people who are not already regularly using prescribed opioid medicines.
- Fental Matrix is a medicine that could be life-threatening to children, even if the patches have been used. Bear in mind that a sticky patch (unused or used) could be tempting to a child and if it sticks to a child's skin or they put it in their mouth, the result may be fatal.
- Store this medicine in a safe and secure place, where other people cannot access it - see section 5 for more information.

Patch sticking to another person

The patch should be used only on the skin of the person for whom it has been prescribed. There have been reports of patches accidentally sticking to a family member while in close physical contact or sharing the same bed as the person wearing the patch. A patch accidentally sticking to another person (particularly a child) can cause the medicine in the patch to go through the skin of the other person and cause serious side effects such as breathing difficulties, with slow or shallow breathing which may be fatal. In case the patch sticks to the skin of another person, take the patch off right away and get medical attention.

Take special care with Fental Matrix

Talk to your doctor or pharmacist before using this medicine if any of the following apply to you - your doctor may need to check you more closely if:

- You have ever had problems with your lungs or breathing
- You have ever had problems with your heart, liver, kidneys, or low blood pressure
- You have ever had a brain tumour
- You have ever had persistent headaches or a head injury
- You are elderly - you may be more sensitive to the effects of this medicine
- You have a condition called ‘myasthenia gravis’ in which muscles become weak and tire easily.

If any of the above apply to you (or you are not sure), talk to your doctor or pharmacist before using Fental Matrix.

While using the patch, tell your doctor if you have breathing problems while sleeping. Opioids like Fental Matrix can cause sleep-related breathing disorders such as sleep apnoea (breathing pauses during sleep) and sleep-related hypoxaemia (low oxygen level in the blood). Tell your doctor if you, your partner or carer notice you have any of the following:

- breathing pauses during sleep
- night awakening due to shortness of breath
- difficulties staying asleep
- excessive drowsiness during the day. Your doctor may decide to change your dose.

While using the patch, tell your doctor if you notice a change in the pain you are feeling. If you feel:

- your pain is no longer relieved by the patch
- an increase in pain
- there is a change in how you feel the pain (for example, you feel pain in another part of your body)
- pain when something touches your body that you wouldn’t expect to hurt you.

Do not change the dose yourself. Your doctor may decide to change your dose or treatment.

Side effects and Fental Matrix

- Fental Matrix may make you unusually drowsy, and make your breathing more slow or shallow. Very rarely these breathing problems can be life-threatening or even fatal, especially in people who have not used strong opioid painkillers (like Fental Matrix or morphine) before. If you, or your partner or carer, notice that the person wearing the patch is unusually drowsy, with slow or shallow breathing:
 - Take the patch off
 - Call a doctor, or go to your nearest hospital straight away
 - Keep the person moving and talking as much as possible
- If you get a fever while using Fental Matrix, tell your doctor - this may increase the amount of medicine that passes through your skin
- Fental Matrix may cause constipation, talk to your doctor or pharmacist for advice on how to prevent or relieve constipation.

See section 4 for a full list of possible side effects.

Fental Matrix, like other opioids, may affect the normal production of hormones in the body such as cortisol, prolactin, or sex hormones, particularly if you have taken Fental Matrix for long periods of

time. The effects of these hormonal changes may include feeling or being sick (including vomiting), loss of appetite, tiredness, weakness, dizziness, low blood pressure, infertility, or decreased sex drive. In addition, female patients may experience changes in menstrual cycle, while male patients may experience impotence or enlarged breasts. If you notice any of these signs, speak to your doctor.

When you are wearing the patch do not expose it to direct heat such as heating pads, electric blankets, hot-water bottles, heated water beds or heat or tanning lamps. Do not sunbathe, have long hot baths or saunas or use hot whirlpool spa baths. If you do, you may increase the amount of medicine you get from the patch.

Long-term use and tolerance

This medicine contains fentanyl which is an opioid medicine. Repeated use of opioid painkillers can result in the drug being less effective (you become accustomed to it, known as drug tolerance). You may also become more sensitive to pain while using Fental Matrix. This is known as hyperalgesia. Increasing the dose of your patches may help to further reduce your pain for a while, but it may also be harmful. If you notice that your medicine becomes less effective, talk to your doctor. Your doctor will decide whether it is better for you to increase the dose or to gradually decrease your use of Fental Matrix.

Dependence and addiction

This medicine contains fentanyl, which is an opioid. It can cause dependence and/or addiction.

Repeated use of Fental Matrix can also lead to dependence, abuse and addiction which may result in life-threatening overdose. The risk of these side effects can increase with a higher dose and longer duration of use. Dependence or addiction can make you feel that you are no longer in control of how much medicine you need to use or how often you need to use it. You might feel that you need to carry on using your medicine, even when it doesn't help to relieve your pain.

The risk of becoming dependent or addicted varies from person to person. You may have a greater risk of becoming dependent on or addicted to Fental Matrix if:

- You or anyone in your family have ever abused or been dependent on alcohol, prescription medicines or illegal drugs ("addiction").
- You are a smoker.
- You have ever had problems with your mood (depression, anxiety or a personality disorder) or have been treated by a psychiatrist for other mental illness.

If you notice any of the following signs whilst using Fental Matrix, it could be a sign that you have become dependent or addicted.

- You need to use the medicine for longer than advised by your doctor
- You need to use more than the recommended dose
- You are using the medicine for reasons other than prescribed, for instance, 'to stay calm' or 'help you sleep'
- You have made repeated, unsuccessful attempts to quit or control the use of the medicine
- When you stop taking the medicine you feel unwell, and you feel better once using the medicine again ('withdrawal effects').

If you notice any of these signs, speak to your doctor to discuss the best treatment pathway for you, including when it is appropriate to stop and how to stop safely.

Withdrawal symptoms when stopping Fental Matrix

Do not suddenly stop taking this medicine. Withdrawal symptoms such as restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, loss of appetite, shaking, shivering or sweating may occur. If you want to stop taking this medicine, talk to your doctor first. Your doctor will tell you how to do this, usually by reducing the dose gradually so that any unpleasant withdrawal effects are kept to a minimum.

Other medicines and Fental Matrix

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines that you buy without a prescription or herbal medicines. You should also tell your pharmacist that you are using Fental Matrix if you buy any medicines from your pharmacy.

Your doctor will know which medicines are safe to take with Fental Matrix. You may need to be closely monitored if you are taking some of the types of medicines listed below or if you stop taking some of the types of medicines listed below, as this may affect the strength of Fental Matrix you need.

In particular, tell your doctor or pharmacist if you are taking:

- Other medicines for pain, such as other opioid painkillers (such as buprenorphine, nalbuphine, or pentazocine) and some painkillers for nerve pain (gabapentin and pregabalin)
- Medicines for helping you sleep (such as temazepam, zaleplon, or zolpidem)
- Medicines to help you calm down (tranquillisers, such as alprazolam, clonazepam, diazepam, hydroxyzine, or lorazepam) and medicines for mental conditions (anti-psychotics, such as aripiprazole, haloperidol, olanzapine, risperidone, or phenothiazines)
- Medicines for relaxing your muscles (such as cyclobenzaprine or diazepam)
- Some medicines used to treat depression called SSRIs or SNRIs (such as citalopram, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline, or venlafaxine). - see below for more information.
- Some medicines used to treat depression or Parkinson's disease called MAOIs (such as isocarboxazid, phenelzine, selegiline or tranylcypromine). You should not take Fental Matrix within 14 days of stopping these medicines - see below for more information.
- Some antihistamines, especially ones that make you sleepy (such as chlorphenamine, clemastine, cyproheptadine, diphenhydramine or hydroxyzine)
- Some antibiotics used to treat infection (such as erythromycin or clarithromycin)
- Medicines used to treat fungal infection (such as itraconazole, ketoconazole, fluconazole, or voriconazole)
- Medicines used to treat HIV infection (such as ritonavir)
- Medicines used to treat an irregular heartbeat (such as amiodarone, diltiazem or verapamil)
- Medicines to treat tuberculosis (such as rifampicin)
- Some medicines used to treat epilepsy (such as carbamazepine, phenobarbital or phenytoin)
- Some medicines used to treat nausea or motion sickness (such as phenothiazines)
- Some medicines used to treat heartburn or ulcers (such as cimetidine)
- Some medicines used to treat angina (chest pain) or high blood pressure (such as nicardipine)
- Some medicines used to treat cancer of the blood (such as idelalisib).

Fental Matrix with antidepressants

The risk of side effects increases if you are taking medicines such as certain antidepressants. Fental Matrix may interact with these medicines and you may experience changes to mental status such as feeling agitated, seeing, feeling, hearing, or smelling things that are not there (hallucinations) and other effects such as changing blood pressure, fast heartbeat, high body temperature, overactive reflexes, lack of coordination, muscle stiffness, nausea, vomiting and diarrhoea.

Use with central nervous system depressants, including benzodiazepines, alcohol and some narcotic medicines

Tell your doctor if you are taking any other medicines that slow down your central nervous system (CNS depressants). CNS depressants are, for example, medicines that make you sleepy, reduce anxiety or decrease awareness (see also "Other medicines and Fental Matrix"); but also alcohol and some narcotic medicines are CNS depressants. Taking these kinds of medicines with Fental Matrix can cause severe drowsiness, decreased awareness, breathing difficulties with slow or shallow breathing, coma and death. Because of this, concomitant use should only be considered when other treatment options are not possible.

However, if your doctor does prescribe Fental Matrix together with sedative medicines, the dose and duration of concomitant treatment should be limited by your doctor. Please follow your doctor's dose

recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

Do not drink alcohol while using Fental Matrix unless you have talked to your doctor first.

Operations

If you think that you are going to receive anaesthesia tell your doctor or dentist that you are using Fental Matrix.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

Fental Matrix should not be used during pregnancy unless you have discussed this with your doctor.

Fental Matrix should not be used during childbirth as the medicine can affect the breathing of the newborn child.

Prolonged use of Fental Matrix during pregnancy can cause withdrawal symptoms (such as high-pitched cry, jitteriness, fits, poor feeding and diarrhoea) in your newborn baby that could be life-threatening if not recognised and treated. Talk to your doctor immediately if you think your baby may have withdrawal symptoms.

Do not use Fental Matrix if you are breast-feeding. You should not breast-feed for 3 days after removing your Fental Matrix patch. This is because the medicine may pass into breast milk.

Driving and using machines

Fental Matrix can affect your ability to drive and use machines or tools as it may make you sleepy or dizzy. If this happens, do not drive or use any tools or machines. Do not drive while using this medicine until you know how it affects you.

Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive while taking this medicine.

Fental Matrix contains soya oil

If you are allergic to peanut or soya, do not use this medicine.

3. How to use Fental Matrix

Always use this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Your doctor will decide which strength of Fental Matrix is most suitable for you, taking into account the severity of your pain, your general condition and type of pain treatment that you have received so far.

Before starting treatment and regularly during treatment, your doctor will also discuss with you what you may expect from using Fental Matrix, when and how long you need to take it, when to contact your doctor, and when you need to stop it (see also section 2 “Withdrawal symptoms when stopping Fental Matrix”).

Using and changing the patches

- There is enough medicine in each patch to last **3 days (72 hours)**
- You should change your patch every third day, unless your doctor has told you differently
- Always remove the old patch **before** applying a new one
- Always change your patch at the **same time** of day every 3 Days (72 hours)
- If you are using more than one patch, change all your patches at the same time

- Make a note of the day, date and time you apply a patch, to remind you when you need to change your patch

The following table shows you when to change your patch:

Apply your patch on	⇔	Change your patch on
Monday	⇔	Thursday
Tuesday	⇔	Friday
Wednesday	⇔	Saturday
Thursday	⇔	Sunday
Friday	⇔	Monday
Saturday	⇔	Tuesday
Sunday	⇔	Wednesday

Where to apply the patch

Adults

- Apply the patch on a flat part of your upper body or arm (not over a joint).

Children

- Always apply the patch to the upper back to make it difficult for your child to reach it or take it off
- Every so often check that the patch remains stuck to the skin
- It is important that your child does not remove the patch and put it in their mouth as this could be life-threatening or even fatal
- Watch your child very closely for 48 hours after:
 - The first patch has been put on
 - A higher dose patch has been put on
- It may take some time for the patch to have its maximum effect. Therefore, your child might need to use other painkillers as well until the patches become effective. Your doctor will talk to you about this.

Adults and Children:

Do not apply the patch on

- The same place twice in a row
- Areas that you move a lot (joints), skin that is irritated or with cuts
- Skin that is very hairy. If there is hair, do not shave it (shaving irritates the skin). Instead, clip the hair as close to the skin as possible.

Putting a patch on

Step 1: Preparing the skin

- Make sure your skin is completely dry, clean and cool before you put the patch on
- If you need to clean the skin, just use cold water
- Do not use soap or any other cleansers, creams, moisturisers, oils or talc before applying the patch
- Do not stick a patch on straight after a hot bath or shower.

Step 2: Open the sachet

- Each patch is sealed in its own sachet
- Tear open the sachet at the notch
- Gently tear off the edge of the sachet completely
- Grasp both sides of the opened sachet and pull apart
- Take the patch out and use straight away
- Keep the empty sachet to dispose of the used patch later
- Use each patch once only
- Do not take the patch out of its sachet until you are ready to use it
- The patches may occasionally show crystallization (white structures at the edges and/or in the matrix surface). The presence of crystals will not affect how the patch works. Do not touch the sticky side of the patch or the crystals.

- Inspect the patch for any damage
- Do not use the patch if it has been divided, cut or looks damaged
- Never divide or cut the patch.

Step 3: Peel and press

- Make sure that the patch will be covered by loose clothing and not stuck under a tight or elasticated band
- Carefully peel the first part of the shiny plastic backing away from the centre of the patch. Try not to touch the sticky side of the patch.
- Press this sticky part of the patch onto the skin
- Remove the other part of the backing and press the whole patch onto the skin with the palm of your hand
- Hold for at least 30 seconds. Make sure it sticks well, especially the edges.

Step 4: Disposing of the patch

- As soon as you take a patch off, fold it firmly in half so that the sticky side sticks to itself
- Put it back in its original sachet and dispose of the sachet as instructed by your pharmacist
- Keep used patches out of sight and reach of children - even used patches contain some medicine which may harm children and may even be fatal.

Step 5: Wash

- Always wash your hands after you have handled the patch using clean water only.

More about using Fental Matrix

Everyday activities while using the patches

- The patches are waterproof
- You can shower or bathe while wearing a patch, but do not scrub the patch itself
- If your doctor agrees, you can exercise or play sport while wearing the patch
- You can also swim while wearing the patch, but:
 - Do not use hot whirlpool spa baths
 - Do not put a tight or elasticated band over the patch
- While you are wearing the patch do not expose it to direct heat such as heating pads, electric blankets, hot-water bottles, heated water beds, heat or tanning lamps. Do not sunbathe, have long hot baths or saunas. If you do, you may increase the amount of medicine you get from the patch.

How quickly will the patches work?

- It may take some time for your first patch to have its maximum effect
- Your doctor may give you other painkillers as well for the first day or so
- After this, the patch should help to relieve pain continuously so that you can stop taking other painkillers. However, your doctor may still prescribe extra painkillers from time to time.

How long will you use the patches for?

- Fental Matrix patches are for long-term pain. Your doctor will be able to tell you how long you can expect to use the patches.

If your pain gets worse

- If your pain suddenly gets worse after placing of your last patch you should check your patch. If it is no longer sticking well or has fallen off you should replace the patch (See also section “If a patch falls off”)
- If your pain gets worse over time while you are using these patches, your doctor may try a higher strength patch, or give you additional painkillers (or both)
- If increasing the strength of the patch does not help, your doctor may decide to stop the use of the patches.

If you use too many patches or the wrong strength patch

If you have stuck on too many patches or the wrong strength patch, take the patches off and contact a

doctor straight away.

Signs of overdose include trouble breathing or shallow breathing, tiredness, extreme sleepiness, being unable to think clearly, walk or talk normally and feeling faint, dizzy or confused. An overdose may also result in a brain disorder known as toxic leukoencephalopathy.

If you forget to change your patch

- If you forget, change your patch as soon as you remember and make note of the day and time
- Change the patch again after **3 days (72 hours)** as usual
- If you are very late changing your patch, you should talk to your doctor because you might need some extra painkillers, but do **not** apply an extra patch.

If a patch falls off

- If a patch falls off before it needs changing, stick a new one on straight away and make note of the day and time. Use a new area of skin on:
 - Your upper body or arm
 - Your child's upper back
- Let your doctor know this has happened and leave the patch on for another **3 days (72 hours)** or as directed by your doctor, before changing the new patch as usual
- If your patches keep falling off, talk to your doctor, pharmacist or nurse.

If you want to stop using the patches

- Do not suddenly stop taking this medicine. If you want to stop taking this medicine, talk to your doctor first. Your doctor will tell you how to do this, usually by reducing the dose gradually so that any unpleasant withdrawal effects are kept to a minimum. See also section 2 "Withdrawal symptoms when stopping Fental Matrix".
- If you have been using them for some time your body may have got used to them. Stopping suddenly may make you feel unwell
- If you stop using the patches, don't start again without asking your doctor first. You might need a different patch strength when you restart.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you or your partner, or carer, notice any of the following about the person wearing the patch, take the patch off and call a doctor, or go to your nearest hospital, straight away. You may need urgent medical treatment.

- Feeling unusually drowsy, breathing that is more slow or shallow than expected.
Follow the advice above and keep the person who was wearing the patch moving and talking as much as possible. Very rarely these breathing difficulties can be life-threatening or even fatal, especially in people who have not used strong opioid painkillers (like Fental Matrix or morphine) before.
(Uncommon, this may affect up to 1 in 100 people)
- Sudden swelling of the face or throat, severe irritation, reddening or blistering of your skin.
These may be signs of a severe allergic reaction (frequency cannot be estimated from the available data)
- Fits (seizures) (Uncommon, this may affect up to 1 in 100 people)
- Reduced consciousness or loss of consciousness (Uncommon, these may affect up to 1 in 100 people).

The following side effects have also been reported

Very common (may affect more than 1 in 10 people)

- Nausea, vomiting, constipation
- Feeling sleepy (somnolence)
- Feeling dizzy

- Headache.

Common (may affect up to 1 in 10 people)

- Allergic reaction
- Loss of appetite
- Difficulty sleeping
- Depression
- Feeling anxious or confused
- Seeing, feeling, hearing or smelling things that are not there (hallucinations)
- Muscle tremors or spasms
- Unusual feeling in the skin, such as tingling or crawling feelings (paraesthesia)
- Spinning sensation (vertigo)
- Heartbeat feels fast or uneven (palpitations, tachycardia)
- High blood pressure
- Being short of breath (dyspnoea)
- Diarrhoea
- Dry mouth
- Stomach pain or indigestion
- Excessive sweating
- Itching, skin rash or redness of the skin
- Being unable to pass urine or empty bladder completely
- Feeling very tired, weak or generally unwell
- Feeling cold
- Swollen hands, ankles or feet (peripheral oedema).

Uncommon (may affect up to 1 in 100 people)

- Feeling agitated or disoriented
- Feeling extremely happy (euphoria)
- Decreased feeling or sensitivity, especially in the skin (hypoesthesia)
- Loss of memory
- Blurred vision
- Slow heartbeat (bradycardia) or low blood pressure
- Blue colour to the skin caused by low oxygen in the blood (cyanosis)
- Loss of contractions of the gut (ileus)
- Itchy skin rash (eczema), allergic reaction or other skin disorders where the patch is placed
- Flu-like illness
- Feeling of body temperature change
- Fever
- Muscle twitching
- Difficulty getting and keeping an erection (impotence) or problems having sex
- Difficulty in swallowing.

Rare side effects (may affect up to 1 in 1000 people)

- Constricted pupils (miosis)
- Stopping breathing from time to time (apnoea).

Frequency not known (frequency cannot be estimated from the available data)

- Lack of male sex hormones (androgen deficiency)
- Delirium (symptoms may include a combination of agitation, restlessness, disorientation, confusion, fear, seeing or hearing things that are not really there, sleep disturbance, nightmares)
- You can become dependent on Fental Matrix (see section 2).

You may notice rashes, redness or slight itching of the skin at the site of the patch. This is usually mild and disappears after you have removed the patch. If it does not, or if the patch irritates your skin

badly, tell your doctor.

Repeated use of the patches can make the medicine become less effective (you become 'tolerant' to it) or you can become dependent on it.

If you switch from a different painkiller to Fental Matrix or if you suddenly stop using Fental Matrix, you may notice withdrawal effects such as sickness, feeling sick, diarrhoea, anxiety or shivering. Tell your doctor if you notice any of these effects.

There have been reports also of newborn infants experiencing withdrawal effects after their mothers have used Fental Matrix for a long time during pregnancy.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system: HPRA Pharmacovigilance; website; www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fental Matrix

Keep all patches (used and unused) out of the sight and reach of children.

Store this medicine in a safe and secure place, where other people cannot access it. It can cause serious harm and be fatal to people who may take this medicine by accident, or intentionally when it has not been prescribed for them.

Do not use this medicine after the expiry date which is stated on the carton and sachet after "EXP". The expiry date refers to the last day of that month.

Store in the original package.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

A used or unused patch accidentally sticking to another person, especially a child, may be fatal. Used patches should be folded firmly in half so that the sticky side of the patch sticks to itself. Then they should be safely discarded by putting them back into the original sachet and stored out of sight and reach of other people, especially children, until safely disposed.

6. Contents of the pack and other information

What Fental Matrix contains

The active substance is fentanyl.

Each transdermal patch (10.5 cm² absorption surface area) contains:

5.78 mg fentanyl (which corresponds to a release rate of 25 microgram/hour of fentanyl)

Each transdermal patch (21 cm² absorption surface area) contains:

11.56 mg fentanyl (which corresponds to a release rate of 50 microgram/hour of fentanyl)

Each transdermal patch (31.5 cm² absorption surface area) contains:

17.34 mg fentanyl (which corresponds to a release rate of 75 microgram/hour of fentanyl)

Each transdermal patch (42 cm² absorption surface area) contains:

23.12 mg fentanyl (which corresponds to a release rate of 100 microgram/hour of fentanyl)

The other ingredients are:

Protective film (removed before the patch is stuck onto the skin):

Poly(ethylene terephthalate) foil, siliconised

Self-adhesive matrix layer:

Colophonium resin (hydrogenated)

Poly(2-ethylhexyl acrylate-co-vinyl acetate)
Soya-bean oil, refined
Water impermeable cover film:
Poly(ethylene terephthalate) foil
Printing ink.

What Fental Matrix looks like and contents of the pack

Fental Matrix is a transparent rounded oblong patch with an adhesive membrane and covered with a protective film (to be removed prior to the application of the patch) so that it can be stuck onto the skin.

Fental Matrix Transdermal patch is available in packs with 3, 5, 7, 10, 14 and 20 transdermal patches.

Hospital packs with 5 transdermal patches.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Rowex Ltd., Bantry, Co. Cork, Ireland.

Manufacturer

Hexal AG, Industriestr. 25, 83607 Holzkirchen, Germany.

This medicinal product is authorised in the Member States of the EEA under the following names:

- DE: Fentapatch Matrix 25 µg/h, transdermales Pflaster
- Fentapatch Matrix 50 µg/h, transdermales Pflaster
- Fentapatch Matrix 75 µg/h, transdermales Pflaster
- Fentapatch Matrix 100 µg/h, transdermales Pflaster
- IE: Fental Matrix 25 micrograms/hour Transdermal patch
- Fental Matrix 50 micrograms/hour Transdermal patch
- Fental Matrix 75 micrograms/hour Transdermal patch
- Fental Matrix 100 micrograms/hour Transdermal patch

This leaflet was last revised in 12/2025.