

Package leaflet: Information for the user

Memantine Sandoz 10 mg Film-coated tablets

Memantine hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Memantine Sandoz is and what it is used for
2. What you need to know before you take Memantine Sandoz
3. How to take Memantine Sandoz
4. Possible side effects
5. How to store Memantine Sandoz
6. Contents of the pack and other information

1. What Memantine Sandoz is and what it is used for

How does Memantine Sandoz work

Memantine Sandoz belongs to a group of medicines known as anti-dementia medicines. Memory loss in Alzheimer's disease is due to a disturbance of message signals in the brain. The brain contains so-called N-methyl-D-aspartate (NMDA)-receptors that are involved in transmitting nerve signals important in learning and memory. Memantine Sandoz belongs to a group of medicines called NMDA-receptor antagonists. Memantine Sandoz acts on these NMDA-receptors improving the transmission of nerve signals and the memory.

What is Memantine Sandoz used for

Memantine Sandoz is used for the treatment of patients with moderate to severe Alzheimer's disease.

2. What you need to know before you take Memantine Sandoz

Do not take Memantine Sandoz

- if you are allergic to memantine hydrochloride or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Memantine Sandoz:

- if you have a history of epileptic seizures
- if you have recently experienced a myocardial infarction (heart attack), or if you are suffering from congestive heart failure or from an uncontrolled hypertension (high blood pressure)

In these situations the treatment should be carefully supervised, and the clinical benefit of Memantine Sandoz reassessed by your doctor on a regular basis.

If you suffer from renal impairment (kidney problems), your doctor should closely monitor your kidney function and if necessary adapt the memantine doses accordingly.

The use of medicinal products called amantadine (for the treatment of Parkinson's disease), ketamine (a substance generally used as an anaesthetic), dextromethorphan (generally used to treat cough) and other NMDA-antagonists at the same time should be avoided.

Children and adolescents

Memantine Sandoz is not recommended for children and adolescents under the age of 18 years.

Other medicines and Memantine Sandoz

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, Memantine Sandoz may change the effects of the following medicines and their dose may need to be adjusted by your doctor:

- amantadine, ketamine, dextromethorphan
- dantrolene, baclofen
- cimetidine, ranitidine, procainamide, quinidine, quinine, nicotine
- hydrochlorothiazide (or any combination with hydrochlorothiazide)
- anticholinergics (substances generally used to treat movement disorders or intestinal cramps)
- anticonvulsants (substances used to prevent and relieve seizures)
- barbiturates (substances generally used to induce sleep)
- dopaminergic agonists (substances such as L-dopa, bromocriptine)
- neuroleptics (substances used in the treatment of mental disorders)
- oral anticoagulants

If you go into hospital, let your doctor know that you are taking Memantine Sandoz.

Memantine Sandoz with food and drink

You should inform your doctor if you have recently changed or intend to change your diet substantially (e.g. from normal diet to strict vegetarian diet) or if you are suffering from states of renal tubular acidosis (RTA, an excess of acid-forming substances in the blood due to renal dysfunction (poor kidney function)) or severe infections of the urinary tract (structure that carries urine), as your doctor may need to adjust the dose of your medicine.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. **Pregnancy**
The use of Memantine Sandoz in pregnant women is **not recommended**.

Breast-feeding

Women taking Memantine Sandoz should not breast-feed.

Driving and using machines

Your doctor will tell you whether your illness allows you to drive and to use machines safely. Also, Memantine Sandoz may change your reactivity, making driving or operating machinery inappropriate.

Memantine Sandoz contains lactose.

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Memantine Sandoz

Always take Memantine Sandoz exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Dosage

The **recommended dose** for adults and elderly patients is 20 mg once a day.

In order to reduce the risk of side effects this dose is achieved gradually by the following daily treatment scheme:

Period of intake	Dosage once daily
week 1	half a 10 mg tablet
week 2	one 10 mg tablet
week 3	one and a half 10 mg tablets
week 4 and beyond	two 10 mg tablets


Dosage in patients with impaired kidney function

If you have impaired kidney function, your doctor will decide upon a dose that suits your condition. In this case, monitoring of your kidney function should be performed by your doctor at specified intervals.

Administration

Memantine Sandoz should be administered orally once a day. To benefit from your medicine you should take it regularly every day at the same time of the day. The tablets should be swallowed with some water. The tablets can be taken with or without food.

The tablet can be divided into 2 equal doses, as seen in the picture. If required, take the tablet in your hands and press the thumbs downwards, over the index fingers.



Duration of treatment

Continue to take Memantine Sandoz as long as it is of benefit to you. Your doctor should assess your treatment on a regular basis.

If you take more Memantine Sandoz than you should

- In general, taking too much Memantine Sandoz should not result in any harm to you. You may experience increased symptoms as described in section 4. “Possible side effects”.
- If you take a large overdose of Memantine Sandoz, contact your doctor or get medical advice, as you may need medical attention.

If you forget to take Memantine Sandoz

- If you find you have forgotten to take your dose of Memantine Sandoz, wait and take your next dose at the usual time.
- Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

In general, the observed side effects are mild to moderate.

Common, may affect up to 1 in 10 people:

- headache, sleepiness, constipation, elevated liver function tests, dizziness, balance disorders, shortness of breath, high blood pressure and drug hypersensitivity

Uncommon, may affect up to 1 in 100 people:

- tiredness, fungal infections, confusion, hallucinations, vomiting, abnormal gait, heart failure and venous blood clotting (thrombosis/thromboembolism)

Very Rare, may affect up to 1 in 10,000 people:

- seizures

Not known, frequency cannot be estimated from the available data:

- inflammation of the pancreas, inflammation of the liver (hepatitis) and psychotic reactions

Alzheimer's disease has been associated with depression, suicidal ideation and suicide. These events have been reported in patients treated with this medicine.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Memantine Sandoz

Keep this medicine out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this medicine after the expiry date which is stated on the blister or bottle label and carton after "EXP". The expiry date refers to the last day of that month. Once opened, the contents of the bottle should be used within 6 months.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Memantine Sandoz contains

The active substance is memantine.

Each film-coated tablet contains 10 mg memantine hydrochloride which is equivalent to 8.31 mg of memantine.

The other ingredients are:

Tablet core: lactose monohydrate, cellulose microcrystalline, silica colloidal anhydrous, magnesium stearate.

Tablet coating: hypromellose (E464), lactose monohydrate, macrogol, triacetin and titanium dioxide (E 171).

What Memantine Sandoz looks like and contents of the pack

Memantine Sandoz are white film-coated tablets of oval shape (6.1 x 11.6 mm) with a breaking line on both sides.

The tablet can be divided into equal doses.

This medicine is available in transparent PVC-Aclar/Aluminium and/or transparent PVC-PVDC/Aluminium blister packs containing 7, 10, 14, 18, 20, 22, 28, 30, 40, 42, 45, 48, 49, 49x1, 50, 56, 56x1, 60, 70, 84, 90, 96, 98, 98x1, 100, 100x1, 112, 980(10x98) or 1000(20x50) film-coated tablets.

This medicine is available in HPDE bottles with PP screw cap with tamper-evident ring and desiccant containing 28, 30, 56, 98, 100 or 112 film-coated tablets.

This medicine is available in amber glass bottles type III containing 20 ml, 50 ml or 100 ml of oral solution. The bottles are closed with HDPE screw cap with tamper-evident ring and packed in a carton box together with an oral syringe (LDPE and PS) connected to a press-in bottle adapter (LDPE). The oral syringe has a main graduation in steps of 0.5 ml and 5 mg (0.5, 1, 1.5 2 ml respectively 5, 10, 15, 20 mg) and a fine graduation in steps of 0.1 ml (= 1 mg).

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturers

Marketing authorization holder:

Rowex Ltd., Bantry, Co. Cork, Ireland.

Manufacturers:

Lek Pharmaceuticals d.d., Verovškova 57, 1526 Ljubljana, Slovenia.

LEK S.A., ul. Domaniewska 50 C, 02-672 Warszawa, Poland.

Salutas Pharma GmbH, Otto-von-Guericke-Allee 1, 39179 Barleben, Germany.

S.C. Sandoz, S.R.L., Str. Livezeni nr. 7A, RO-540472 Targu-Mures, Romania.

This medicinal product is authorised in the Member States of the EEA under the following names:

NL	Memantine Sandoz 10 mg, filmomhulde tabletten
AT	Memantin Sandoz 10 mg - Filmtabletten
BE	Memantine Sandoz 10 mg filmomhulde tabletten
CY	Memantine Sandoz 10 mg tabs
CZ	Memantin Sandoz 10 mg
DE	Memantin HEXAL 10 mg Filmtabletten
DK	Memantine Sandoz
EL	Memantine/Sandoz
ES	Memantina Sandoz 10 mg comprimidos recubiertos con película EFG
FI	Memantine Sandoz 10 mg tabletti, kalvopäällysteinen
FR	Memantine Sandoz 10 mg, comprimé pelliculé sécable
IE	Memantine Sandoz 10 mg Film-coated tablets
IS	Memantine Sandoz
LU	Memantine Sandoz 10 mg comprimés pelliculés
MT	Memantine 10 mg Film-coated Tablets
NO	Memantine Sandoz
PT	Memantina Sandoz
SE	Memantine Sandoz
UK	Memantine 10 mg Film-coated Tablets

This leaflet was last revised in 09/2015.