

Front

53 mm

You should immediately contact your doctor if you think you are having a hypersensitivity reaction to Abacavir/Lamivudine Rowex. Write your doctor's details below:

Doctor:

Tel:

If your doctor is not available, you must urgently seek alternative medical advice (e.g. the emergency unit of the nearest hospital).

For general Abacavir/Lamivudine Rowex information enquiries, contact Rowex Ltd. Tel.: 027 50077

IMPORTANT - ALERT CARD
Abacavir/Lamivudine Rowex Tablets
(abacavir / lamivudine)
Carry this card with you at all times

Since Abacavir/Lamivudine Rowex contains abacavir some patients taking Abacavir/Lamivudine Rowex may develop a hypersensitivity reaction (serious allergic reaction) which **can be life-threatening** if treatment with Abacavir/Lamivudine Rowex is continued.

85 mm

Back

CONTACT YOUR DOCTOR IMMEDIATELY for advice on whether you should stop taking Abacavir/Lamivudine Rowex if:

1) you get a skin rash OR
2) you get one or more symptoms from at least TWO of the following groups

- fever
- shortness of breath, sore throat or cough
- nausea or vomiting or diarrhoea or abdominal pain
- severe tiredness or achiness or generally feeling ill

If you have discontinued Abacavir/Lamivudine Rowex due to this reaction, **YOU MUST NEVER TAKE** Abacavir/Lamivudine Rowex, or any medicine containing abacavir (e.g. Zigen, Trizivir, Trimeq) again as **within hours** you may experience a life-threatening lowering of your blood pressure or death.

(see reverse of card)