

How to take Emtricitabine/tenofovir disoproxil Rowex

Always take Emtricitabine/tenofovir disoproxil Rowex exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- The recommended dose is one Emtricitabine/tenofovir disoproxil Rowex tablet each day. Whenever possible, Emtricitabine/tenofovir disoproxil Rowex should be taken with food. Take Emtricitabine/tenofovir disoproxil Rowex every day, not just when you think you have been at risk of getting HIV infection
- Do not miss any doses of Emtricitabine/tenofovir disoproxil Rowex, or stop taking it unless told to by your doctor. Missing doses may increase your risk of getting HIV infection. If you miss a dose of Emtricitabine/tenofovir disoproxil Rowex and you notice:
 - Within 12 hours of the time you usually take Emtricitabine/tenofovir disoproxil Rowex tablet, take the tablet preferably with food as soon as possible. Then take the next dose at your usual time
 - 12 hours or more after the time you usually take Emtricitabine/tenofovir disoproxil Rowex forget about the missed dose. Wait and take the next dose, preferably with food, at your usual time
- If you vomit less than 1 hour after taking Emtricitabine/tenofovir disoproxil Rowex, take another tablet. You do not need to take another tablet if you were sick more than 1 hour after taking Emtricitabine/tenofovir disoproxil Rowex

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. Your doctor may ask if you want to participate in a study (called the Antiretroviral Pregnancy Registry) which monitors the outcomes of women taking this medicine in pregnancy.

Side effects

Like all medicines, Emtricitabine/tenofovir disoproxil Rowex can cause side effects, although not everybody gets them.

Read the patient information leaflet provided in the box containing your medicine for a full list of possible side effects.

Emtricitabine/tenofovir disoproxil Rowex can cause serious side effects, including new or worsening of kidney problems and bone problems. Before and during treatment, your doctor may order blood tests to measure kidney function. Tell your doctor if you have had kidney disease, or if tests have shown problems with your kidneys, or **if you think you have any of the serious side effects listed in the patient information leaflet.**

Very common side effects in people who take Emtricitabine/tenofovir disoproxil Rowex include diarrhoea, being sick (vomiting), feeling sick (nausea), dizziness, headache, rash and feeling weak.

Tell your doctor if you have any side effect that bothers you or does not go away. This includes any possible side effects not listed in the patient information leaflet.

How to store Emtricitabine/tenofovir disoproxil Rowex

- Keep this medicine out of the sight and reach of children
- Do not use this medicine after the expiry date which is stated on the bottle and carton after {EXP}. The expiry date refers to the last day of that month
- Store in the original package in order to protect from moisture. Keep the bottle tightly closed

! IMPORTANT INFORMATION

Important information about emtricitabine/tenofovir disoproxil Rowex to Reduce the Risk of Getting Human Immunodeficiency Virus (HIV) Infection

Information for individuals who have been prescribed Emtricitabine/tenofovir disoproxil Rowex for Pre-Exposure Prophylaxis (PrEP)

Please report any adverse events suspected to be caused by the use of Emtricitabine/tenofovir disoproxil Rowex to Rowex Ltd, Bantry, Co.Cork. Tel 027 50077; fax 027 50417

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in the patient information leaflet. You can also report side effects directly via the national reporting system:

Ireland

HPRA Pharmacovigilance

Earlsfort Terrace

IRL - Dublin

Tel: +353 1 6764971

Fax: +353 1 6762517

Website: www.hpra.ie

e-mail: medsafety@hpra.ie

This guide is intended for information only and should complement (not replace) the patient information leaflet supplied with your medicine. For a full list of side effects and other important information, please refer to the patient information leaflet provided in the box containing your medicine. Ask your healthcare provider if you have any questions.

PA0711/271/001

Date of preparation: October 2017

CCF 19519

What is Emtricitabine/tenofovir disoproxil Rowex for PrEP?

Emtricitabine/tenofovir disoproxil Rowex is a prescription medicine to help reduce the risk of getting HIV infection in adults, together with safer sex practices.

What you should know before and while taking Emtricitabine/tenofovir disoproxil Rowex to reduce your risk of getting HIV infection

You must be HIV negative to start taking Emtricitabine/tenofovir disoproxil Rowex for PrEP. Emtricitabine/tenofovir disoproxil Rowex can only help reduce your risk of getting HIV before you are infected. You must get tested to make sure you do not already have HIV. Do not take Emtricitabine/tenofovir disoproxil Rowex to reduce your risk unless you are confirmed to be HIV negative.

Tell your doctor about any flu-like illnesses either in the month before starting Emtricitabine/tenofovir disoproxil Rowex or at any time while taking Emtricitabine/tenofovir disoproxil Rowex. If you have engaged in sex practices that may put you at risk of getting HIV, these may be signs of HIV infection:

- TIREDNESS
- FEVER
- JOINT OR MUSCLE ACHES
- HEADACHE
- VOMITING OR DIARRHOEA
- RASH
- NIGHT SWEATS
- ENLARGED LYMPH NODES IN THE NECK OR GROIN

Take Emtricitabine/tenofovir disoproxil Rowex as prescribed by your doctor. Do not miss any doses of Emtricitabine/tenofovir disoproxil Rowex, or stop taking it, unless directed by a healthcare professional. Missing doses and erratic intake of doses will increase your risk of getting HIV and to possibly developing resistance to the drug in case you get infected with HIV

- Know your HIV status and the HIV status of your partners
- Get tested for HIV regularly (e.g. at least every 3 months)
- If you think you were infected with HIV, tell your doctor right away. They may want to do more tests to be sure you are still HIV negative
- **Just taking Emtricitabine/tenofovir disoproxil Rowex may not stop you getting HIV. You must continue using safer sex practises** while you are taking Emtricitabine/tenofovir disoproxil Rowex to reduce your risk of getting HIV:
 - Use condoms to reduce contact with semen, vaginal fluids, or blood and avoid being exposed to other sexually transmitted infections
 - Do not share personal items that can have blood or body fluids on them, such as toothbrushes and razor blades
 - Do not share or reuse needles or other injection or drug equipment
- **Take action to stay negative while taking Emtricitabine/tenofovir disoproxil Rowex for PrEP:**
 - Get tested for other sexually transmitted infections, such as syphilis and gonorrhoea. These infections make it easier for HIV to infect you
 - Get information and support to help reduce sexual behaviour that may put you at higher risk
- **Know your hepatitis B virus (HBV) infection status before starting Emtricitabine/tenofovir disoproxil Rowex:**

If you have HBV infection, there is a serious risk that liver problems may become worse after you stop taking Emtricitabine/tenofovir disoproxil Rowex