

Package leaflet: Information for the user

Tevanate Once Weekly 70 mg Tablets alendronic acid

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- It is particularly important to understand the information in section 3. before taking this medicine.

What is in this leaflet

1. What Tevanate Once Weekly 70 mg Tablets is and what it is used for
2. What you need to know before you take Tevanate Once Weekly 70 mg Tablets
3. How to take Tevanate Once Weekly 70 mg Tablets
4. Possible side effects
5. How to store Tevanate Once Weekly 70 mg Tablets
6. Contents of the pack and other information

1. What Tevanate Once Weekly 70 mg Tablets is and what it is used for

What is Tevanate Once Weekly 70 mg Tablets?

Tevanate Once Weekly 70 mg Tablets is a tablet containing the active substance alendronic acid (commonly called alendronate) and belongs to a group of non-hormonal medicines called bisphosphonates. Tevanate Once Weekly 70 mg Tablets prevents the loss of bone that occurs in women after they have been through the menopause, and helps to rebuild bone. It reduces the risk of spine and hip fractures.

What is Tevanate Once Weekly 70 mg Tablets used for?

Your doctor has prescribed Tevanate Once Weekly 70 mg Tablets to treat your osteoporosis. It reduces the risk of spine and hip fractures.

Tevanate Once Weekly 70 mg Tablets is a once weekly treatment.

What is osteoporosis?

Osteoporosis is a thinning and weakening of the bones. It is common in women after the menopause. At the menopause, the ovaries stop producing the female hormone, oestrogen, which helps to keep a woman's skeleton healthy. As a result, bone loss occurs and bones become weaker. The earlier a woman reaches the menopause, the greater the risk of osteoporosis.

Early on, osteoporosis usually has no symptoms. If left untreated, however, it can result in broken bones. Although these usually hurt, breaks in the bones of the spine may go unnoticed until they cause height loss. Broken bones can happen during normal, everyday activity, such as lifting, or from minor injury that would not generally break normal bone. Broken bones usually occur at the hip, spine, or wrist and can lead not only to pain but also to considerable problems like stooped posture ("dowager's hump") and loss of mobility.

How can osteoporosis be treated?

As well as your treatment with Tevanate Once Weekly 70 mg Tablets, your doctor may suggest you make changes to your lifestyle to help your condition, such as:

<i>Stopping smoking</i>	Smoking appears to increase the rate at which you lose bone and, therefore, may increase your risk of broken bones.
<i>Exercise</i>	Like muscles, bones need exercise to stay strong and healthy. Consult your doctor before you begin any exercise programme.
<i>Eating a balanced diet</i>	Your doctor can advise you about your diet or whether you should take any dietary supplements (especially calcium and Vitamin D).

2. What you need to know before you take Tevanate Once Weekly 70 mg Tablets

Do not take Tevanate Once Weekly 70 mg Tablets

- if you are allergic to alendronic acid or any of the other ingredients of this medicine (listed in section 6).
- if you have certain problems with your gullet (oesophagus – the tube that connects your mouth with your stomach) such as narrowing or difficulty swallowing.
- if you cannot stand or sit upright for at least 30 minutes.
- if your doctor has told you that you have low blood calcium.

If you think any of these apply to you, do not take the tablets. Talk to your doctor first and follow the advice given.

Warnings and precautions

Talk to your doctor or pharmacist before taking Tevanate Once Weekly 70 mg Tablets if

- you suffer from kidney problems,
- you have, or have recently had, any swallowing or digestive problems,
- your doctor has told you that you have Barrett's oesophagus (a condition associated with changes in the cells that line the lower oesophagus),
- you have been told you have trouble absorbing minerals in your stomach or intestines (malabsorption syndrome),
- you have been told you have low blood calcium,
- you have poor dental health, gum disease, a planned dental extraction or you don't receive routine dental care,
- you have cancer,
- you are undergoing chemotherapy or radiotherapy,
- you are taking angiogenesis inhibitors (such as bevacizumab, or thalidomide) which are used in the treatment of cancer,
- you are taking corticosteroids (such as prednisone or dexamethasone) which are used in the treatment of such conditions as asthma, rheumatoid arthritis, and severe allergies,
- you are or have been a smoker (as this may increase the risk of dental problems)

You may be advised to have a dental check-up before starting treatment with Tevanate Once Weekly 70 mg Tablets.

It is important to maintain good oral hygiene when being treated with Tevanate Once Weekly 70 mg Tablets. You should have routine dental check-ups throughout your treatment and you should contact your doctor or dentist if you experience any problems with your mouth or teeth such as loose teeth, pain or swelling.

Irritation, inflammation or ulceration of the gullet (oesophagus – the tube that connects your mouth with your stomach) often with symptoms of chest pain, heartburn, or difficulty or pain upon swallowing may occur, especially if patients do not drink a full glass of water and/or if they lie down less than 30 minutes after taking Tevanate Once Weekly 70 mg Tablets. These side effects may worsen if patients continue to take Tevanate Once Weekly 70 mg Tablets after developing these symptoms.

Children and adolescents

Tevanate Once Weekly 70 mg Tablets should not be given to children and adolescents less than 18 years of age.

Other medicines and Tevanate Once Weekly 70 mg Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

It is likely that calcium supplements, antacids, and some oral medicines will interfere with the absorption of Tevanate Once Weekly 70 mg Tablets if taken at the same time. Therefore, it is important that you follow the advice given in section 3.

Certain medicines for rheumatism or long-term pain called NSAIDs (e.g. acetylsalicylic acid or ibuprofen) might cause digestive problems. Therefore, caution should be used when these medicines are taken at the same time as Tevanate Once Weekly 70 mg Tablets.

Tevanate Once Weekly 70 mg Tablets with food and drink

It is likely that food and beverages (including mineral water) will make Tevanate Once Weekly 70 mg Tablets less effective if taken at the same time. Therefore, it is important that you follow the advice given in section 3.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. Tevanate Once Weekly 70 mg Tablets is only intended for use in postmenopausal women. You should not take Tevanate Once Weekly 70 mg Tablets if you are or think you may be pregnant or if you are breast-feeding.

Driving and using machines

There have been side effects (for example blurred vision, dizziness and severe bone, muscle or joint pain) reported with Tevanate Once Weekly 70 mg Tablets that may affect your ability to drive or operate machinery (see section 4).

Tevanate Once Weekly 70 mg Tablets contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially “sodium-free”.

3. How to take Tevanate Once Weekly 70 mg Tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Take one Tevanate Once Weekly 70 mg Tablets tablet once a week.

Follow these instructions carefully:

- 1) Choose the day of the week that best fits your schedule. Every week, take one Tevanate Once Weekly 70 mg Tablet on your chosen day

It is very important to follow instructions 2), 3), 4) and 5) to help the Tevanate Once Weekly 70 mg Tablets tablet reach your stomach quickly and help reduce the chance of irritating your gullet (oesophagus - the tube that connects your mouth with your stomach).

- 2) After getting up for the day and before taking any food, drink, or other medicine, swallow your Tevanate Once Weekly 70 mg Tablets tablet whole with a full glass of water only (not mineral water) (not less than 200 ml), so that Tevanate Once Weekly 70 mg Tablets is adequately absorbed.
 - Do not take with mineral water (still or sparkling).
 - Do not take with coffee or tea.

- Do not take with juice or milk.

Do not crush or chew the tablet or allow it to dissolve in your mouth because of the possibility of mouth ulceration.

- 3) Do not lie down - stay fully upright (sitting, standing or walking) - for at least 30 minutes after swallowing the tablet. Do not lie down until after your first food of the day.
- 4) Do not take Tevanate Once Weekly 70 mg Tablets at bedtime or before getting up for the day.
- 5) If you develop difficulty or pain upon swallowing, chest pain, or new or worsening heartburn, stop taking Tevanate Once Weekly 70 mg Tablets and contact your doctor.
- 6) After swallowing your Tevanate Once Weekly 70 mg Tablets tablet, wait at least 30 minutes before taking your first food, drink, or other medicine of the day, including antacids, calcium supplements and vitamins. Tevanate Once Weekly 70 mg Tablets is effective only if taken when your stomach is empty.

If you take more Tevanate Once Weekly 70 mg Tablets than you should

If you take too many tablets by mistake, drink a full glass of milk and contact your doctor immediately. Do not make yourself vomit and do not lie down.

If you forget to take Tevanate Once Weekly 70 mg Tablets

If you miss a dose, just take one tablet on the morning after you remember. *Do not take two tablets on the same day.* Return to taking one tablet once a week as originally scheduled on your chosen day. Do not take a double dose to make up for a forgotten tablet.

If you stop taking Tevanate Once Weekly 70 mg Tablets

It is important that you take Tevanate Once Weekly 70 mg Tablets for as long as your doctor prescribes the medicine. Since it is not known how long you should take Tevanate Once Weekly 70 mg Tablets, you should discuss the need to stay on this medicine with your doctor periodically to determine if Tevanate Once Weekly 70 mg Tablets is still right for you.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

See your doctor immediately if you notice any of the following side effects, which may be serious, and for which you may need urgent medical treatment:

Common (may affect up to 1 in 10 people)

- heartburn; difficulty swallowing; pain upon swallowing; ulceration of the gullet (oesophagus - the tube that connects your mouth with your stomach) which can cause chest pain, heartburn or difficulty or pain upon swallowing.

Rare (may affect up to 1 in 1,000 people)

- allergic reactions, such as hives, swelling of the face, lips, tongue and/or throat, possibly causing difficulty breathing or swallowing, severe skin reactions,
- pain in the mouth, and/or jaw, swelling or sores inside the mouth, numbness or a feeling of heaviness in the jaw, or loosening of a tooth. These could be signs of bone damage in the jaw (osteonecrosis) generally associated with delayed healing and infection, often following tooth extraction. Contact your doctor and dentist if you experience such symptoms,
- unusual fracture of the thigh bone particularly in patients on long-term treatment for osteoporosis may occur rarely. Contact your doctor if you experience pain, weakness or

discomfort in your thigh, hip or groin as this may be an early indication of a possible fracture of the thigh bone.

- bone, muscle and/or joint pain which is severe.

Other side effects include

Very common (may affect more than 1 in 10 people)

- bone, muscle and/or joint pain which is sometimes severe.

Common (may affect up to 1 in 10 people)

- joint swelling,
- abdominal pain, uncomfortable feeling in the stomach or belching after eating; constipation; full or bloated feeling in the stomach, diarrhoea, flatulence,
- hair loss, itching,
- headache, dizziness
- tiredness, swelling in the hands or legs.

Uncommon (may affect up to 1 in 100 people)

- nausea, vomiting,
- irritation or inflammation of the gullet (oesophagus – the tube that connects your mouth with your stomach) or stomach,
- black or tar-like stools,
- blurred vision; pain or redness in the eye,
- rash; redness of the skin,
- transient flu-like symptoms, such as aching muscles, generally feeling unwell and sometimes with fever usually at the start of treatment,
- taste disturbance.

Rare (may affect up to 1 in 1,000 people)

- symptoms of low blood calcium levels including muscle cramps or spasms and/or tingling sensation in the fingers or around the mouth,
- stomach or peptic ulcers (sometimes severe or with bleeding),
- narrowing of the gullet (oesophagus – the tube that connects your mouth with your stomach),
- rash made worse by sunlight,
- mouth ulcers.

Very rare (may affect up to 1 in 10,000 people)

- talk to your doctor if you have ear pain, discharge from the ear, and/or an ear infection. These could be signs of bone damage in the ear.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via HPRC Pharmacovigilance, Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Tevanate Once Weekly 70 mg Tablets

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after “EXP”. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Tevanate Once Weekly 70 mg Tablets contains

- The active ingredient is alendronic acid. Each tablet contains 70 mg alendronic acid (as sodium monohydrate).
- The other ingredients are microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

What Tevanate Once Weekly 70 mg Tablets looks like and contents of the pack

- The name of your medicine is Tevanate Once Weekly 70 mg Tablets. The product is available in pack sizes of 2, 4, 8, 12, 40 and 50 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Teva Pharma B.V.
Swensweg 5
2031GA Haarlem
The Netherlands

Manufacturer

Teva Pharmaceutical Works Private Limited Company, Pallagi út 13, 4042 Debrecen,
Hungary.

Pharmachemie B.V., Swensweg 5, Postbus 552, 2003 RN Haarlem, The Netherlands.

This medicinal product is authorised in the Member States of the EEA under the following names:

Belgium	Alendronate TEVA 70 mg tabletten
Czech Republic	Alendronate – Teva 70 mg
Denmark	Alendronat TEVA
Germany	Tevanate® 70 mg Tabletten
Greece	TEVANATE Δισκία 70 mg
Hungary	Trabecan-Teva 70 mg Tabletta
Ireland	Tevanate Once Weekly 70 mg Tablets
Italy	Alendronato Teva
Lithuania	TEVANATE 70 mg tablets
Netherlands	Alendroninezuur 70 PCH, tabletten 70 mg
Norway	Alendronat TEVA 70 mg tablett
Portugal	Acido Alendronico Teva 70 mg comprimidos
Spain	Acido Alendronico Teva 70 mg comprimidos EFG
Sweden	Alendronat Teva Veckotablett 70 mg tabletter
United Kingdom	Alendronic acid 70 mg Tablets

This leaflet was last revised in August 2021.