

# Omeprazole Rowa 10mg gastro-resistant capsules, hard Omeprazole Rowa 20mg gastro-resistant capsules, hard Omeprazole Rowa 40mg gastro-resistant capsules, hard

Omeprazole

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

## What is in this leaflet

1. What Omeprazole Rowa is and what it is used for
2. What you need to know before you take Omeprazole Rowa
3. How to take Omeprazole Rowa
4. Possible side effects
5. How to store Omeprazole Rowa
6. Contents of the pack and other information

Marketed by  
**ROWEX**<sup>®</sup>

## 1. What Omeprazole Rowa is and what it is used for

Omeprazole Rowa contains the active substance omeprazole. It belongs to a group of medicines called 'proton pump inhibitors'. They work by reducing the amount of acid that your stomach produces.

Omeprazole Rowa is used to treat the following conditions:

In adults:

- Gastro-esophageal reflux disease' (GERD). This is where acid from the stomach escapes into the gullet (the tube which connects your throat to your stomach) causing pain, inflammation and heartburn.
- Ulcers in the upper part of the intestine (duodenal ulcer) or stomach (gastric ulcer).
- Ulcers which are infected with bacteria called '*Helicobacter pylori*'. If you have this condition, your doctor may also prescribe antibiotics to treat the infection and allow the ulcer to heal.
- Ulcers caused by medicines called NSAIDs (Non-Steroidal Anti-Inflammatory Drugs). Omeprazole Rowa can also be used to stop ulcers from forming if you are taking NSAIDs.
- Too much acid in the stomach caused by a growth in the pancreas (Zollinger-Ellison syndrome).

In children:

*Children over 1 year of age and ≥ 10 kg*

- Gastro-oesophageal reflux disease' (GERD). This is where acid from the stomach escapes into the gullet (the tube which connects your throat to your stomach) causing pain, inflammation and heartburn.

In children, the symptoms of the condition can include the return of stomach contents into the mouth (regurgitation), being sick (vomiting) and poor weight gain.

*Children and adolescents over 4 years of age*

- Ulcers which are infected with bacteria called '*Helicobacter pylori*'. If your child has this condition, your doctor may also prescribe antibiotics to treat the infection and allow the ulcer to heal.

## 2. What you need to know before you take Omeprazole Rowa

### Do not take Omeprazole Rowa

- If you are allergic to omeprazole or any of the other ingredients of this medicine (listed in section 6)
- If you are allergic to medicines containing other proton pump inhibitors (e.g. pantoprazole, lansoprazole, rabeprazole, esomeprazole)
- If you are taking a medicine containing nelfinavir (used for HIV infection).

Do not take Omeprazole Rowa if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Omeprazole Rowa.

### Warnings and precautions

Talk to your doctor or pharmacist before taking Omeprazole Rowa.

Omeprazole Rowa may hide the symptoms of other diseases. Therefore, if any of the following happen to you before you start taking Omeprazole Rowa or while you are taking it, talk to your doctor straight away:

- You lose a lot of weight for no reason and have problems swallowing
- You get stomach pain or indigestion
- You begin to vomit food or blood
- You pass black stools (blood-stained faeces)
- You experience severe or persistent diarrhoea, as omeprazole has been associated with a small increase in infectious diarrhoea
- You have severe liver problems
- You have reduced body stores or risk factors for reduced vitamin B<sub>12</sub> and receive omeprazole long-term treatment. As with all acid reducing agents, omeprazole may lead to a reduced absorption of vitamin B<sub>12</sub>.
- If you have ever had a skin reaction after treatment with a medicine similar to Omeprazole Rowa that reduces stomach acid
- If you are due to have a specific blood test (Chromogranin A).

If you take Omeprazole Rowa on a long-term basis (longer than 1 year) your doctor will probably keep you under regular surveillance. You should report any new and exceptional symptoms and circumstances whenever you see your doctor.

Taking a proton pump inhibitor like Omeprazole Rowa, especially over a period of more than one year, may slightly increase your risk of fracture in the hip, wrist or spine. Tell your doctor if you have osteoporosis or if you are taking corticosteroids (which can increase the risk of osteoporosis).

If you get a rash on your skin, especially in areas exposed to the sun tell your doctor as soon as you can, as you may need to stop your treatment with Omeprazole Rowa. Remember to also mention any other ill-effects like pain in your joints.

### Children

Some children with chronic illnesses may require long-term treatment although it is not recommended. Do not give this medicine to children under 1 year of age or < 10 kg.

### Other medicines and Omeprazole Rowa

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. This includes medicines that you buy without a prescription. This is because Omeprazole Rowa can affect the way some

medicines work and some medicines can have an effect on Omeprazole Rowa.

Do not take Omeprazole Rowa if you are taking a medicine containing **nelfinavir** (used to treat HIV infection).

Tell your doctor or pharmacist if you are taking any of the following medicines:

- Ketoconazole, itraconazole, posaconazole or voriconazole (used to treat infections caused by a fungus)
- Digoxin (used to treat heart problems)
- Diazepam (used to treat anxiety, relax muscles or in epilepsy)
- Phenytoin (used in epilepsy). If you are taking phenytoin, your doctor will need to monitor you when you start or stop taking Omeprazole Rowa
- Medicines that are used to thin your blood, such as warfarin or other vitamin K blockers. Your doctor may need to monitor you when you start or stop taking Omeprazole Rowa.
- Rifampicin (used to treat tuberculosis)
- Atazanavir (used to treat HIV infection)
- Tacrolimus (in cases of organ transplantation)
- St John's wort (*Hypericum perforatum*) (used to treat mild depression)
- Cilostazol (used to treat intermittent claudication)
- Saquinavir (used to treat HIV infection)
- Clopidogrel (used to prevent blood clots (thrombi))
- Erlotinib (used to treat cancer)
- Methotrexate (a chemotherapy medicine used in high doses to treat cancer) – if you are taking a high dose of methotrexate, your doctor may temporarily stop your Omeprazole Rowa treatment.

If your doctor has prescribed the antibiotics amoxicillin and clarithromycin as well as omeprazole to treat ulcers caused by *Helicobacter pylori* infection, it is very important that you tell your doctor about any other medicines you are taking.

### Omeprazole Rowa with food and drink

See section 3.

### Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Omeprazole is excreted in breast milk but is not likely to influence the child when therapeutic doses are used. Your doctor will decide whether you can take Omeprazole Rowa if you are breast-feeding.

### Driving and using machines

Omeprazole Rowa is not likely to affect your ability to drive or use any tools or machines. Side effects such as dizziness and visual disturbances may occur (see section 4). If affected, you should not drive or operate machinery.

### Omeprazole Rowa contains sucrose

This medicine contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### Omeprazole Rowa contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per capsule that is to say essentially 'sodium free'.

## 3. How to take Omeprazole Rowa

Always take Omeprazole Rowa exactly as your doctor or pharmacist has told you. You should check with your doctor or pharmacist if you are not sure.

Your doctor will tell you how many capsules to take and how long to take them for. This will depend on your condition and how old you are.

The recommended dose is:

**Adults:**

To treat symptoms of GERD such as **heartburn and acid regurgitation:**

- If your doctor has found that your food pipe (gullet) has been slightly damaged, the usual dose is 20 mg once a day for 4-8 weeks. Your doctor may tell you to take a dose of 40 mg for a further 8 weeks if your gullet has not yet healed.
- The usual dose once the gullet has healed is 10 mg once a day.
- If your gullet has not been damaged, the usual dose is 10 mg once a day.

To treat **ulcers in the upper part of the intestine** (duodenal ulcer):

- The usual dose is 20 mg once a day for 2 weeks. Your doctor may tell you to take the same dose for a further 2 weeks if your ulcer has not yet healed.
- If the ulcer does not fully heal, the dose can be increased to 40 mg once a day for 4 weeks.

To treat **ulcers in the stomach** (gastric ulcer):

- The usual dose is 20 mg once a day for 4 weeks. Your doctor may tell you to take the same dose for a further 4 weeks if your ulcer has not yet healed.
- If the ulcer does not fully heal, the dose can be increased to 40 mg once a day for 8 weeks.

To **prevent the duodenal and stomach ulcers** from coming back:

- The usual dose is 10 mg or 20 mg once a day. Your doctor may increase the dose to 40 mg once a day.

To treat duodenal and stomach **ulcers caused by NSAIDs** (Non-Steroidal Anti-Inflammatory Drugs):

- The usual dose is 20 mg once a day for 4-8 weeks.

To **prevent duodenal and stomach ulcers** if you are taking **NSAIDs**:

- The usual dose is 20 mg once a day.

To treat ulcers caused by *Helicobacter pylori* infection and to stop them coming back:

- The usual dose is 20 mg Omeprazole Rowa twice a day for one week.
- Your doctor will also tell you to take two antibiotics among amoxicillin, clarithromycin and metronidazole.

To treat too much acid in the stomach caused by a **growth in the pancreas (Zollinger-Ellison syndrome)**:

- The recommended dose is 60 mg daily.
- Your doctor will adjust the dose depending on your needs and will also decide how long you need to take the medicine for.

Use in children and adolescents:

To treat symptoms of GERD such as **heartburn and acid regurgitation**:

- Children over 1 year of age and with a body weight of more than 10 kg may take Omeprazole Rowa. The dose for children is based on the child's weight and the doctor will decide the correct dose.

To treat **ulcers caused by *Helicobacter pylori*** infection and to stop them coming back:

- Children aged over 4 years may take Omeprazole Rowa. The dose for children is based on the child's weight and the doctor will decide the correct dose.
- Your doctor will also prescribe two antibiotics called amoxicillin and clarithromycin for your child.

#### Taking this medicine

- It is recommended that you take your capsules in the morning
- You can take your capsules with food or on an empty stomach
- Swallow your capsules whole with half a glass of water. Do not chew or crush the capsules. This is because the capsules contain coated pellets which stop the medicine from being broken down by the acid in your stomach. It is important not to damage the pellets.

#### What to do if you or your child have trouble swallowing the capsules

- If you or your child have trouble swallowing the capsules:
  - Open the capsules and swallow the contents directly with half a glass of water or put the contents into a glass of still (non-fizzy) water, any acidic fruit juice (e.g. apple, orange or pineapple) or apple sauce
  - Always stir the mixture just before drinking it (the mixture will not be clear). Then drink the mixture straight away or within 30 minutes.
  - To make sure that you have drunk all of the medicine, rinse the glass very well with half a glass of water and drink it. The solid pieces contain the medicine - do not chew or crush them.

#### If you take more Omeprazole Rowa than you should

If you take more Omeprazole Rowa than prescribed by your doctor, talk to your doctor or pharmacist straight away.

#### If you forget to take Omeprazole Rowa

If you forget to take a dose, take it as soon as you remember it. However, if it is almost time for your next dose, skip the missed dose. Do not take a double dose to make up for a forgotten dose.

#### If you stop taking Omeprazole Rowa

Do not stop taking Omeprazole Rowa without first talking to your doctor or pharmacist.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

#### If you notice any of the following rare but serious side effects, stop taking Omeprazole Rowa and contact a doctor immediately:

- Sudden wheezing, swelling of your lips, tongue and throat or body, rash, fainting or difficulties in swallowing (severe allergic reaction)
- Reddening of the skin with blisters or peeling. There may also be severe blisters and bleeding in the lips, eyes, mouth, nose and genitals. This could be 'Stevens-Johnson syndrome' or 'toxic epidermal necrolysis'.
- Yellow skin, dark urine and tiredness which can be symptoms of liver problems.

Other side effects include:

#### Common side effects (may affect up to 1 in 10 people)

- Headache
- Effects on your stomach or gut: diarrhoea, stomach pain, constipation, wind (flatulence)
- Feeling sick (nausea) or being sick (vomiting)
- Benign polyps in the stomach.

#### Uncommon side effects (may affect up to 1 in 100 people)

- Swelling of the feet and ankles
- Disturbed sleep (insomnia)
- Dizziness, tingling feelings such as "pins and needles", feeling sleepy
- Spinning feeling (vertigo)
- Changes in blood tests that check how the liver is working
- Skin rash, lumpy rash (hives) and itchy skin
- Generally feeling unwell and lacking energy
- Fracture of the wrist, hip or spine.

#### Rare side effects (may affect up to 1 in 1,000 people)

- Blood problems such as a reduced number of white cells or platelets. This can cause weakness, bruising or make infections more likely.
- Allergic reactions, sometimes very severe, including swelling of the lips, tongue and throat, fever, wheezing
- Low levels of sodium in the blood. This may cause weakness, being sick (vomiting) and cramps.
- Feeling agitated, confused or depressed
- Taste changes
- Eyesight problems such as blurred vision
- Suddenly feeling wheezy or short of breath (bronchospasm)
- Dry mouth
- An inflammation of the inside of the mouth
- An infection called "thrush" which can affect the gut and is caused by a fungus
- Liver problems, including jaundice which can cause yellow skin, dark urine, and tiredness
- Hair loss (alopecia)
- Skin rash on exposure to sunshine
- Joint pains (arthralgia) or muscle pains (myalgia)
- Severe kidney problems (interstitial nephritis)
- Increased sweating.

#### Very rare side effects (may affect up to 1 in 10,000 people)

- Changes in blood count including agranulocytosis (lack of white blood cells)
- Aggression
- Seeing, feeling or hearing things that are not there (hallucinations)
- Severe liver problems leading to liver failure and inflammation of the brain
- Sudden onset of a severe rash or blistering or peeling skin. This may be associated with a high fever and joint pains (Erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis).
- Muscle weakness
- Enlarged breasts in men.

#### Not known (frequency cannot be estimated from the available data)

- Inflammation in the gut (leading to diarrhoea)
- If you are on Omeprazole Rowa for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary contractions, disorientation, convulsions, dizziness, increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium.
- Rash, possibly with pain in the joints.

Omeprazole Rowa may in very rare cases affect the white blood cells leading to immune deficiency. If you have an infection with symptoms such as fever with a **severely** reduced general condition or fever with symptoms of a local infection such as pain in the neck, throat or mouth or difficulties in urinating, you must consult your doctor as soon as possible so that a lack of white blood cells (agranulocytosis) can be ruled out by a blood test. It is important for you to give information about your medicine at this time.

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance; website: [www.hpra.ie](http://www.hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Omeprazole Rowa

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, bottle label or blister after "EXP:". The expiry date refers to the last day of that month.

Store below 25°C.

OPA/Al/PVC-Al blister: Store in the original package to protect from moisture.

HDPE bottle: Keep the bottle tightly closed to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

#### What Omeprazole Rowa contains:

- The active substance is omeprazole. Omeprazole Rowa capsules contain 10 mg, 20 mg or 40 mg of omeprazole.
- The other ingredients are:
  - *Capsule content*: sugar spheres (consisting of maize starch and sucrose), sodium laurilsulfate, disodium phosphate, mannitol, hypromellose 6 cP, macrogol 6000, talc, polysorbate 80, titanium dioxide (E171), and methacrylic acid-ethylacrylate copolymer (1:1).
  - *Capsule shell*: gelatin. The 10 and 20 mg capsules also contain the colouring agents quinoline yellow (E104) and titanium dioxide (E171). The 40 mg capsules contain indigo carmine (E132) and titanium dioxide (E171).

#### What Omeprazole Rowa looks like and contents of the pack

- Omeprazole Rowa 10 mg: Opaque yellow capsule containing off-white to cream-white spherical microgranules
- Omeprazole Rowa 20 mg: Opaque yellow capsule containing off-white to cream-white spherical microgranules
- Omeprazole Rowa 40 mg: Opaque blue and opaque white capsule containing off-white to cream white spherical microgranules.

The capsules are supplied in blisters of 7, 14, 15, 28, 30, 50, 56, 60, 90, 100, 140, 280 and 500 capsules and in HDPE bottles of 28, 30 and 100 capsules (only 10 mg strength), 14, 28, 30 and 100 capsules (only 40 mg strength) and bottles of 28, 30, 60 (2x30) and 100 capsules (only 20 mg strength).

Not all pack sizes may be marketed.

#### Marketing Authorisation Holder and Manufacturer:

##### Marketing Authorisation Holder:

Rowa Pharmaceuticals Ltd., Bantry, Co. Cork, Ireland.

##### Manufacturer

LABORATORIOS LICONSA, S.A.  
Av. Miralcampo, N° 7, Polígono Industrial Miralcampo  
19200 Azuqueca de Henares (Guadalajara)  
SPAIN

#### This medicinal product is authorised in the Member States of the EEA under the following names:

Belgium:	Omeprazol Aurobindo harde maagsapresistente capsules /gélules gastro-resistantes/ magensaftresistente Hartkapseln
Ireland:	Omeprazole Rowa 10/20/40 mg gastro-resistant capsules
Luxembourg:	Omeprazol Aurobindo gélules gastro-resistantes
The Netherlands:	Omeprazol capsules 10/20/40 mg, maagsapresistente capsules

This leaflet was last revised in August 2022.

I.M. L/969a-c 08-22

<b>Artwork Checklist</b>	<b>Job Description</b> Omeprazole Rowia 10mg 20mg 40mg PIL RCF15593, Var. 55	<b>Version Number</b> 03	<b>Size</b> 140 x 540 mm	<b>Inks</b> Black								<b>ACCF</b> 24867
	<b>Market</b> IM	<b>Proof Date</b> 04/08/2022	<b>Cutter Ref.</b> 231	<b>Colour</b> Black	keyline							
				Front								
				Back								