

Package leaflet: Information for the patient

Ramipril Teva Pharma 1.25 mg Tablets
Ramipril Teva Pharma 2.5 mg Tablets
Ramipril Teva Pharma 5 mg Tablets
Ramipril Teva Pharma 10 mg Tablets

ramipril

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Ramipril Teva Pharma is and what it is used for
2. What you need to know before you take Ramipril Teva Pharma
3. How to take Ramipril Teva Pharma
4. Possible side effects
5. How to store Ramipril Teva Pharma
6. Contents of the pack and other information

1. What Ramipril Teva Pharma is and what it is used for

Ramipril Teva Pharma contains a medicine called ramipril. This belongs to a group of medicines called ACE inhibitors (Angiotensin Converting Enzyme Inhibitors).

Ramipril Teva Pharma works by:

- Decreasing your body's production of substances that could raise your blood pressure
- Making your blood vessels relax and widen
- Making it easier for your heart to pump blood around your body.

Ramipril Teva Pharma can be used:

- To treat high blood pressure (hypertension)
- To reduce the risk of you having a heart attack or stroke
- To reduce the risk or delay the worsening of kidney problems (whether or not you have diabetes)
- To treat your heart when it cannot pump enough blood to the rest of your body (heart failure)
- As treatment following heart attack (myocardial infarction) complicated with heart failure.

2. What you need to know before you take Ramipril Teva Pharma

Do not take Ramipril Teva Pharma

- If you are allergic to ramipril, any other ACE inhibitor medicine or any of the other ingredients of this medicine (listed in section 6).
Signs of an allergic reaction may include a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue

- If you have ever had a serious allergic reaction called “angioedema”. The signs include itching, hives (urticaria), red marks on the hands, feet and throat, swelling of the throat and tongue, swelling around the eyes and lips, difficulty breathing and swallowing
- If you are having dialysis or any other type of blood filtration. Depending on the machine that is used, Ramipril Teva Pharma may not be suitable for you
- If you have kidney problems where the blood supply to your kidney is reduced (renal artery stenosis)
- During the last 6 months of pregnancy. (see section “Pregnancy and breast-feeding”)
- If your blood pressure is abnormally low or unstable. Your doctor will need to make this assessment.
- If you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren
- If you have taken or are currently taking sacubitril/valsartan, a medicine used to treat a type of long-term (chronic) heart failure in adults, as the risk of angioedema (rapid swelling under the skin in an area such as the throat) is increased.

Do not take Ramipril Teva Pharma if any of the above applies to you. If you are not sure, talk to your doctor before taking Ramipril Teva Pharma.

Warnings and precautions

Talk to your doctor or pharmacist before taking Ramipril Teva Pharma

- If you have heart, liver or kidney problems
- If you have lost a lot of body salts or fluids (through being sick (vomiting), having diarrhoea, sweating more than usual, being on a low salt diet, taking diuretics (water tablets) for a long time or having had dialysis)
- If you are going to have treatment to reduce your allergy to bee or wasp stings (desensitization)
- If you are taking any of the following medicines used to treat high blood pressure:
 - an angiotensin II receptor blocker (ARBs) (also known as sartans - for example valsartan, telmisartan, irbesartan), in particular if you have diabetes-related kidney problems.
 - aliskiren

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.

See also information under the heading “Do not take *Ramipril Teva Pharma*”

- If you are going to receive an anaesthetic. This may be given for an operation or any dental work. You may need to stop your Ramipril Teva Pharma treatment one day beforehand; ask your doctor for advice
- If you have high amounts of potassium in your blood (shown in blood test results)
- If you are taking any of the following medicines, the risk of angioedema, a serious allergic reaction (rapid swelling under the skin in an area such as the throat), may be increased:
 - racecadotril, a medicine used to treat diarrhoea;
 - medicines used to prevent organ transplant rejection and for cancer (e.g., temsirolimus, sirolimus, everolimus);
 - vildagliptin, a medicine used to treat diabetes.
- If you have collagen vascular disease such as scleroderma or systemic lupus erythematosus
- You must tell your doctor if you think that you are (or might become) pregnant. Ramipril Teva Pharma is not recommended in the first 3 months of pregnancy, and may cause serious harm to your baby after 3 months of pregnancy (see section “Pregnancy and breast-feeding”).

Children and adolescents

Ramipril Teva Pharma is not recommended for use in children and adolescents below 18 years of age because safety and efficacy of Ramipril Teva Pharma in children has not yet been established.

If any of the above applies to you (or you are not sure), talk to your doctor before taking Ramipril Teva Pharma.

Other medicines and Ramipril Teva Pharma

Tell your doctor or pharmacist if you are taking/using, have recently taken/used or might take/use any other medicines. This is because Ramipril Teva Pharma can affect the way some other medicines work. Also some medicines can affect the way Ramipril Teva Pharma works.

Tell your doctor if you are taking any of the following medicines. They can make Ramipril Teva Pharma work less well:

- Medicines used to relieve pain and inflammation (e.g. Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) such as ibuprofen or indometacin and aspirin)
- Medicines used for the treatment of low blood pressure, shock, cardiac failure, asthma or allergies such as ephedrine, noradrenaline or adrenaline. Your doctor will need to check your blood pressure.

Tell your doctor if you are taking any of the following medicines. They can increase the chance of getting side effects if you take them with Ramipril Teva Pharma:

- Sacubitril/valsartan – used for treating a type of long term (chronic) heart failure in adults (see section 2 ‘Do not take Ramipril Teva Pharma’)
- Medicines used to relieve pain and inflammation (e.g. Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) such as ibuprofen or indometacin and aspirin)
- Medicines for cancer (chemotherapy)
- Diuretics (water tablets) such as furosemide
- Potassium supplements (including salt substitutes), potassium-sparing diuretics such as spironolactone, triamterene, amiloride and other medicines that can increase the amount of potassium in your blood (e.g. trimethoprim and co-trimoxazole for infections caused by bacteria; ciclosporin, an immunosuppressant medicine used to prevent organ transplant rejection; and heparin, a medicine used to thin blood to prevent clots)
- Steroid medicines for inflammation such as prednisolone
- Allopurinol (used to lower the uric acid in your blood)
- Procainamide (for heart rhythm problems)
- Medicines which are most often used to avoid rejection of transplanted organs (sirolimus, everolimus and other medicines belonging to the class of mTOR inhibitors). See section “Warnings and precautions”
- Vildagliptin (used for treating type 2 diabetes)
- Racecadotril (used against diarrhoea).

Your doctor may need to change your dose and/or to take other precautions:

- If you are taking an angiotensin II receptor blocker (ARB) or aliskiren (see also information under the headings “Do not take *Ramipril Teva Pharma*” and “Warnings and precautions”)

Tell your doctor if you are taking any of the following medicines. They may be affected by Ramipril Teva Pharma:

- Medicines for diabetes such as oral glucose lowering medicines and insulin. Ramipril Teva Pharma may lower your blood sugar amounts. Check your blood sugar amounts closely while taking Ramipril Teva Pharma
- Lithium (for mental health problems). Ramipril Teva Pharma may increase the amount of lithium in your blood. Your lithium amount will need to be closely checked by your doctor.

If any of the above applies to you (or you are not sure), talk to your doctor before taking Ramipril Teva Pharma.

Ramipril Teva Pharma with food and alcohol

- Drinking alcohol with Ramipril Teva Pharma may make you feel dizzy or light-headed. If you are concerned about how much you can drink while you are taking Ramipril Teva Pharma, discuss this with your doctor. Medicines used to reduce blood pressure and alcohol may mutually amplify their effect, which may cause an excessive reduction of blood pressure.
- Ramipril Teva Pharma may be taken with or without food.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

You must tell your doctor if you think that you are (or might become) pregnant. You should not take Ramipril Teva Pharma in the first 12 weeks of pregnancy, and you must not take them at all after the 13th week as their use during pregnancy may possibly be harmful to the baby.

If you become pregnant while on Ramipril Teva Pharma, tell your doctor immediately. A switch to a suitable alternative treatment should be carried out in advance of a planned pregnancy.

Breast-feeding

You should not take Ramipril Teva Pharma if you are breast-feeding.

Driving and using machines

You may feel dizzy, while taking Ramipril Teva Pharma. This is more likely to happen when you start taking Ramipril Teva Pharma or start taking a higher dose. If this happens, do not drive or use any tools or machines.

Ramipril Teva Pharma contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Ramipril Teva Pharma contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Ramipril Teva Pharma

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

Treatment of high blood pressure

- The usual starting dose is 1.25 mg or 2.5 mg once daily.
- Your doctor will adjust the amount you take until your blood pressure is controlled.
- The maximum dose is 10 mg once daily.
- If you are already taking diuretics (water tablets), your doctor may stop or reduce the amount of the diuretic you take before beginning treatment with Ramipril Teva Pharma.

To reduce the risk of you having a heart attack or stroke

- The usual starting dose is 2.5 mg once daily.
- Your doctor may then decide to increase the amount you take.
- The usual dose is 10 mg once daily.

Treatment to reduce or delay the worsening of kidney problems

- You may be started on a dose of 1.25 mg or 2.5 mg once daily.
- Your doctor will adjust the amount you are taking.
- The usual dose is 5 mg or 10 mg once daily.

Treatment of heart failure

- The usual starting dose is 1.25 mg once daily.
- Your doctor will adjust the amount you take.
- The maximum dose is 10 mg daily. Two administrations per day are preferable.

Treatment after you have had a heart attack

- The usual starting dose is 1.25 mg once daily to 2.5 mg twice daily.
- Your doctor will adjust the amount you take.

- The usual dose is 10 mg daily. Two administrations per day are preferable.

Elderly

- Your doctor will reduce the initial dose and adjust your treatment more slowly.

Taking this medicine

- Take this medicine by mouth at the same time of the day each day.
- Swallow the tablets with liquid.
- Do not crush or chew the tablets.

2.5 mg/5 mg/10 mg

- The tablet can be divided into equal doses.

If you take more Ramipril Teva Pharma than you should

Tell a doctor or go to the nearest hospital casualty department straight away. Do not drive to the hospital, get somebody else to take you or call for an ambulance. Take the medicine pack with you. This is so the doctor knows what you have taken.

If you forget to take Ramipril Teva Pharma

- If you miss a dose, take your normal dose when it is next due.
- Do not take a double dose to make up for a forgotten tablet.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Ramipril Teva Pharma and see a doctor straight away, if you notice any of the following serious side effects - you may need urgent medical treatment:

- Swelling of the face, lips or throat which make it difficult to swallow or breathe, as well as itching and rashes. This could be a sign of a severe allergic reaction to Ramipril Teva Pharma
- Severe skin reactions including rash, ulcers in your mouth, worsening of a pre-existing skin disease, reddening, blistering or detachment of skin (such as Stevens-Johnson syndrome, toxic epidermal necrolysis or erythema multiform).

Tell your doctor immediately if you experience:

- Faster heart rate, uneven or forceful heartbeat (palpitations), chest pain, tightness in your chest or more serious problems including heart attack and stroke
- Shortness of breath or a cough. These could be signs of lung problems
- Bruising more easily, bleeding for longer than normal, any sign of bleeding (e.g. bleeding from the gums), purple spots blotching on the skin or getting infections more easily than usual, sore throat and fever, feeling tired, faint, dizzy or having pale skin. These can be signs of blood or bone marrow problems
- Severe stomach pain which may reach through to your back. This could be a sign of pancreatitis (inflammation of the pancreas).
- Fever, chills, tiredness, loss of appetite, stomach pain, feeling sick, yellowing of your skin or eyes (jaundice). These can be signs of liver problems such as hepatitis (inflammation of the liver) or liver damage.

Other side effects include:

Tell your doctor if any of the following gets serious or lasts longer than a few days.

Common: (may affect up to 1 in 10 people)

- Headache or feeling tired
- Feeling dizzy. This is more likely to happen when you start taking Ramipril Teva Pharma or start taking a higher dose

- Fainting, hypotension (abnormally low blood pressure), especially when you stand or sit up quickly
- Dry tickly cough, inflammation of your sinuses (sinusitis) or bronchitis, shortness of breath
- Stomach or gut pain, diarrhoea, indigestion, feeling or being sick
- Skin rash with or without raised area
- Chest pain
- Cramps or pain in your muscles
- Blood tests showing more potassium than usual in your blood.

Uncommon: (may affect up to 1 in 100 people)

- Balance problems (vertigo)
- Itching and unusual skin sensations such as numbness, tingling, pricking, burning or creeping on your skin (paraesthesia)
- Loss or change in the way things taste
- Sleep problems
- Feeling depressed, anxious, more nervous than usual or restless
- Blocked nose, difficulty breathing or worsening of asthma
- A swelling in your gut called “intestinal angioedema” presenting with symptoms like abdominal pain, vomiting and diarrhoea
- Heartburn, constipation or dry mouth
- Passing more water (urine) than usual over the day
- Sweating more than usual
- Loss or decrease of appetite (anorexia)
- Increased or irregular heartbeats
- Swollen arms and legs. This may be a sign of your body holding onto more water than usual
- Flushing
- Blurred vision
- Pain in your joints
- Fever
- Sexual inability in men, reduced sexual desire in men or women
- An increased number of certain white blood cells (eosinophilia) found during a blood test
- Blood tests showing changes in the way your liver, pancreas or kidneys are working.

Rare: (may affect up to 1 in 1,000 people)

- Feeling shaky or confused
- Red and swollen tongue
- Severe flaking or peeling of the skin, itchy, lumpy rash
- Nail problem (e.g. loosening or separation of a nail from its bed)
- Skin rash or bruising
- Blotches on your skin and cold extremities
- Red, itchy, swollen or watery eyes
- Disturbed hearing and ringing in your ears
- Feeling weak
- Blood tests showing a decrease in the number of red blood cells, white blood cells or platelets or in the amount of haemoglobin.

Very rare: (may affect up to 1 in 10,000 people)

- Being more sensitive to the sun than usual.

Other side effects reported:

Tell your doctor if any of the following gets serious or lasts longer than a few days.

- Difficulty concentrating
- Swollen mouth
- Blood tests showing too few blood cells in your blood
- Blood tests showing less sodium than usual in your blood

- Concentrated urine (dark in color), feel or are sick, have muscle cramps, confusion and fits which may be due to inappropriate ADH (anti-diuretic hormone) secretion. If you have these symptoms contact your doctor as soon as possible.
- Fingers and toes changing colour when you are cold and then tingling or feeling painful when you warm up (Raynaud's phenomenon)
- Breast enlargement in men
- Slowed or impaired reactions
- Burning sensation
- Change in the way things smell
- Hair loss.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRRA Pharmacovigilance Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Ramipril Teva Pharma

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the outer carton and blister packs after EXP. The expiry date refers to the last day of that month.

Do not store above 25 °C.

Store in the original package in order to protect from moisture and light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Ramipril Teva Pharma contains

- The active substance is: ramipril.

1.25 mg

Each tablet contains 1.25 mg ramipril.

2.5 mg

Each tablet contains 2.5 mg ramipril.

5 mg

Each tablet contains 5 mg ramipril.

10 mg

Each tablet contains 10 mg ramipril.

- The other ingredients are:

1.25 mg

Sodium hydrogen carbonate, lactose monohydrate, croscarmellose sodium, pregelatinised starch (from maize starch), sodium stearyl fumarate, iron oxide yellow (E172).

2.5 mg

Sodium hydrogen carbonate, lactose monohydrate, croscarmellose sodium, pregelatinised starch (from maize starch), sodium stearyl fumarate, iron oxide yellow (E172).

5 mg

Sodium hydrogen carbonate, lactose monohydrate, croscarmellose sodium, pregelatinised starch (from maize starch), sodium stearyl fumarate, iron oxide yellow (E172), iron oxide red (E172).

10 mg

Sodium hydrogen carbonate, lactose monohydrate, croscarmellose sodium, pregelatinised starch (from maize starch), sodium stearyl fumarate.

What Ramipril Teva Pharma looks like and contents of the pack

Tablets.

1.25 mg

yellow, round, biconvex tablets without break score and a diameter of approximately 6.0 mm.

2.5 mg

yellow, round, biconvex tablets with break score on one side and a diameter of 8.0 mm.

5 mg

pink, round, biconvex tablets with break score on one side and a diameter of 6.5 mm.

10 mg

white to off-white, round, biconvex tablets with break score on one side and a diameter of 9.0 mm

Blister packs (OPA/Al/PVC – Al).

Packages with 7, 20, 21, 28, 30, 50, 56, 60, 90, 98, 100, 50x1 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Teva Pharma B.V.

Swensweg 5

2031GA Haarlem

The Netherlands

Manufacturers

Merckle GmbH

Ludwig-Merckle-Strasse 3,

89143 Blaubeuren

Germany

Teva Operations Poland Sp. z.o.o.

ul Mogilska 80

31-546 Kraków

Poland

This medicinal product is authorised in the Member States of the EEA under the following names:

Denmark	Ramipril Teva
Greece	Ramipril Teva 1.25 mg, 2.5 mg, 5 mg, 10 mg δισκία
Ireland	Ramipril Teva Pharma 1.25 mg, 2.5 mg, 5 mg, 10 mg Tablets
Netherlands	Ramipril Teva 1.25 mg, 2.5 mg, 5 mg, 10 mg tabletten
Norway	Ramipril Teva
Poland	Ramipril Teva Pharma
Portugal	Ramipril Teva
Romania	Ramipril Teva 5 mg, 10 mg comprimate
Spain	Ramipril Tevagen 1.25 mg, 2.5 mg, 5 mg, 10 mg comprimidos EFG
Sweden	Ramipril Teva
UK(NI)	Ramipril 1.25 mg, 2.5 mg, 5 mg, 10 mg Tablets

This leaflet was last revised in July 2023