

PACKAGE LEAFLET: INFORMATION FOR THE USER

FARMADURA™ XL 4mg

FARMADURA™ XL 8mg

Prolonged-release Tablets

doxazosin (as mesilate)

Important things that you **SHOULD** know about your medicine:

- Farmadura XL is a treatment for high blood pressure
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- You should take Farmadura XL regularly to get the maximum benefit, even if you are feeling well.
- Most people do not have serious problems when taking Farmadura XL but side effects can occur – see Section 4 for details. If you experience swelling of the face, tongue or windpipe, see your doctor immediately. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.
- Taking other medicines may sometimes cause problems. Check with your doctor or pharmacist before taking this or any other medicines.
- If you are, or are trying to become, pregnant, do not take Farmadura XL and tell your doctor.

Please read the rest of this leaflet. It includes other important information on the safe and effective use of this medicine that might be especially important to you. If you have any further questions, ask your doctor or pharmacist.

In this leaflet:

1. What Farmadura XL is and what it is used for
2. Before you take Farmadura XL
3. How to take Farmadura XL
4. Possible side effects
5. How to Store Farmadura XL
6. Further information

1. What Farmadura XL is and what it is used for

Farmadura XL is one of a group of medicines called alpha-blockers. It is used to treat high blood pressure.

Farmadura XL can be used to treat high blood pressure (hypertension), by relaxing blood vessels so that blood passes through them more easily. This helps to lower blood pressure.

2. Before You Take Farmadura XL

⚠ Do not take Farmadura XL:

- If you have had an allergic reaction in the past to Farmadura XL, the active ingredient doxazosin, other types of quinazolines (such as prazosin or terazosin) or any of the other

ingredients listed in section 6. This may have been itching, reddening of the skin or difficulty in breathing.

- If you are breast-feeding
- If you have any form of obstruction of the digestive tract
- If you are under 12 years of age
- If you have a history of a condition known as 'orthostatic hypotension' which is a form of low blood pressure that causes you to feel dizzy or light-headed when you stand up from sitting or lying down
- If you have an enlarged prostate gland with one of the following: any kind of congestion or blockage in your urinary tract, a longstanding infection of the urinary tract or you have bladder stones.
- If you have an enlarged prostate gland and have overflow incontinence (you do not feel the urge to urinate), or anuria (your body is not producing any urine) with or without kidney problems.

Take special care with Farmadura XL:

The following are reasons why Farmadura XL may not be suitable for you:

- If you are pregnant or trying to become pregnant
- If you have liver, kidney or heart disease
- If you are also taking other medicines
- If you have problems with your heart

If you are undergoing eye surgery because of cataract (cloudiness of the lens) please inform your eye specialist before the operation that you are using or have previously used Farmadura XL. This is because this medicine may cause complications during the surgery which can be managed if your specialist is prepared in advance.

When you start to take Farmadura XL you may experience faintness or dizziness caused by low blood pressure, when getting up from sitting or lying down. If you feel faint or dizzy, you should sit or lie down until you feel better and avoid situations where you might fall or hurt yourself.

Taking other medicines:

There are some medicines that may interact with Farmadura XL

- Medicines for erectile dysfunction (impotence) such as sildenafil, tadalafil or vardenafil
Some patients who take alpha-blocker therapy for the treatment of high blood pressure or prostate enlargement may experience dizziness or light-headedness, which may be caused by low blood pressure upon sitting or standing up quickly. Certain patients have experienced these symptoms when taking drugs for erectile dysfunction (impotence) with alpha-blockers. In order to reduce the likelihood that these symptoms occur, you should be on a regular daily dose of your alpha blocker before you start drugs for erectile dysfunction.
- Farmadura XL may also increase the effects of drugs used to treat high blood pressure (hypertension)

You should always tell your doctor if you are taking or have recently taken any other medicine, even those not prescribed, because they might interact with Farmadura XL.

☒ Taking Farmadura XL with food and drink:

Farmadura XL can be taken with or without food.

☒ Pregnancy and Breast-feeding:

The safety of using Farmadura XL during pregnancy has not been established. If you are pregnant or trying to become pregnant speak to your doctor who will decide if this medicine is suitable for you. Do not take Farmadura XL if you are breast-feeding.

☒ Driving and using machinery:

Take care if you drive or operate machinery. Your tablets may affect your ability to drive or operate machinery safely, particularly when you first start to take them. They may cause drowsiness and could also make you feel weak or dizzy. If you are affected, do not drive or operate machinery and contact your doctor immediately.

3. How To Take Farmadura XL

Farmadura XL is a prolonged-release tablet. The medicine is contained within a non-absorbable shell that has been specially designed to slowly release the medicine. Once swallowed, the medicine (doxazosin) is slowly released into the body from the tablet, until the tablet is empty. Since the empty tablet is eliminated from the body in bowel movements, you may occasionally observe in the stools, something that looks like a tablet. This is to be expected and you should not be concerned.

How to take your medicine

- The usual dose of Farmadura XL is one 4mg prolonged-release tablet taken once a day. Your doctor may wish to increase your dose to 8mg. This is the maximum dose of this medicine.
- Swallow your medicine whole with water. **Do not chew, divide or crush** the tablets.
- Take your tablets as your doctor told you.
- It is important to keep taking your tablets. They help to control your blood pressure.
- Don't change the dose or stop taking the tablets without first checking with your doctor.
- Don't wait until your tablets are finished before seeing your doctor.
- If you are still not sure how to take your tablets, ask your doctor or pharmacist.

If you take more Farmadura XL than you should:

Taking too many tablets at once may make you unwell. If several tablets are taken it may be dangerous. Tell your doctor immediately or go to your nearest hospital casualty department immediately.

If you miss a dose of Farmadura XL:

Do not worry. If you forget to take a tablet, leave that dose out completely. Then go on as before.

4. Possible side effects

Like all medicines, Farmadura XL can cause side effects although not everybody gets them.

STOP taking Farmadura XL and call an ambulance immediately if you experience any of the following:

- Heart attack
- Increased, decreased or irregular heartbeat
- Weakness of arms, legs or problems speaking which may be symptoms of a stroke

- Allergic reactions (hypersensitivity). Symptoms include sudden wheeziness, tightness in chest, difficulty in breathing, swelling of the face, tongue or throat, rash or itching (especially affecting the whole body)

Tell your doctor **immediately** if you experience any of the following symptoms after taking Farmadura XL:

- Chest pain (angina), wheezing, shortness of breath, difficulty breathing
- Feeling your heartbeat (palpitations)
- Fainting
- Yellowing of the skin or the eyes (jaundice)
- Low numbers of white blood cells or blood platelets, which may result in bruising or easy bleeding

The following events have been reported in patients being treated with Farmadura XL. If any of these side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor.

Common side effects

These can affect between 1 in 10 and 1 in 100 patients:

- dizziness, feeling of spinning or rotation of surroundings (vertigo), headache
- low blood pressure
- swelling of feet, ankles or fingers
- bronchitis, coughing, respiratory tract (nose, throat, lungs) infection
- nasal stuffiness, sneezing and/or runny nose caused by inflammation of the lining of the nose (rhinitis)
- stomach/abdominal pains, feeling/being sick
- urinary tract infection, cystitis, urinary incontinence (inability to control passing urine)
- sleepiness, general weakness
- indigestion, dry mouth
- itching
- back pain, painful muscles
- flu-like symptoms

Uncommon side effects:

These can affect between 1 in 100 and 1 in 1,000 patients:

- constipation, wind, inflammation of the stomach and intestines (gastroenteritis) which can cause diarrhoea and vomiting
- pain or discomfort on passing urine, increased frequency in passing urine, blood in urine
- inflammation of the joints (gout), painful joints, general pain
- sleeplessness, anxiety, depression
- reduced or altered sense of touch or sensation of the hands and feet
- increased appetite or loss of appetite, weight gain
- nose bleeds
- skin rash
- ringing or noise in the ears, tremor

- failure/ inability to achieve penile erection
- liver enzyme increases which may have an effect on some medical tests

Very Rare side effects

These can affect less than 1 in 10,000 patients:

- faintness or dizziness caused by low blood pressure when getting up from a sitting or lying position
- agitation or nervousness
- hepatitis (liver inflammation) or bile disorder
- hives, hair loss, red or purple patches on the skin, bleeding under the skin
- tingling or numbness of the hands and feet
- tiredness, generally feeling unwell
- blurred vision
- hot flushes
- muscle cramps, muscle weakness
- disorder in passing urine, needing to pass urine at night, increased number of times of passing urine, increased volume of urine passed
- discomfort or enlargement of the breasts in men
- painful persistent erection of the penis

Other side effects

The frequency of these side effects is unknown:

- little or no semen ejaculated at sexual climax, cloudy urine following sexual climax
- eye problems may occur during eye surgery for cataract (cloudiness of the lens of the eye). See section “Take special care with Farmadura XL”.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor.

5. How To Store Farmadura XL

Keep your tablets out of the reach and sight of children.

Do not use Farmadura XL after the expiry date which is stated on the carton and tablet blister after Exp. The expiry date refers to the last day of that month.

Do not store above 30°C.

Keep your tablets in the original packaging. This medicine is sensitive to moisture.

The tablets should be whole to work effectively, if you notice any defects (such as broken tablets) consult your pharmacist immediately.

6. Further Information

What Farmadura XL contains and its package

- The active substance is doxazosin (as doxazosin mesilate)
- The other ingredients are polyethylene oxide, sodium chloride, hypromellose, red ferric oxide (E172), titanium dioxide (E171), magnesium stearate, cellulose acetate, Macrogol, pharmaceutical glaze, black iron oxide (E172), ammonium hydroxide and propylene glycol

Farmadura XL tablets come in two strengths, 4mg and 8mg.

Each tablet contains doxazosin mesilate equivalent to 4mg or 8mg of doxazosin, within a shell that has been specially designed to slowly release the drug.

The 4mg tablets are white, round biconvex shaped with a hole in one side, marked CXL4.

The 8mg tablets are white, round biconvex shaped with a hole in one side, marked CXL8.

Farmadura XL tablets are available in packs of 28 tablets.

Marketing Authorisation Holder

Pfizer Healthcare Ireland
9 Riverwalk
National Digital Park
Citywest Business Campus
Dublin 24
Republic of Ireland

Manufacturer

Pfizer Manufacturing
Deutschland GmbH,
Heinrich-Mack-Strasse 35,
89257 Illertissen,
Germany

Company contact address:

For further information on your medicine contact Medical Information at the following address:
Pfizer Healthcare Ireland, 9 Riverwalk, National Digital Park, Citywest Business Campus,
Dublin 24, Ireland
Telephone 1800 633 363

This leaflet was last revised in 04/2013

Ref: FD2_1 IE