

PACKAGE LEAFLET: INFORMATION FOR THE USER

Meropenem Hospira 500 mg and 1 g Powder for Solution for Injection or Infusion
meropenem

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to you doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Meropenem Hospira is and what it is used for
2. What you need to know before you use Meropenem Hospira
3. How to use Meropenem Hospira
4. Possible side effects
5. How to store Meropenem Hospira
6. Contents of the pack and other information

1. WHAT MEROPENEM HOSPIRA IS AND WHAT IT IS USED FOR

Meropenem Hospira belongs to a group of medicines called carbapenem antibiotics. It works by killing bacteria, which can cause serious infections.

- Infection affecting the lungs (pneumonia)
- Lung and bronchial infections in patients suffering from cystic fibrosis
- Complicated urinary tract infections
- Complicated infections in the abdomen
- Infections that you can catch during or after delivery
- Complicated skin and soft tissue infections
- Acute bacterial infection of the brain (meningitis)

Meropenem Hospira may be used in the management of neutropenic patients with fever that is suspected to be due to bacterial infection.

Meropenem may be used to treat bacterial infection of the blood which might be associated with a type of infection mentioned above.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE MEROPENEM HOSPIRA

Do not use Meropenem Hospira

- if you are allergic to meropenem or any of the other ingredients of Meropenem Hospira (listed in Section 6)
- if you are allergic (hypersensitive) to other antibiotics such as penicillins, cephalosporins, or carbapenems as you may also be allergic to meropenem

Warnings and precautions

Talk to your doctor or nurse before using Meropenem Hospira:

- if you have health problems, such as liver or kidney problems.
- if you have had severe diarrhoea after taking other antibiotics.

You may develop a positive test (Coombs test) which indicates the presence of antibodies that may destroy red blood cells. Your doctor will discuss this with you.

If you are not sure if any of the above applies to you, talk to your doctor or nurse before using Meropenem Hospira.

Other medicines and Meropenem

Tell your doctor if you are taking, have recently taken or might take any other medicines. This is because Meropenem Hospira can affect the way some medicines work and some medicines can have an effect on Meropenem Hospira.

In particular, tell your doctor or nurse if you are taking any of the following medicines:

- Probenecid (used to treat gout).
- Valproic acid/sodium valproate/valpromide (used to treat epilepsy).
Meropenem Hospira should not be used because it may decrease the effect of sodium valproate.
- Oral anti-coagulant agent (used to treat or prevent blood clots).

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. It is preferable to avoid the use of meropenem during pregnancy.

Your doctor will decide whether you should use Meropenem Hospira.

It is important that you tell your doctor if you are breast-feeding or if you intend to breast-feed before receiving Meropenem Hospira. Small amounts of this medicine may pass into the breast milk and it may affect the baby. Therefore, your doctor will decide whether you should use Meropenem Hospira while breast-feeding.

Ask your doctor for advice before taking any medicine.

Driving and using machines

No studies on the effect on the ability to drive and use machines have been performed.

Meropenem Hospira contains sodium.

Meropenem Hospira 500 mg: This medicinal product contains approximately 2.0 mEq of sodium per 500 mg dose which should be taken into consideration by patients on a controlled sodium diet.

Meropenem Hospira 1 g: This medicinal product contains approximately 4.0 mEq of sodium per 1 g dose which should be taken into consideration by patients on a controlled sodium diet.

If you have a condition which requires you to monitor your sodium intake please inform your doctor or nurse.

3. HOW TO USE MEROPENEM HOSPIRA

Always use this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

Use in adults

- The dose depends on the type of infection that you have, where the infection is in the body and how serious the infection is. Your doctor will decide on the dose that you need.
- The dose for adults is usually between 500 mg (milligrams) and 2 g (gram). You will usually receive a dose every 8 hours. However you may receive a dose less often if your kidneys do not work very well.

Use in children and adolescents

- The dose for children over 3 months old and up to 12 years of age is decided using the age and weight of the child. The usual dose is between 10 mg and 40 mg of Meropenem Hospira for each kilogram (kg) that the child weighs. A dose is usually given every 8 hours. Children who weigh over 50 kg will be given an adult dose.

Method of administration

- Meropenem Hospira will be given to you as an injection or infusion into a large vein.
- Your doctor or nurse will normally give Meropenem Hospira to you.
- However, some patients, parents and carers are trained to give Meropenem Hospira at home. Instructions for doing this are provided in this leaflet (in the section called 'Instructions for giving Meropenem Hospira to yourself or someone else at home'). Always use Meropenem Hospira exactly as your doctor has told you. You should check with your doctor if you are not sure.
- Your injection should not be mixed with or added to solutions that contain other medicines.
- The injection may take about 5 minutes or between 15 and 30 minutes. Your doctor will tell you how to give Meropenem Hospira.
- You should normally have your injections at the same times each day.

If you use more Meropenem Hospira than you should

If you accidentally use more than your prescribed dose, contact your doctor or nearest hospital straight away.

If you forget to use Meropenem Hospira

If you miss an injection, you should have it as soon as possible. However, if it is almost time for your next injection, skip the missed injection. Do not take a double dose (two injections at the same time) to make up for a forgotten dose.

If you stop using Meropenem Hospira

Do not stop having Meropenem Hospira until your doctor tells you to.

If you have any further questions on the use of this product, ask your doctor or nurse.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The frequency of possible side effects listed below;

Severe allergic reactions

If you have a severe allergic reaction, **stop having Meropenem Hospira and see a doctor straight away**. You may need urgent medical treatment. The signs may include a sudden onset of:

- Severe rash, itching or hives on the skin.
- Swelling of the face, lips, tongue or other parts of the body.
- Shortness of breath, wheezing or trouble breathing.

Severe reaction (rare)

You may develop DRESS (Drug Reaction with Eosinophilia and Systemic Symptoms). If you have any of the following signs, **stop having Meropenem Hospira and see a doctor straight away**. You may need urgent medical treatment.

The signs may include:

- Flu-like symptoms
- Rash on the face and then an extended rash with a high temperature.
- Increased level of liver enzymes, seen in blood tests.
- Increase in a type of white blood cell (eosinophilia), seen in blood tests.
- Enlarged lymph nodes.

Damage to red blood cells (not known)

The signs include:

- Being breathless when you do not expect it.
- Red or brown urine.

If you notice any of the above, **see a doctor straight away**.

Other possible side effects:

Common (may affect up to 1 in 10 people)

- Abdominal (stomach) pain.
- Feeling sick (nausea).
- Being sick (vomiting).
- Diarrhoea.
- Headache.
- Skin rash, itchy skin.
- Pain and inflammation.

- Increased number of platelets in your blood (shown in a blood test).
- Changes in blood tests, including tests that show how well your liver is working.

Uncommon (may affect up to 1 in 100 people)

- Changes in your blood. These include reduced numbers of platelets (which may make you bruise more easily), increased numbers of some white blood cells, decreased numbers of other white cells and increased amounts of a substance called ‘bilirubin’. Your doctor may do blood tests from time to time.
- Changes in blood tests, including tests that show how well your kidneys are working.
- A tingling feeling (pins and needles).
- Infections of the mouth or the vagina that are caused by a fungus (thrush).
- Inflammation of the bowel with diarrhoea.
- Sore veins where Meropenem Hospira is injected.
- Other changes in your blood. The symptoms include frequent infections, high temperature and sore throat. Your doctor may do blood tests from time to time.
- Sudden onset of a severe rash or blistering or peeling skin. This may be associated with a high fever and joint pains.

Rare (may affect up to 1 in 1000 people)

- Fits (convulsions).
-

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

HPRA Pharmacovigilance
 Earlsfort Terrace
 IRL - Dublin 2
 Tel: +353 1 6764971
 Fax: +353 1 6762517
 Website: www.hpra.ie
 e-mail: medsafety@hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE MEROPENEM HOSPIRA

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the vial label and carton after ‘EXP’. The expiry date refers to the last day of that month.

Prior to reconstitution or dilution, this medicinal product does not require any special storage conditions.

After reconstitution: The reconstituted solutions for intravenous injection or infusion should be used immediately. The product should be inspected visually for particulate matter, damage to the container or discolouration (solution should be colourless to pale yellow) prior to administration. Discard the product if such defects are observed.

The time interval between the beginning of reconstitution and the end of intravenous injection or infusion should not exceed one hour.

Do not freeze the reconstituted solution.

Do not throw away any medicine via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Meropenem Hospira contains

Each 0.5 g vial contains meropenem trihydrate equivalent to 500 mg anhydrous meropenem.

Each 1 g vial contains meropenem trihydrate equivalent to 1 g anhydrous meropenem.

The other ingredient is sodium carbonate, anhydrous.

What Meropenem Hospira looks like and contents of the pack

- Meropenem Hospira is a white to pale yellow crystalline powder for solution for injection or infusion in vials. Pack sizes of 1 or 10 vials.

Marketing Authorisation Holder and Manufacturer

The Marketing Authorisation Holder and manufacturer (responsible for batch release of the product in the EEA) is Hospira UK Limited, Horizon, Honey Lane, Hurley, Maidenhead, SL6 6RJ UK.

This medicinal product is authorised in the Member States of the EEA under the following names:

Austria: Meropenem Pfizer
Belgium: Meropenem Hospira
Bulgaria: Meropenem Hospira
Denmark: Meropenem Hospira
Germany: Meropenem Hospira
Estonia: Meropenem Hospira
Ireland: Meropenem Hospira
Luxembourg: Meropenem Hospira
Netherlands: Meropenem Hospira
Norway: Meropenem Hospira
Poland: Meropenem Hospira
Romania: Meropenem Hospira
Sweden: Meropenem Hospira

This leaflet was last revised in 12/ 2016

Advice/medical education

Antibiotics are used to treat infections caused by bacteria. They have no effect against infections caused by viruses.

Sometimes an infection caused by bacteria does not respond to a course of an antibiotic. One of the commonest reasons for this to occur is because the bacteria causing the infection are resistant to the antibiotic that is being taken. This means that they can survive and even multiply despite the antibiotic.

Bacteria can become resistant to antibiotics for many reasons. Using antibiotics carefully can help to reduce the chance of bacteria becoming resistant to them.

When your doctor prescribes a course of an antibiotic it is intended to treat only your current illness. Paying attention to the following advice will help prevent the emergence of resistant bacteria that could stop the antibiotic working.

1. It is very important that you take the antibiotic at the right dose, at the right times and for the right number of days. Read the instructions on the label and if you do not understand anything ask your doctor or pharmacist to explain.
2. You should not take an antibiotic unless it has been prescribed specifically for you and you should use it only to treat the infection for which it was prescribed.
3. You should not take antibiotics that have been prescribed for other people even if they had an infection that was similar to yours.
4. You should not give antibiotics that were prescribed for you to other people.
5. If you have any antibiotic left over when you have taken the course as directed by your doctor you should take the remainder to a pharmacy for appropriate disposal.

The following information is intended for medical or healthcare professionals only:

Instructions for giving Meropenem Hospira to yourself or someone else at home

Some patients, parents and carers are trained to give Meropenem Hospira at home.

Warning - You should only give this medicine to yourself or someone else at home after a doctor or nurse has trained you.

- The medicine must be mixed with another liquid (the diluent). Your doctor will tell you how much of the diluent to use.
- Use the medicine straight after preparing it. Do not freeze it.

How to prepare this medicine

1. Wash your hands and dry them very well. Prepare a clean working area.
2. Remove the Meropenem Hospira vial from the packaging. Check the vial and the expiry date. Check that the vial is intact and has not been damaged.
3. Remove the cap and clean the rubber stopper with an alcohol wipe. Allow the rubber stopper to dry.
4. Connect a new sterile needle to a new sterile syringe, without touching the ends.
5. Draw up the recommended amount of sterile 'Water for Injections' into the syringe. The amount of liquid that you need is shown in the table below:

Dose of Meropenem Hospira	Amount of 'Water for Injections' needed for dilution
500 mg (milligrams)	10 ml (millilitres)
1 g (gram)	20 ml
1.5 g	30 ml
2 g	40 ml

Please note: If your prescribed dose of Meropenem Hospira is more than 1 g, you will need to use more than 1 vial of Meropenem Hospira. You can then draw the liquid in the vials into the one syringe.

6. Put the needle of the syringe through the centre of the rubber stopper and inject the recommended amount of Water for Injections into the vial or vials of Meropenem Hospira.
7. Remove the needle from the vial and shake the vial well for about 5 seconds, or until all the powder has dissolved. Clean the rubber stopper once more with a new alcohol wipe and allow the rubber stopper to dry.
8. With the plunger of the syringe pushed fully into the syringe, put the needle back through the rubber stopper. You must then hold both the syringe and the vial and turn the vial upside down.
9. Keeping the end of the needle in the liquid, pull back the plunger and draw all the liquid in the vial into the syringe.
10. Remove the needle and syringe from the vial and throw the empty vial away in a safe place.

11. Hold the syringe upright, with the needle pointing upwards. Tap the syringe so that any bubbles in the liquid rise to the top of the syringe.
12. Remove any air in the syringe by gently pushing the plunger until all the air has gone.
13. If you are using Meropenem Hospira at home, dispose of any needles and infusion lines that you have used in an appropriate way. If your doctor decides to stop treatment, dispose of any unused Meropenem Hospira in an appropriate way.

Giving the injection

You can either give this medicine through a short cannula or venflon, or through a port or central line.

Giving Meropenem Hospira through a short cannula or venflon

1. Remove the needle from the syringe and throw the needle away carefully in your sharps bin.
2. Wipe the end of the short cannula or venflon with an alcohol wipe and allow it to dry. Open the cap on your cannula and connect the syringe.
3. Slowly push the plunger of the syringe to give the antibiotic steadily over about 5 minutes.
4. Once you have finished giving the antibiotic and the syringe is empty, remove the syringe and use a flush as recommended by your doctor or nurse.
5. Close the cap of your cannula and carefully throw the syringe away in your sharps bin.

Giving Meropenem Hospira through a port or central line

1. Remove the cap on the port or line, clean the end of the line with an alcohol wipe and allow it to dry.
2. Connect the syringe and slowly push the plunger on the syringe to give the antibiotic steadily over about 5 minutes.
3. Once you have finished giving the antibiotic, remove the syringe and use a flush as recommended by your doctor or nurse.
4. Place a new clean cap on your central line and carefully throw the syringe away in your sharps bin.

