

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### **LEMSIP MAX COLD & FLU Blackcurrant 1000mg Powder for Oral solution Paracetamol**

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

This medicine is available without prescription. However, you still need to take Lemsip Max Cold & Flu Blackcurrant carefully to get the best results from it.

#### **1. WHAT LEMSIP MAX COLD & FLU BLACKCURRANT IS AND WHAT IT IS USED FOR?**

Each sachet of Lemsip Max Cold & Flu Blackcurrant 1000mg Powder for Oral Solution contains 1000mg paracetamol. Paracetamol has pain killing (analgesic) and fever reduction (antipyretic) properties.

Lemsip Max Cold & Flu Blackcurrant can be used to relieve the symptoms of common cold or influenza such as headache, sore throat, fever, body aches and pains.

#### **2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE LEMSIP MAX COLD & FLU BLACKCURRANT**

##### **Do not take Lemsip Max Cold & Flu Blackcurrant**

- If you are allergic (hypersensitive) to paracetamol or any of the other ingredients listed in section 6
- If you are taking any other paracetamol containing products
- If you are less than 16 years of age

##### **Take special care with Lemsip Max Cold & Flu Blackcurrant**

- If you have impaired liver or kidney function
- If you are underweight (less than 50 kg) or are malnourished
- If you use large quantities of alcohol regularly (alcoholism)
- If you suffer from severe dehydration
- If you are taking other drugs that affect the liver (see Taking other medicines)
- If you are elderly
- If you have glutathione deficiency (a metabolic disorder which may lead to difficulty in breakdown of paracetamol)
- If you have an enzyme deficiency called Glucose-6-phosphate dehydrogenase deficiency
- If you have haemolytic anaemia (abnormal breakdown of red blood cells)

If any of the above applies to you, you may need to reduce the amount of paracetamol that you take or avoid taking this medicinal product altogether.

During treatment with Lemsip Max Cold & Flu Blackcurrant, tell your doctor straight away if:

- If you have severe illnesses, including severe renal impairment or sepsis (when bacteria and their toxins circulate in the blood leading to organ damage), or you suffer from malnutrition, chronic alcoholism or if you are also taking flucloxacillin (an antibiotic). A serious condition called metabolic acidosis (a blood and fluid abnormality) has been reported in patients in these situations when paracetamol is used at regular doses for a prolonged period or when paracetamol is taken together with flucloxacillin. Symptoms of metabolic acidosis may include: serious breathing difficulties with deep rapid breathing, drowsiness, feeling sick (nausea) and being sick (vomiting).

If you are diabetic, paracetamol contained in this medicine may falsely elevate continuous blood glucose monitor readings compared to finger prick readings. This is not the case with all continuous blood glucose monitors. Please refer to the information provided by the manufacturer of your monitor. Contact your doctor, pharmacist or nurse if you are unsure or should you require further advice.

Prolonged use, except under medical supervision can be harmful. If you have a high fever, or if symptoms get worse or persist for longer than 3 days, consult your doctor. Only take when necessary. Do not exceed the stated dose.

##### **Taking other medicines**

Care should be taken while taking this product if you are currently taking:

- Drugs to stop you feeling sick such as metoclopramide or domperidone: may increase absorption of paracetamol.
- Cholesterol drugs such as cholestyramine: may reduce absorption of paracetamol
- Anticoagulants such as warfarin: their effect may be enhanced by long term use of paracetamol.
- Drugs that affect the liver such as monoamine oxidase inhibitors and antidepressants: may increase the hepatotoxicity (liver toxicity) of paracetamol.

- Flucloxacillin. Please inform your doctor or pharmacist if you are taking flucloxacillin (antibiotic), due to a serious risk of blood and fluid abnormality (called metabolic acidosis) that must have urgent treatment.

### **Taking Lemsip Max Cold & Flu Blackcurrant with food, drink and alcohol**

Alcohol may increase the adverse effects of paracetamol.

### **Pregnancy, breast-feeding and fertility:**

Lemsip Max Cold & Flu Blackcurrant can be used during pregnancy. You should use the lowest possible dose that reduces your pain and/or fever and use it for the shortest time possible. Contact your doctor if the pain and/or fever are not reduced or if you need to take the medicine more often.

Paracetamol is excreted in breast milk, but not in a clinically significant amount. You can take this product whilst breastfeeding.

### **Important information about some of the ingredients of Lemsip Max Cold & Flu Blackcurrant**

- This medicinal product contains 61.5 mg aspartame in each sachet. Aspartame is a source of phenylalanine. It may be harmful if you have phenylketonuria (PKU), a rare genetic disorder in which phenylalanine builds up because the body cannot remove it properly.
- This medicinal product contains 2.11 g of sucrose per sachet. This should be taken into account in patients with diabetes mellitus. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

This medicinal product contains 128.71 mg sodium (main component of cooking/table salt) in each sachet. This is equivalent to 6.4 % of the recommended maximum daily dietary intake of sodium for an adult. This product is considered high in salt and should be taken in to account if on a low salt diet

### **3. HOW TO TAKE LEMSIP MAX COLD & FLU BLACKCURRANT**

Always use the lowest effective dose for the shortest time necessary to relieve your symptoms. Do not exceed the stated dose. For oral administration. Dilute the contents of one sachet in a mug of hot water, do not boil the water. Stir until dissolved. If preferred, sweeten to taste with sugar, honey or usual sweetener.

Adults, the elderly and children aged 16 years and over: The usual dose is 1 sachet up to 4 times daily. The dose may be repeated in 4 hours with a maximum of four doses in a 24 hour period.

Do not give to children under 16 years of age.

For children under 16 years, other formulations and dosage strengths are available which may be more appropriate.

Do not take more than 2 sachets a day (2 g/day) in the following situations, unless directed by your doctor:

- Weight less than 50 kg
- Use of large quantities of alcohol regularly (alcoholism)
- Dehydration
- Malnutrition

Use in kidney or liver impairment:

If you have been diagnosed with kidney or liver problems, please consult your doctor or pharmacist. You may need to take a lower dose or increase the length of time between doses.

### **If you take more Lemsip Max Cold & Flu Blackcurrant than you should:**

Immediate medical advice should be sought in the event of overdose because of the risk of irreversible liver damage. There is an increased risk of liver damage from paracetamol overdosing associated patients who consume alcohol in excess of recommended amounts, patients on long-term treatment with enzyme-inducing drugs and patients likely to be lacking glutathione e.g. eating disorders, cystic fibrosis, HIV infection, starvation, cachexia. Symptoms of overdose may present as pallor, nausea, vomiting, loss of appetite and abdominal pain.

### **If you forget to take Lemsip Max Cold & Flu Blackcurrant**

Take the next dose as soon as you remember, if needed. Do not double dose. This product should only be taken when necessary. If you have any further questions on the use of this product, ask your pharmacist.

### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, Lemsip Max Cold & Flu Blackcurrant can cause side effects although not everybody gets them. Side effects may include hypersensitivity reactions such as skin rashes, skin and tissue disorders such as shedding of your skin, blisters on your skin and mucous membranes, facial swelling, tongue swelling. If skin rash occurs discontinue use immediately. There have also been reports of blood disorders.

Frequency “Very rare”:

- Very rare cases of serious skin reactions have been reported.

Frequency “Not known” (frequency cannot be estimated from the available data):

- A serious condition that can make blood more acidic (called metabolic acidosis), in patients with severe illness using paracetamol (see section 2).

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRa Pharmacovigilance, Website: [www.hpra.ie](http://www.hpra.ie)  
By reporting side effects you can help provide more information on the safety of this medicine.

## **5. HOW TO STORE LEMSIP MAX COLD & FLU BLACKCURRANT**

Keep out of the sight and reach of children.

Do not use Lemsip Max Cold & Flu Blackcurrant after the expiry date (EXP Day/Month/Year) which is stated on the carton and sachet. Do not store above 25°C. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **6. CONTENTS OF THE PACK AND OTHER INFORMATION**

### **What Lemsip Max Cold & Flu Blackcurrant contains**

Each sachet contains 1000mg of paracetamol as the active ingredient. The other ingredients are sucrose, citric acid anhydrous granular, blackcurrant witham flavour, saccharin sodium, aspartame (E951), sodium citrate, ascorbic acid and enocyanin.

### **What Lemsip Max Cold & Flu Blackcurrant looks like and contents of the pack**

Lilac coloured homogenous powder for oral solution with a taste and odour of blackcurrant. Each carton contains 5 or 10 sachets, each containing 4.8g of product. Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

MA Holder: Reckitt Benckiser Ireland Ltd, 7 Riverwalk, Citywest Business Campus, Dublin 24, Ireland.

Manufacturers: Reckitt Benckiser Healthcare (UK) Ltd., Hull, HU8 7DS, UK and RB NL Brands B.V. Schiphol Blvd 207, 1118 BH Schiphol, NL.

For product queries please call: 01 6305429

Date of revision of text: upon approval