

## **Package leaflet: Information for the patient**

**Atorvastatin Billev Pharma 10 mg film-coated tablets**  
**Atorvastatin Billev Pharma 20 mg film-coated tablets**  
**Atorvastatin Billev Pharma 40 mg film-coated tablets**

Atorvastatin

### **Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Atorvastatin Billev Pharma is and what it is used for
2. What you need to know before you take Atorvastatin Billev Pharma
3. How to take Atorvastatin Billev Pharma
4. Possible side effects
5. How to store Atorvastatin Billev Pharma
6. Contents of the pack and other information

### **1. What Atorvastatin Billev Pharma is and what it is used for**

Atorvastatin Billev Pharma belongs to a group of medicines known as statins, which are lipid (fat) regulating medicines.

Atorvastatin Billev Pharma is used to lower lipids known as cholesterol and triglycerides in the blood when a low fat diet and life style changes on their own have failed. If you are at an increased risk of heart disease, Atorvastatin Billev Pharma can also be used to reduce such risk even if your cholesterol levels are normal. You should maintain a standard cholesterol lowering diet during treatment.

### **2. What you need to know before you take Atorvastatin Billev Pharma**

#### **Do not take Atorvastatin Billev Pharma**

- if you are allergic to atorvastatin or to any similar medicines used to lower blood lipids or to any of the other ingredients of the medicine (listed in section 6).
- if you have or have ever had a disease which affects the liver
- if you have had any unexplained abnormal blood tests for liver function
- if you are a woman able to have children and not using reliable contraception
- if you are pregnant or trying to become pregnant
- if you are breast-feeding.

#### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Atorvastatin Billev Pharma.

The following are reasons why Atorvastatin Billev Pharma may not be suitable for you:

- if you have had a previous stroke with bleeding into the brain, or have small pockets of fluid in the brain from previous strokes
- if you have kidney problems
- if you have an under-active thyroid gland (hypothyroidism)
- if you have had repeated or unexplained muscle aches or pains, a personal history or family history of muscle problems
- if you have had previous muscular problems during treatment with other lipid-lowering medicines (e.g. other ‘-statin’ or ‘-fibrate’ medicines)
- if you regularly drink a large amount of alcohol
- if you have a history of liver disease
- if you are older than 70 years.

**Check with your doctor or pharmacist before taking Atorvastatin Billev Pharma**

- if you have severe respiratory failure.

If any of these apply to you, your doctor will need to carry out a blood test before and possibly during your Atorvastatin Billev Pharma treatment to predict your risk of muscle related side effects. The risk of muscle related side effects e.g. rhabdomyolysis is known to increase when certain medicines are taken at the same time (see Section 2 “Other medicines and Atorvastatin Billev Pharma”).

While you are on this medicine your doctor will monitor you closely if you have diabetes or are at risk of developing diabetes. You are likely to be at risk of developing diabetes if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure.

**Other medicines and Atorvastatin Billev Pharma**

There are some medicines that may change the effect of Atorvastatin Billev Pharma or their effect may be changed by Atorvastatin Billev Pharma. This type of interaction could make one or both of the medicines less effective. Alternatively it could increase the risk or severity of side-effects, including the important muscle wasting condition known as rhabdomyolysis described in Section 4:

- Medicines used to alter the way your immune system works, e.g. ciclosporin
- Certain antibiotics or antifungal medicines, e.g. erythromycin, clarithromycin, telithromycin, ketoconazole, itraconazole, voriconazole, fluconazole, posaconazole, rifampin, fusidic acid
- Other medicines to regulate lipid levels, e.g. gemfibrozil, other fibrates, colestipol
- Some calcium channel blockers used for angina or high blood pressure, e.g. amlodipine, diltiazem
- Medicines to regulate your heart rhythm e.g. digoxin, verapamil, amiodarone
- Medicines used in the treatment of HIV e.g. ritonavir, lopinavir, atazanavir, indinavir, darunavir, the combination of tipranavir/ritonavir etc.
- Some medicines used in the treatment of hepatitis C e.g. telaprevir
- Other medicines known to interact with Atorvastatin Billev Pharma include ezetimibe (which lowers cholesterol), warfarin (which reduces blood clotting), oral contraceptives, stiripentol (an anti-convulsant for epilepsy), cimetidine (used for heartburn and peptic ulcers), phenazone (a painkiller), colchicine (used to treat gout), and antacids (indigestion products containing aluminium or magnesium) and boceprevir (used to treat liver disease such as hepatitis C)
- Medicines obtained without a prescription: St John’s Wort

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

**Atorvastatin Billev Pharma with food and drink**

See Section 3 for instructions on how to take Atorvastatin Billev Pharma. Please note the following:

*Grapefruit juice*

Do not take more than one or two small glasses of grapefruit juice per day because large quantities of grapefruit juice can change the effects of Atorvastatin Billev Pharma.

#### *Alcohol*

Avoid drinking too much alcohol while taking this medicine. See Section 2 “Warnings and precautions” for details

#### **Pregnancy and breast-feeding**

Do not take Atorvastatin Billev Pharma if you are pregnant, or if you are trying to become pregnant.

Do not take Atorvastatin Billev Pharma if you are able to become pregnant unless you use reliable contraceptive measures.

Do not take Atorvastatin Billev Pharma if you are breast-feeding.

The safety of Atorvastatin Billev Pharma during pregnancy and breast-feeding has not yet been proven. Ask your doctor or pharmacist for advice before taking any medicine.

#### **Driving and using machines**

Normally this medicine does not affect your ability to drive or operate machines. However, do not drive if this medicine affects your ability to drive. Do not use any tools or machines if your ability to use them is affected by this medicine.

#### **Atorvastatin Billev Pharma contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

### **3. How to take Atorvastatin Billev Pharma**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Before starting treatment, your doctor will place you on a low-cholesterol diet, which you should maintain also during therapy with Atorvastatin Billev Pharma.

The usual starting dose of Atorvastatin Billev Pharma is 10 mg once a day in adults and children aged 10 years or older. This may be increased if necessary by your doctor until you are taking the amount you need. Your doctor will adapt the dose at intervals of 4 weeks or more. The maximum dose of Atorvastatin Billev Pharma is 80 mg once daily for adults and 20 mg once daily for children.

Atorvastatin Billev Pharma tablets should be swallowed whole with a drink of water, and can be taken at any time of day, with or without food. However, try to take your tablet at the same time every day.

#### **The duration of treatment with Atorvastatin Billev Pharma is determined by your doctor.**

Please ask your doctor if you think that the effect of Atorvastatin Billev Pharma is too strong or too weak.

#### **If you take more Atorvastatin Billev Pharma than you should**

If you accidentally take too many Atorvastatin Billev Pharma tablets (more than your usual daily dose), contact your doctor or nearest hospital for advice.

#### **If you forget to take Atorvastatin Billev Pharma**

If you forget to take a dose, just take your next scheduled dose at the correct time. Do not take a double dose to make up for a forgotten dose.

#### **If you stop taking Atorvastatin Billev Pharma**

If you have any further questions on the use of this medicine or wish to stop your treatment, ask your doctor or pharmacist.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If you experience any of the following serious side effects, stop taking your tablets and tell your doctor immediately or go to the nearest hospital accident and emergency department.**

Rare (may affect up to 1 in 1,000 people):

- Serious allergic reaction which causes swelling of the face, tongue and throat that can cause great difficulty in breathing.
- Serious illness with severe peeling and swelling of the skin, blistering of the skin, mouth, eyes genitals and fever. Skin rash with pink-red blotches especially on palms of hands or soles of feet which may blister.
- Muscle weakness, tenderness or pain and particularly, if at the same time, you feel unwell or have a high temperature it may be caused by an abnormal muscle breakdown. The abnormal muscle breakdown does not always go away, even after you have stopped taking atorvastatin, and it can be life-threatening and lead to kidney problems.

Very rare (may affect up to 1 in 10,000 people):

- If you experience problems with unexpected or unusual bleeding or bruising, this may be suggestive of a liver complaint. You should consult your doctor as soon as possible.

#### **Other possible side effects with Atorvastatin Billev Pharma:**

Common side effects (may affect up to 1 in 10 people) include:

- inflammation of the nasal passages, pain in the throat, nose bleed
- allergic reactions
- increases in blood sugar levels (if you have diabetes continue careful monitoring of your blood sugar levels), increase in blood creatine kinase
- headache
- nausea, constipation, wind, indigestion, diarrhoea
- joint pain, muscle pain and back pain
- blood test results that show your liver function can become abnormal

Uncommon side effects (may affect up to 1 in 100 people) include:

- anorexia (loss of appetite), weight gain, decreases in blood sugar levels (if you have diabetes you should continue careful monitoring of your blood sugar levels)
- having nightmares, insomnia
- dizziness, numbness or tingling in the fingers and toes, reductions of sensation to pain or touch, change in sense of taste, loss of memory
- blurred vision
- ringing in the ears and/or head
- vomiting, belching, abdominal pain upper and lower, pancreatitis (inflammation of the pancreas leading to stomach pain)

- hepatitis (liver inflammation)
- rash, skin rash and itching, hives, hair loss
- neck pain, muscle fatigue
- fatigue, feeling unwell, weakness, chest pain, swelling especially in the ankles (oedema), raised temperature
- urine tests that are positive for white blood cells

Rare side effects (may affect up to 1 in 1,000 people) include:

- visual disturbance
- unexpected bleeding or bruising
- cholestasis (yellowing of the skin and whites of the eyes)
- tendon injury

Very rare side effects (may affect up to 1 in 10,000 people) include:

- an allergic reaction – symptoms may include sudden wheezing and chest pain or tightness, swelling of the eyelids, face, lips, mouth, tongue or throat, difficulty breathing, collapse
- hearing loss
- gynecomastia (breast enlargement in men)

Possible side effects reported with some statins (medicines of the same type):

- Sexual difficulties
- Depression
- Breathing problems including persistent cough and/or shortness of breath or fever
- Diabetes. This is more likely if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure. Your doctor will monitor you while you are taking this medicine.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

HPRA Pharmacovigilance

Earlsfort Terrace

IRL - Dublin 2

Tel: +353 1 6764971

Fax: +353 1 6762517

Website: [www.hpra.ie](http://www.hpra.ie)

e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Atorvastatin Billev Pharma**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the packaging after EXP. The expiry date refers to the last day of that month.

Store in the original package in order to protect from light and moisture.

This medicine does not require any special temperature storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Atorvastatin Billev Pharma contains

- The active substance is atorvastatin. Each film-coated tablet contains 10 mg, 20 mg or 40 mg atorvastatin as atorvastatin calcium.
- The other ingredients are sodium hydroxide, sodium laurilsulfate, hydroxypropylcellulose, lactose monohydrate (see section 2 “Atorvastatin Billev Pharma contains lactose”), microcrystalline cellulose, croscarmellose sodium, crospovidone, magnesium stearate in the tablet core and polyvinyl alcohol, titanium dioxide (E171), macrogol 3000 and talc in the film-coating.

### What Atorvastatin Billev Pharma looks like and contents of the pack

10 mg film-coated tablets are white, round, slightly convex and bevel-edged, with diameter 6 mm.  
20 mg film-coated tablets are white, round, slightly convex and bevel-edged, with diameter 8 mm.  
40 mg film-coated tablets are white, round, slightly convex and bevel-edged, with diameter 10 mm.

Boxes of 4, 7, 10, 14, 20, 28, 30, 50, 56, 60, 84, 90, 98 and 100 film-coated tablets in blisters packs are available.

Not all pack sizes may be marketed.

### Marketing Authorisation Holder

Billev Pharma Aps, Fuglebækgaard, Elmegårdsvej 1A, Tørslev, DK-3630 Jægerspris, Denmark

### Manufacturer

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

### This medicinal product is authorised in the member states of the EEA under the following names:

Name of the Member State	Name of the medicinal product
Sveden	Atovans
Netherlands	Atorvastatine Jacobsen
United Kingdom	Atorvastatin
Bulgaria, Estonia, Greece, Lithuania, Latvia, Romania, Slovakia	Atorvastatin Billev
Hungary, Poland	Atorvastatin Krka
Malta	Atorvastatin TAD
Slovenia, Ireland	Atorvastatin Billev Pharma