

Pharmacode position may change as per Supplier's m/c requirement & additional small pharma code may appear on the front / back panel



#### Package leaflet: Information for the user

**Levetiracetam Aurobindo 250 mg film-coated tablets**  
**Levetiracetam Aurobindo 500 mg film-coated tablets**  
**Levetiracetam Aurobindo 750 mg film-coated tablets**  
**Levetiracetam Aurobindo 1000 mg film-coated tablets**  
Levetiracetam

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4

#### What is in this leaflet

1. What Levetiracetam Aurobindo is and what it is used for
2. What you need to know before you take Levetiracetam Aurobindo
3. How to take Levetiracetam Aurobindo
4. Possible side effects
5. How to store Levetiracetam Aurobindo
6. Contents of the pack and other information

#### 1. What Levetiracetam Aurobindo is and what it is used for

Levetiracetam is an antiepileptic medicine (a medicine used to treat seizures in epilepsy).

Levetiracetam Aurobindo is used

- on its own in adults and adolescents from 16 years of age with newly diagnosed epilepsy, to treat a certain form of epilepsy. Epilepsy is a condition where the patients have repeated fits (seizures). Levetiracetam is used for the epilepsy form in which the fits initially affect only one side of the brain, but could thereafter extend to larger areas on both sides of the brain (partial onset seizures with or without secondary generalisation). Levetiracetam has been given to you by your doctor to reduce the number of fits.
- as an add-on to other antiepileptic medicines to treat:
- partial onset seizures with or without generalisation in adults, adolescents, children and infants from one month of age
- myoclonic seizures (short, shock-like jerks of a muscle or group of muscles) in adults and adolescents from 12 years of age with juvenile myoclonic epilepsy.
- primary generalised tonic-clonic seizures (major fits, including loss of consciousness) in adults and adolescents from 12 years of age with idiopathic generalised epilepsy (the type of epilepsy that is thought to have a genetic cause).

#### 2. What you need to know before you take Levetiracetam Aurobindo

##### Do not take Levetiracetam Aurobindo:

- If you are allergic to levetiracetam pyrrolidone derivatives or any of the other ingredients of this medicine (listed in Section 6).

##### Warnings and precautions

Talk to your doctor or pharmacist before taking Levetiracetam Aurobindo:

- If you suffer from kidney problems, follow your doctor's instructions. He/she may decide if your dose should be adjusted.
- If you notice any slow down in the growth or unexpected puberty development of your child, please contact your doctor.
- A small number of people being treated with anti-epileptics such as Levetiracetam Aurobindo have had thoughts of harming or killing themselves. If you have any symptoms of depression and/or suicidal ideation, please contact your doctor.

##### Children and adolescents

- Levetiracetam Aurobindo is not indicated in children and adolescents below 16 years on its own (monotherapy)

##### Other medicines and Levetiracetam Aurobindo

Tell your doctor or pharmacist if you are taking or have recently taken or might take any other medicines.

Do not take macrogol (a drug used as laxative) for one hour before and one hour after taking levetiracetam as this may result in a loss of its effect.

##### Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Levetiracetam Aurobindo should not be used during pregnancy unless clearly necessary. A risk of birth defects for your unborn child cannot be completely excluded... Levetiracetam has shown unwanted reproductive effects in animal studies at dose levels higher than you would need to control your seizures.

Breast-feeding is not recommended during treatment.

##### Driving and using machines:

Levetiracetam Aurobindo may impair your ability to drive or operate any tools or machinery, as it may make you feel sleepy. This is more likely at the beginning of treatment or after an increase in the dose. You should not drive or use machines until it is established that your ability to perform such activities is not affected.

**Levetiracetam Aurobindo contains Sunset Yellow FCF (E110):** Levetiracetam Aurobindo 750 mg contains Sunset Yellow FCF (E110). It may cause allergic reactions.

#### 3. How to take Levetiracetam Aurobindo

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Take the number of tablets following your doctor's instructions. Levetiracetam Aurobindo must be taken twice a day, once in the morning and once in the evening, at about the same time each day.

##### Monotherapy

##### Dose in adults and adolescents (from 16 years of age):

General dose: between 1000 mg and 3,000 mg each day.

When you will first start taking Levetiracetam Aurobindo, your doctor will prescribe you a **lower dose** during 2 weeks before giving you the lowest general dose.

*Example: if your daily dose is 1000 mg, your reduced starting dose is 2 tablets of 250 mg in the morning and 2 tablets of 250 mg in the evening.*

##### Add-on therapy

##### Dose in adults and adolescents (12 to 17 years) weighing 50 kg or more:

• General dose: between 1,000 mg and 3,000 mg each day.

*Example: if your daily dose is 1,000 mg, you might take 2 tablets of 250 mg in the morning and 2 tablets of 250 mg in the evening.*

##### Dose in infants (1 month to 23 months), children (2 to 11 years) and adolescents (12 to 17 years) weighing less than 50 kg:

Your doctor will prescribe the most appropriate pharmaceutical form of Levetiracetam according to the age, weight and dose.

Levetiracetam 100 mg/ml oral solution is a formulation more appropriate to infants and children under the age of 6 years and to children and adolescent (from 6 to 17 years) weighing less than 50kg and when tablets don't allow accurate dosage.

##### Method of administration:

Swallow Levetiracetam Aurobindo tablets with a sufficient quantity of liquid (e.g. a glass of water).

You may take Levetiracetam Aurobindo with or without food

##### Duration of treatment:

- Levetiracetam Aurobindo is used as a chronic treatment. You should continue Levetiracetam Aurobindo treatment for as long as your doctor has told you.
- Do not stop your treatment without your doctor's advice as this could increase your seizures.

##### If you take more Levetiracetam Aurobindo than you should:

The possible side effects of an overdose of Levetiracetam Aurobindo are sleepiness, agitation, aggression, decrease of alertness, inhibition of breathing and coma. Contact your doctor if you took more tablets than you should. Your doctor will establish the best possible treatment of overdose.

##### If you forget to take Levetiracetam Aurobindo:

Contact your doctor if you have missed one or more doses. Do not take a double dose to make up for a forgotten tablet.


##### If you stop taking Levetiracetam Aurobindo:

If stopping treatment, Levetiracetam Aurobindo should be discontinued gradually to avoid an increase of seizures. Should your doctor decide to stop your Levetiracetam Aurobindo treatment, he/she will instruct you about the gradual withdrawal of Levetiracetam Aurobindo.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.



The most frequently reported side effects are nasopharyngitis, somnolence (sleepiness), headache, fatigue and dizziness. At the beginning of the treatment or at dose increase side effects like sleepiness, tiredness and dizziness may be more common. These effects should however decrease over time.

**Very common:** may affect more than 1 in 10 people

- nasopharyngitis;
- somnolence (sleepiness), headache.

**Common:** may affect up to 1 in 10 people

- anorexia (loss of appetite);
- depression, hostility or aggression, anxiety, insomnia, nervousness or irritability;
- convulsion, balance disorder (equilibrium disorder), dizziness (sensation of unsteadiness), lethargy (lack of energy and enthusiasm), tremor (involuntary trembling);
- vertigo (sensation of rotation);
- cough;
- abdominal pain, diarrhoea, dyspepsia (indigestion), vomiting, nausea;
- rash;
- asthenia/fatigue (tiredness).

**Uncommon:** may affect up to 1 in 100 people

- decreased number of blood platelets, decreased number of white blood cells;
- weight decrease, weight increase;
- suicide attempt and suicidal ideation, mental disorder, abnormal behaviour, hallucination, anger, confusion, panic attack, emotional instability/mood swings, agitation;
- amnesia (loss of memory), memory impairment (forgetfulness), abnormal coordination/ataxia (impaired coordinated movements), paraesthesia (tingling), disturbance in attention (loss of concentration);
- diplopia (double vision), vision blurred;
- elevated/abnormal values in a liver function test
- hair loss, eczema, pruritus;
- muscle weakness, myalgia (muscle pain);
- injury.

**Rare:** may affect up to 1 in 1,000 people

- infection;
- decreased number of all blood cell types;
- severe allergic reactions (DRESS anaphylactic reaction [severe and important allergic reaction], Quincke's oedema [swelling of the face, lips, tongue and throat]);
- decreased blood sodium concentration;
- suicide, personality disorders (behavioural problems), thinking abnormal (slow thinking, unable to concentrate);
- uncontrollable muscle spasms affecting the head, torso and limbs, difficulty in controlling movements, hyperkinesia (hyperactivity);
- pancreatitis;
- liver failure, hepatitis;
- skin rash, which may form blisters and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) (*erythema multiforme*), a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (*Stevens-Johnson syndrome*), and a more severe form causing skin peeling in more than 30% of the body surface (*toxic epidermal necrolysis*).

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

#### Ireland

HPRA Pharmacovigilance  
Earlsfort Terrace  
IRL - Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.hpra.ie](http://www.hpra.ie)  
e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

#### 5. How to store Levetiracetam Aurobindo

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label, carton & bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### 6. Contents of the pack and other information

##### What Levetiracetam Aurobindo contains

The active substance is levetiracetam. Each film-coated tablet contains 250 mg/ 500 mg/ 750 mg/ 1000 mg levetiracetam.

The other ingredients are:

Tablet core: Maize starch, Silica colloidal anhydrous (E551), Povidone (K-30) (E1201), Talc (E553b), Magnesium stearate (E470b).

Film-coating:

**250 mg:** Hypromellose 3cp & 6cp (E464), titanium dioxide (E 171), macrogol 4000, indigo carmine aluminium lake (E132).

**500 mg:** Hypromellose 3cp & 6cp (E464), titanium Dioxide (E 171), macrogol 4000, iron oxide yellow (E172).

**750 mg:** Hypromellose 3cp & 6cp (E464), titanium Dioxide (E 171), macrogol 4000, indigo carmine aluminium lake (E132), sunset yellow aluminium lake (E110), iron oxide red. (E172)

**1000 mg:** Hypromellose 5 cp (E464), titanium dioxide (E 171), macrogol 400.

##### What Levetiracetam Aurobindo looks like and contents of the pack

Film-coated tablet.

##### Levetiracetam Aurobindo 250 mg film-coated tablets

Blue oval shaped biconvex film-coated tablets debossed with a deep break line separating 'E' and '10' on one side and plain on the other side. The tablet can be divided into equal doses.

##### Levetiracetam Aurobindo 500 mg film-coated tablets

Yellow oval shaped biconvex film-coated tablets debossed with a deep break line separating 'E' and '11' on one side and plain on the other side. The tablet can be divided into equal doses.

##### Levetiracetam Aurobindo 750 mg film-coated tablets

Orange oval shaped biconvex film-coated tablets debossed with a deep break line separating 'E' and '12' on one side and plain on the other side. The tablet can be divided into equal doses.

##### Levetiracetam Aurobindo 1000 mg film-coated tablets

White modified oval shaped, biconvex film-coated tablets debossed with a deep break line separating 'E' and '13' on one side and plain on the other side. The tablet can be divided into equal doses.

Levetiracetam Aurobindo film-coated tablets are packaged in blister pack or HDPE bottle.

##### Pack sizes:

Blister pack: 20, 30, 50, 60, 100, 200 and 500 film-coated tablets  
Bottle pack: 30, 100, 200 and 500 film-coated tablets

Not all pack sizes may be marketed.

##### Marketing Authorisation Holder

Milpharm Limited  
Ares, Odyssey Business Park  
West End Road  
South Ruislip HA4 6QD  
United Kingdom

##### Manufacturer

APL Swift Services (Malta) Limited  
HF26, Hal Far Industrial Estate, Hal Far  
Birzebbugia, BBG 3000  
Malta

**This medicinal product is authorised in the Member States of the EEA under the following names:**

|                |  |
|----------------|--|
| France:        | Levetiracetam Arrow Lab 250 mg/ 500 mg/ 1000 mg, comprimé pelliculé sécable              |
| Germany        | Levetiracetam Aurobindo 250 mg/ 500 mg/ 750 mg/ 1000 mg Filmtabletten                    |
| Ireland        | Levetiracetam Aurobindo 250 mg/ 500 mg/ 750 mg/ 1000 mg film-coated tablets              |
| Italy          | Levetiracetam Aurobindo 500 mg/ 1000 mg compresse rivestite con film                     |
| Netherlands    | Levetiracetam Aurobindo 250 mg/ 500 mg/ 750 mg/ 1000 mg. filmomhulde tabletten           |
| Poland         | Levetiracetam Aurobindo  |
| Romania        | Levetiracetam Aurobindo 250 mg/ 500 mg/ 1000 mg comprimate filmate                       |
| Spain          | Levetiracetam Aurobindo 250 mg/ 500 mg/ 1000 mg comprimidos recubiertos con película EFG |
| United Kingdom | Levetiracetam Milpharm 250 mg/ 500 mg/ 750 mg/ 1000 mg film-coated tablets               |

This leaflet was last revised in 04/2016.

