

Package leaflet: information for the user

Risedronate Aurobindo Once a Week 35 mg film-coated tablets risedronate sodium

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their sign of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Risedronate Aurobindo Once a Week is and what it is used for
2. What you need to know before you take Risedronate Aurobindo Once a Week
3. How to take Risedronate Aurobindo Once a Week
4. Possible side effects
5. How to store Risedronate Aurobindo Once a Week
6. Contents of the pack and other information

1. What Risedronate Aurobindo is and what it is used for

What Risedronate Aurobindo Once a Week is

Risedronate Aurobindo Once a Week belongs to a group of non-hormonal medicines called bisphosphonates which are used to treat bone diseases. It works directly on your bones to make them stronger and therefore less likely to break.

Bone is a living tissue. Old bone is constantly removed from your skeleton and replaced with new bone.

Postmenopausal osteoporosis is a condition occurring in women after the menopause where the bones become weaker, more fragile and more likely to break after a fall or strain.

Osteoporosis can also occur in men due to a number of causes including ageing and/or a low level of the male hormone, testosterone.

The spine, hip and wrist are the most likely bones to break, although this can happen to any bone in your body. Osteoporosis – related fractures can also cause back pain, height loss and a curved back. Many patients with osteoporosis have no symptoms and you may not even have known that you had it.

What Risedronate Aurobindo Once a Week is used for

The treatment of osteoporosis

- in postmenopausal women, even if osteoporosis is severe. It reduces the risk of spinal and hip fractures.
- in men.

2. What you need to know before you take Risedronate Aurobindo

Do not take Risedronate Aurobindo Once a Week

- If you are allergic to risedronate sodium or any of the other ingredients of this medicine (listed in section 6)
- If your doctor has told you that you have a condition called hypocalcaemia (a low blood calcium level)

- If you may be pregnant, are pregnant or planning to become pregnant
- If you are breast-feeding
- If you have severe kidney problems.

Warnings and precautions

Talk to your doctor or pharmacist before taking Risedronate Aurobindo

- If you are unable to stay in an upright position (sitting or standing) for at least 30 minutes.
- If you have abnormal bone and mineral metabolism (for example lack of vitamin D, parathyroid hormone abnormalities, both leading to a low blood calcium level).
- If you have had problems in the past with your oesophagus (the tube that connects your mouth with your stomach). For instance you may have had pain or difficulty in swallowing food or you have previously been told that you have Barrett's oesophagus (a condition associated with changes in the cells that line the lower oesophagus).
- If you have been told by your doctor that you have an intolerance to some sugars (such as lactose).
- If you have had or have pain, swelling or numbness of the jaw or a “heavy jaw feeling” or loosening of a tooth.
- If you are under dental treatment or will undergo dental surgery, tell your dentist that you are being treated with Risedronate Aurobindo Once a Week.

Your doctor will advise you on what to do when taking Risedronate Aurobindo Once a Week if you have any of the above.

Children and adolescents

Risedronate is not recommended for use in children and adolescents below age 18 due to insufficient data on safety and efficacy.

Other medicines and Risedronate Aurobindo

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Medicines containing one of the following lessen the effect of Risedronate Aurobindo Once a Week if taken at the same time:

- calcium
- magnesium
- aluminium (for example some indigestion mixtures)
- iron

Take these medicines at least 30 minutes after your Risedronate Aurobindo Once a Week tablet.

Risedronate Aurobindo with food and drink

It is very important that you do NOT take your Risedronate Aurobindo Once a Week tablet with food or drinks (other than plain water) so that it can work properly. In particular do not take this medicine at the same time as dairy products (such as milk) as they contain calcium (see section 2, “Other medicines and Risedronate Aurobindo”).

Take food and drinks (other than plain water) at least 30 minutes after your Risedronate Aurobindo Once a Week tablet.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Do NOT take Risedronate Aurobindo Once a Week if you may be pregnant, are pregnant or planning to become pregnant (see section 2, “Do not take Risedronate Aurobindo Once a Week”). The potential risk associated with the use of risedronate sodium (active substance in Risedronate Aurobindo Once a Week) in pregnant women is unknown.

Do NOT take Risedronate Aurobindo Once a Week if you are breast-feeding (see section 2, “Do not take Risedronate Aurobindo Once a Week”).

Risedronate Aurobindo Once a Week should only be used to treat postmenopausal women and men. Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Risedronate Aurobindo Once a Week is not known to affect your ability to drive and use machines.

Risedronate Aurobindo Once a Week contains lactose.

Risedronate Aurobindo Once a Week contains a small amount of lactose (see section 2, “Warnings and precautions”). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Risedronate Aurobindo

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

The recommended dose is:

Take ONE Risedronate Aurobindo Once a Week tablet (35 mg of risedronate sodium) once a week.

Choose one day of the week that best fits your schedule. Every week, take the Risedronate Aurobindo Once a Week tablet on your chosen day.

For your convenience, so that you take your tablet on the right day every week, there is a feature included with the Risedronate Aurobindo Once a Week pack:

When to take the Risedronate Aurobindo Once a Week tablet

Take your Risedronate Aurobindo Once a Week tablet at least 30 minutes before the first food, drink (other than plain water) or other medicine of the day.

How to take the Risedronate Aurobindo Once a Week tablet

- Take the tablet whilst you are in an upright position (you may sit or stand) to avoid heartburn.
- Swallow it with at least one glass (120 ml) of plain water.
- Swallow it whole. Do not suck or chew it.
- Do not lie down for 30 minutes after taking your tablet.

Your doctor will tell you if you need calcium and vitamin supplements, if you are not taking enough from your diet.

If you take more Risedronate Aurobindo Once a Week than you should

If you or somebody else has accidentally taken more Risedronate Aurobindo Once a Week tablets than prescribed, drink one full glass of milk and seek medical attention.

If you forget to take Risedronate Aurobindo Once a Week

If you have forgotten to take your tablet on your chosen day, take it on the day you remember. Return to taking one tablet once a week on the day the tablet is normally taken.

Do not take a double dose to make up for a forgotten dose.

If you stop taking Risedronate Aurobindo Once a Week

If you stop treatment you may begin to lose bone mass. Please talk to your doctor before you consider stopping treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Risedronate Aurobindo and contact a doctor immediately if you experience any of the following:

- Symptoms of a severe allergic reaction such as;
 - Swelling of face, tongue or throat
 - Difficulties in swallowing
 - Hives and difficulties in breathing
- Severe skin reactions that can include blistering of the skin.

Tell your doctor promptly if you experience the following side effects:

- Eye inflammation, usually with pain, redness and light sensitivity.
- Bone necrosis of the jaw (osteonecrosis) associated with delayed healing and infection, often following tooth extraction (see section 2, “warnings and precautions”).
- Symptoms from oesophagus such as pain when you swallow, difficulties in swallowing, chest pain or new or worsened heartburn.

However in clinical studies the other side effects that were observed were usually mild and did not cause the patient to stop taking their tablets.

Common (may affect up to 1 in 10 people)

- Indigestion, feeling sick, stomach ache, stomach cramps or discomfort, constipation, feelings of fullness, bloating, diarrhoea.
- Pain in your bones, muscles or joints.
- Headache.

Uncommon (may affect up to 1 in 100 people)

- Inflammation or ulcer of the oesophagus (the tube that connects your mouth with your stomach) causing difficulty and pain in swallowing (see also section 2, “Warnings and precautions”), inflammation of the stomach and duodenum (bowel draining the stomach).
- Inflammation of the coloured part of the eye (iris) (red painful eyes with a possible change in vision).

Rare (may affect up to 1 in 1000 people)

- Inflammation of the tongue (red swollen, possibly painful), narrowing of the oesophagus (the tube that connects your mouth with your stomach).
- Abnormal liver tests have been reported. These can only be diagnosed from a blood test.

Very rare (may affect up to 1 in 10,000 people):

- Talk to your doctor if you have ear pain, discharge from the ear, and/or an ear infection. These could be signs of bone damage in the ear.

During post-marketing experience, the following have been reported (unknown frequency);

- Hair loss
- Liver disorders, some cases were severe.

Rarely, at the beginning of treatment, a patient’s blood calcium and phosphate levels may fall. These changes are usually small and cause no symptoms.

Unusual fracture of the thigh bone particularly in patients on long-term treatment for osteoporosis may occur rarely. Contact your doctor if you experience pain, weakness or discomfort in your thigh, hip or groin as this may be an early indication of a possible fracture of the thigh bone.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via

HPRA Pharmacovigilance

Earlsfort Terrace

IRL - Dublin 2

Tel: +353 1 6764971

Fax: +353 1 6762517

Website: www.hpra.ie

e-mail: medsafety@hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Risedronate Aurobindo

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines no you longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Risedronate Aurobindo Once a Week contains

The active substance is risedronate sodium. Each film-coated tablet contains 35 mg risedronate sodium (equivalent to 32.4 mg risedronic acid).

The other ingredients are:

Tablet core: Lactose monohydrate, crospovidone, hydroxy propyl cellulose, magnesium stearate and cellulose microcrystalline.

Film coating: Hypromellose, titanium dioxide (E171), macrogol 400, hydroxy propyl cellulose, iron oxide yellow (E172), macrogol 8000, iron oxide red (E172) and silica colloidal anhydrous.

What Risedronate Aurobindo Once a Week looks like and contents of the pack

Film-coated tablet.

Light orange coloured, circular shaped film coated biconvex tablets debossed with 'F27' on one side and plain on the other side.

Transparent PVC / PE / PVdC / Aluminium blisters in a cardboard box, Packs 1, 2, 4, 10, 12 or 16 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Milpharm Limited

Ares, Odyssey Business Park

West End Road

South Ruislip HA4 6QD

United Kingdom

Manufacturer

APL Swift Services (Malta) Limited
HF26, Hal Far Industrial Estate, Hal Far
Birzebbugia, BBG 3000
Malta

or

Milpharm Limited,
Ares Block, Odyssey Business Park,
West End Road, Rusilp HA46QD.
United Kingdom

This medicinal product is authorised in the Member States of the EEA under the following names:

France	RISEDRONATE ARROW 35 mg, comprimé pelliculé
Germany	Risedronat Aurobindo einmal wöchentlich 35 mg Filmtabletten
Ireland	Risedronate Aurobindo Once a Week 35 mg film-coated tablets
Italy	Risedronato Aurobindo 35 mg
Malta	Risedronate Aurobindo Once a Week 35 mg film-coated tablets
Romania	Risedronat Aurobindo 35 mg comprimate filmate
Spain	Risedronat Semanal Aurobindo 35 mg comprimidos recubiertos con película EFG
The Netherlands	Risedronaatnatrium Aurobindo Wekelijks 35 mg, filmomhulde tabletten
United Kingdom.	Risedronate sodium 35mg film-coated tablets

This leaflet was last revised in 09/2016