

PACKAGE LEAFLET

Package leaflet: Information for the user

Amlodipine Pinewood 5 mg tablets

Amlodipine Pinewood 10 mg tablets

amlodipine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Amlodipine Pinewood is and what it is used for
2. What you need to know before you take Amlodipine Pinewood
3. How to take Amlodipine Pinewood
4. Possible side effects
5. How to store Amlodipine Pinewood
6. Contents of the pack and other information

1. What Amlodipine Pinewood is and what it is used for

Amlodipine Pinewood tablets contain the active substance amlodipine which belongs to a group of medicines called calcium antagonists.

Amlodipine Pinewood is used to treat high blood pressure (hypertension) or a certain type of chest pain called angina, a rare form of which is Prinzmetal's or variant angina.

In patients with high blood pressure your medicine works by relaxing blood vessels, so that blood passes through them more easily. In patients with angina Amlodipine Pinewood works by improving blood supply to the heart muscle which then receives more oxygen and as a result chest pain is prevented. Your medicine does not provide immediate relief of chest pain from angina.

2. What you need to know before you take Amlodipine Pinewood

Do not take Amlodipine Pinewood:

- if you are allergic to amlodipine or any of the other ingredients of this medicine (listed in section 6), or to any other calcium antagonists. This may be itching, reddening of the skin or difficulty in breathing.
- if you have severe low blood pressure (hypotension)
- if you have narrowing of the aortic heart valve (aortic stenosis) or cardiogenic shock (a condition where your heart is unable to supply enough blood to the body).
- if you suffer from heart failure after a heart attack.

Warnings and precautions

Talk to your doctor or pharmacist before taking Amlodipine Pinewood.

You should inform your doctor if you have or have had any of the following conditions:

- Recent heart attack
- Heart failure
- Severe increase in blood pressure (Hypertensive crisis)
- Liver disease
- You are elderly and your dose needs to be increased

Children and adolescents

Amlodipine Pinewood has not been studied in children under the age of 6 years. Amlodipine Pinewood should only be used for hypertension in children and adolescents from 6 years to 17 years of age (see section 3). For more information, talk to your doctor.

Other medicines and Amlodipine Pinewood

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Amlodipine Pinewood may affect or be affected by other medicines, such as:

- ketoconazole, itraconazole (anti-fungal medicines)
- ritonavir, indinavir, nelfinavir (so called protease inhibitors used to treat HIV)
- rifampicin, erythromycin, clarithromycin (antibiotics)
- hypericum perforatum (St. John's Wort)
- verapamil, diltiazem (heart medicines)
- dantrolene (infusion for severe body temperature abnormalities)
- tacrolimus, sirolimus, temsirolimus, and everolimus (medicine used to alter the way your immune system works)
- simvastatin (a cholesterol lowering medicine)
- cyclosporine (an immunosuppressant)

Amlodipine Pinewood may lower your blood pressure even more if you are already taking other medicines to treat your high blood pressure.

Amlodipine Pinewood with food and drink

Grapefruit juice and grapefruit should not be consumed by people who are taking Amlodipine Pinewood. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active ingredient amlodipine, which can cause an unpredictable increase in the blood pressure lowering effect of Amlodipine Pinewood.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

The safety of amlodipine in human pregnancy has not been established. If you think you might be pregnant, or are planning to get pregnant, you must tell your doctor before you take Amlodipine Pinewood.

Breast-feeding:

Amlodipine Pinewood has been shown to pass into breast milk in small amounts. If you are breast-feeding or about to start breast-feeding you must tell your doctor before taking Amlodipine Pinewood.

Driving and using machines

Amlodipine Pinewood may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Amlodipine Pinewood contains sodium.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, which means it is essentially ‘sodium-free’.

3. How to take Amlodipine Pinewood

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended initial dose is Amlodipine Pinewood 5 mg once daily. The dose can be increased to Amlodipine Pinewood 10 mg once daily.

Your medicine can be used before or after food and drinks. You should take your medicine at the same time each day with a drink of water. Do not take Amlodipine Pinewood with grapefruit juice.

Use in children and adolescents

For children and adolescents (6 -17 years old), the recommended usual starting dose is 2.5 mg a day. The maximum recommended dose is 5 mg a day. The 2.5 mg dose can be obtained by breaking the Amlodipine Pinewood 5mg tablet along the breakline, and taking half the tablet.

It is important to keep taking the tablets. Do not wait until your tablets are finished before seeing your doctor.

If you take more Amlodipine Pinewood than you should

Taking too many tablets may cause your blood pressure to become low or even dangerously low. You may feel dizzy, lightheaded, faint or weak. If blood pressure drop is severe enough shock can occur. Your skin could feel cool and clammy and you could lose consciousness.

Excess fluid may accumulate in your lungs (pulmonary oedema) causing shortness of breath that may develop up to 24-48 hours after intake.

Seek immediate medical attention if you take too many Amlodipine Pinewood tablets.

If you forget to take Amlodipine Pinewood

Do not worry. If you forget to take a tablet, leave out that dose completely. Take your next dose at the right time. Do not take a double dose to make up for a missed dose.

If you stop taking Amlodipine Pinewood

Your doctor will advise you how long to take your medicine. Your condition may return if you stop using your medicine before you are advised.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Visit your doctor **immediately** if you experience any of the following side effects after taking this medicine.

- Sudden wheeziness, chest pain, shortness of breath or difficulty in breathing

- Swelling of eyelids, face or lips
- Swelling of the tongue and throat which causes great difficulty breathing
- Severe skin reactions including intense skin rash, hives, reddening of the skin over your whole body, severe itching, blistering, peeling and swelling of the skin, inflammation of mucous membranes (Stevens Johnson Syndrome, toxic epidermal necrolysis) or other allergic reactions
- Heart attack, abnormal heart beat
- Inflamed pancreas which may cause severe abdominal and back pain accompanied with feeling very unwell

The following **very common side effect** has been reported. If this causes you problems or if it **lasts for more than one week**, you should **contact your doctor**.

Very common: may affect more than 1 in 10 people

- Oedema (fluid retention)

The following **common side effects** have been reported. If any of these cause you problems or if they **last for more than one week**, you should contact your doctor.

Common: may affect up to 1 in 10 people

- Headache, dizziness, sleepiness (especially at the beginning of treatment)
- Palpitations (awareness of your heart beat), flushing
- Abdominal pain, feeling sick (nausea)
- Altered bowel habits, diarrhoea, constipation, indigestion
- Tiredness, weakness
- Visual disturbances, double vision
- Muscle cramps
- Ankle swelling

Other side-effects that have been reported include the following list. If any of these get serious, or if you notice any side-effects not listed in this leaflet, please tell your doctor or pharmacist.

Uncommon: may affect up to 1 in 100 people

- Mood changes, anxiety, depression, sleeplessness
- Trembling, taste abnormalities, fainting
- Numbness or tingling sensation in your limbs; loss of pain sensation
- Ringing in the ears
- Low blood pressure
- Sneezing/running nose caused by inflammation of the lining of the nose (rhinitis)
- Cough
- Dry mouth, vomiting (being sick)
- Hair loss, increased sweating, itchy skin, red patches on skin, skin discolouration
- Disorder in passing urine, increased need to urinate at night, increased number of times of passing urine
- Inability to obtain an erection; discomfort or enlargement of the breasts in men
- Pain, feeling unwell
- Joint or muscle pain, back pain
- Weight increase or decrease

Rare: may affect up to 1 in 1,000 people

- Confusion

Very rare: may affect up to 1 in 10,000 people

- Decreased numbers of white blood cells, decrease in blood platelets which may result in unusual bruising or easy bleeding (red blood cell damage)
- Excess sugar in the blood (hyperglycaemia)
- A disorder of the nerves which can cause muscular weakness, tingling or numbness
- Swelling of the gums, bleeding gums
- Abdominal bloating (gastritis)
- Abnormal liver function, inflammation of the liver (hepatitis), yellowing of the skin (jaundice), liver enzyme increase which may have an effect on some medical tests
- Increased muscle tension
- Inflammation of blood vessels, often with skin rash
- Sensitivity to light
- Disorders combining rigidity, tremor, and/or movement disorders

Not known: frequency cannot be estimated from the available data

- Trembling, rigid posture, mask-like face, slow movements and a shuffling, unbalanced walk

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects direct via HPRA Pharmacovigilance, Website: www.hpra.ie.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Amlodipine Pinewood

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton label and blister foil after 'EXP'. The expiry date refers to the last day of that month.

Store below 25°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Amlodipine Pinewood Tablets contain

- The active substance is amlodipine as amlodipine besilate
- Amlodipine Pinewood 5 mg: Each tablet contains 5 mg amlodipine as amlodipine besilate.
- Amlodipine Pinewood 10 mg: Each tablet contains 10 mg amlodipine as amlodipine besilate.
- The other ingredients are microcrystalline cellulose, calcium hydrogen phosphate, sodium starch glycolate (Type A) and magnesium stearate.

What Amlodipine Pinewood Tablets look like and contents of the pack

Amlodipine Pinewood 5 mg tablets:

White to off-white, round, flat faced, bevelled edged tablets debossed with 'U' and 'L' on either side of the breakline on one side and breakline on the other side.

Amlodipine Pinewood 10 mg tablets:

White to off-white, round, flat faced, bevelled edged tablets with 'UH' debossed on one side and plain on the other side.

The 5 mg and 10 mg tablets are available in blister packs of 10, 14, 15, 20, 28, 30, 50, 56, 60, 84, 90, 98 and 100 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Pinewood Laboratories Limited,
Ballymacarbry,
Clonmel,
Co. Tipperary,
Ireland

Manufacturer

Niche Generics Ltd,
Unit 5, 151 Baldoyle Industrial Estate,
Dublin 13,
Ireland.

This leaflet was last revised in 10/2025