

Reason for update: Type 1A variation to remove Ware as Batch Release Site (no impact to PIL artwork)  
Market: IE  
Agency Approval Date: N/A T1A  
Text Date: 09/02/2021  
Text Issue and Draft No.: Issue 6 draft 1


## GSK Logo

### Package Leaflet: Information for the User

# Serevent Diskus

50 microgram per metered dose inhalation powder, pre-dispensed

salmeterol (as xinafoate)



**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side-effects, talk to your doctor or pharmacist. This includes any possible side-effects not listed in this leaflet. See section 4.

#### **In this leaflet:**

- 1 What Serevent Diskus is and what it is used for
- 2 What you need to know before you use Serevent Diskus
- 3 How to use Serevent Diskus
- 4 Possible side effects
- 5 How to store Serevent Diskus
- 6 Contents of the pack and other information

## **1 What Serevent Diskus is and what it is used for**

- Serevent Diskus contains the medicine salmeterol. It is a 'long-acting bronchodilator'. It helps the airways in the lungs to stay open. This makes it easier for air to get in and out. The effects are usually felt within 10 to 20 minutes and last for 12 hours or more.
- The doctor has prescribed it to help prevent breathing problems. These could be caused by asthma. Taking Serevent Diskus regularly will help prevent asthma attacks. This also includes asthma brought on by exercise or at night.
- Taking Serevent Diskus regularly will also help prevent breathing problems caused by other chest illnesses such as Chronic Obstructive Pulmonary Disease (COPD), chronic bronchitis and emphysema.

Reason for update: Type 1A variation to remove Ware as Batch Release Site (no impact to PIL artwork)

Market: IE

Agency Approval Date: N/A – T1A

Text Date: 09/02/2021

Text Issue and Draft No.: Issue 6 draft 1

- Serevent Diskus helps to stop breathlessness and wheezing coming on. It does not work once you are breathless or wheezy. If that happens, you need to use a fast acting 'reliever' medicine, such as salbutamol.
- Serevent is supplied to you in an inhaler called the diskus. You breathe the medicine directly into your lungs.

**If you are being treated for asthma, you should always be given both a Serevent and a steroid inhaler to use together.**

## 2 What you need to know before you use Serevent Diskus

### **Do not take Serevent Diskus if:**

- you are allergic to salmeterol xinafoate, or to the other ingredient lactose monohydrate.

### **Warnings and Precautions**

Talk to your doctor or pharmacist before using Serevent Diskus.

- **If you are using Serevent for asthma your doctor will want to regularly check your symptoms.**
- **If your asthma or breathing gets worse tell your doctor straight away.** You may find that you feel more wheezy, your chest feels tight more often or you may need to use more of your fast acting 'reliever' medicine. If any of these happen, do not increase your number of puffs of Serevent. Your chest condition may be getting worse and you could become seriously ill. See your doctor as you may need a change in asthma treatment.
- **If you have breathing difficulties or wheezing that gets worse straight after taking Serevent, stop using your Serevent Diskus.** Use your fast-acting 'reliever' inhaler to help your breathing and **tell your doctor straight away**
- Once your asthma is well controlled your doctor may consider it appropriate to gradually reduce the dose of Serevent.
- If you have been prescribed Serevent for your asthma, continue to use any other asthma medication you are already taking. These could include a steroid inhaler or steroid tablets. Continue taking the same doses as before, unless your doctor tells you otherwise. Do this even if you feel much better. **Do not stop taking your steroid inhaler (or any steroid tablets) when you start using Serevent.**
- Your doctor may want to check your health regularly if you have an overactive thyroid gland, diabetes mellitus (Serevent may increase your blood sugar) or heart disease, including an irregular or fast heartbeat.
- If you are of black African or Afro-Caribbean ancestry, tell your doctor if your asthma symptoms remained uncontrolled or worsen whilst using Serevent.

### **Other medicines and Serevent**

Reason for update: Type 1A variation to remove Ware as Batch Release Site (no impact to PIL artwork)

Market: IE

Agency Approval Date: N/A – T1A

Text Date: 09/02/2021

Text Issue and Draft No.: Issue 6 draft 1

- Tell your doctor if you are taking, have recently taken or might take any other medicines. This includes those for asthma or any other medicines obtained without a prescription. This is because Serevent may not be suitable to be taken with other medicines.
- Please inform your doctor before using Serevent if you are currently being treated for any fungal infections with medicines containing ketoconazole or itraconazole or with the antibiotic telithromycin, or if you are being treated for HIV with ritonavir. These medicines may increase the risk of you experiencing side effects with Serevent, including irregular heartbeats, or may make side-effects worse.
- Beta-blockers should be avoided when taking Serevent, unless your doctor tells you to. Beta-blockers, include atenolol, propranolol and sotalol, are mostly used for high blood pressure or other heart conditions. Please tell your doctor if you are taking beta-blockers or have recently been prescribed beta-blockers as they may reduce or abolish the effects of salmeterol.
- Serevent can reduce the amount of potassium in your blood. If this happens you may notice an uneven heartbeat, muscle weakness or cramp. This is more likely to happen if you take Serevent with medicines used to treat high blood pressure (diuretics – water tablets) and other medicines used to treat breathing problems such as theophylline or steroids. Your doctor may ask for you to have blood tests to check the amount of potassium in your blood. If you have any concerns discuss them with your doctor.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

The possible side effects associated with Serevent are unlikely to affect your ability to drive or use machines.

### **Serevent Diskus contains lactose monohydrate**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

## **3 How to use Serevent Diskus**

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- **If you are being treated for asthma, you should always be given both a Serevent and a steroid inhaler to use together.**
- Use Serevent every day, until your doctor advises you to stop.
- You will start to feel your medicine working within the first day of use.

Serevent is for inhalation by mouth only.

Reason for update: Type 1A variation to remove Ware as Batch Release Site (no impact to PIL artwork)

Market: IE

Agency Approval Date: N/A – T1A

Text Date: 09/02/2021

Text Issue and Draft No.: Issue 6 draft 1

## **Adults with Asthma, Chronic Obstructive Pulmonary Disease (COPD) including bronchitis and emphysema**


- The usual starting dose is 1 puff twice a day.
- For people with more severe airways obstruction, your doctor may increase your dose to 2 puffs twice a day.

### **Instructions for use**

- Your doctor, nurse or pharmacist should show you how to use your inhaler. They should check how you use it from time to time. Not using the Serevent Diskus properly or as prescribed may mean that it will not help your asthma or COPD as it should.
- The Serevent Diskus device holds blisters containing Serevent as a powder.
- There is a counter on top of the Serevent Diskus which tells you how many doses are left. It counts down to 0. The numbers 5 to 0 will appear in red to warn you when there are only a few doses left. Once the counter shows 0, your inhaler is empty.

### **Using your Diskus**

- 1** To open your Serevent Diskus, hold the outer case in one hand and put the thumb of your other hand on the thumbgrip. Push your thumb away from you as far as it will go. You will hear a click. This will open a small hole in the mouthpiece.



- 2** Hold your Serevent Diskus with the mouthpiece towards you. You can hold it in either your right or left hand. Slide the lever away from you as far as it will go. You will hear a click. This places a dose of your medicine in the mouthpiece.



Every time the lever is pushed back a blister is opened inside and the powder made ready for you to inhale. Do not play with the lever as this opens the blisters and wastes medicine.

- 3** Hold the Serevent Diskus away from your mouth, breathe out as far as is comfortable. Do not breathe into your Serevent Diskus.

Reason for update: Type 1A variation to remove Ware as Batch Release Site (no impact to PIL artwork)

Market: IE

Agency Approval Date: N/A – T1A

Text Date: 09/02/2021

Text Issue and Draft No.: Issue 6 draft 1


- 4 Put the mouthpiece to your lips; breathe in steadily and deeply through the Serevent Diskus, not through your nose.

Remove the Serevent Diskus from your mouth.


Hold your breath for about 10 seconds or for as long as is comfortable.

Breathe out slowly.

You may not be able to taste or feel the powder on your tongue, even if you have used the Serevent Diskus correctly.



- 5 To close the Serevent Diskus, slide the thumb grip back towards you, as far as it will go. You will hear a click. The lever will return to its original position and is reset.



Your Serevent Diskus is now ready for you to use again.

### **Cleaning your inhaler**

Wipe the mouthpiece of the Serevent Diskus with a dry tissue to clean it.

### **If you use more Serevent than you should**

It is important to use the inhaler as instructed. If you accidentally take a larger dose than recommended, talk to your doctor or pharmacist. You may notice your heart beating faster than usual and that you feel shaky or dizzy. You may also have a headache, muscle weakness and aching joints.

### **If you forget to use Serevent**

Do not take a double dose to make up for a forgotten dose. Just take your next dose at the usual time.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4 Possible side effects**

Reason for update: Type 1A variation to remove Ware as Batch Release Site (no impact to PIL artwork)  
Market: IE  
Agency Approval Date: N/A – T1A  
Text Date: 09/02/2021  
Text Issue and Draft No.: Issue 6 draft 1

Like all medicines, this medicine can cause side effects, although not everybody gets them. To reduce the chances of side effects, your doctor will prescribe the lowest dose of Serevent to control your asthma or COPD. These are the side effects reported by people taking Serevent.

**Allergic reactions: you may notice your breathing suddenly gets worse after using Serevent.** You may be very wheezy and cough. You may also notice itching and swelling (usually of the face, lips, tongue or throat). If you get these effects or they happen suddenly after using Serevent, **tell your doctor straight away**. Allergic reactions to Serevent are very rare (they affect less than 1 person in 10,000).

Other side effects are listed below:

**Common (affects less than 1 person in 10):**

- Muscle cramps
- Feeling shaky; fast or uneven heartbeat (palpitations), headache, shaking hands (tremor). Tremors are more likely if you are taking more than one puff twice daily. These side effects do not last long and happen less as treatment with Serevent continues.

**Uncommon (affects less than 1 person in 100):**

- Rash
- Very fast heart rate (tachycardia). This is more likely to happen if you are taking more than one puff twice daily
- Feeling nervous.

**Rare (affects less than 1 person in 1,000):**

- Feeling dizzy
- Being unable to sleep or finding sleep difficult
- A reduction in the amount of potassium in your blood (you may get an uneven heartbeat, muscle weakness, cramp).

**Very rare (affects less than 1 person in 10,000):**

- **Breathing difficulties or wheezing that gets worse straight after taking Serevent.** If this happens **stop using your Serevent Diskus**. Use your fast-acting 'reliever' inhaler to help your breathing and **tell your doctor straight away**
- Uneven heartbeat or your heart gives an extra beat (arrhythmias). If this happens do not stop using Serevent but tell your doctor
- Sore mouth or throat
- Feeling sick (nausea)
- Aching, swollen joints, or chest pain.
- Increases in the amount of sugar (glucose) in your blood (hyperglycaemia). If you have diabetes, more frequent blood sugar monitoring and possibly adjustment of your usual diabetic treatment may be required.

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Website [www.hpra.ie](http://www.hpra.ie).

Reason for update: Type 1A variation to remove Ware as Batch Release Site (no impact to PIL artwork)  
Market: IE  
Agency Approval Date: N/A – T1A  
Text Date: 09/02/2021  
Text Issue and Draft No.: Issue 6 draft 1

By reporting side effects you can help provide more information on the safety of this medicine.

## 5 How to store Serevent Diskus

- **Keep this medicine out of the sight and reach of children.**
- Do not store above 30°C.
- Do not use this medicine after the expiry date which is stated on the label and carton after 'Exp'. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## 6 Contents of the pack and other information

### What Serevent Diskus contains

- The active substance is 50 micrograms salmeterol (as xinafoate).
- The other ingredient is lactose monohydrate (which contains milk proteins).

### What Serevent Diskus looks like and contents of the pack

- The Serevent Diskus contains a foil strip containing individual doses of a white inhalation powder. The foil protects the powder for inhalation from the effects of the atmosphere.
- Each dose is pre-dispensed.
- The devices are packed in cartons which hold one Diskus containing 60 inhalations

### Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:  
GlaxoSmithKline (Ireland) Ltd.  
12 Riverwalk  
Citywest Business Campus  
Dublin 24  
Ireland

Manufacturers:  
Glaxo Wellcome Production  
Zone Industrielle No. 2  
23 Rue Lavoisier  
27000 Evreux  
France

**This leaflet was last revised in May 2020.**

Trade marks are owned by or licensed to the GSK group of companies.

© 2020 GSK group of companies or its licensor