

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Priorix-Tetra powder and solvent for solution for injection in pre-filled syringe Measles, mumps, rubella and varicella vaccine (live)

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

After reconstitution, 1 dose (0.5 ml) contains:

Measles virus¹ Schwarz strain (live, attenuated) not less than $10^{3.0}$ CCID₅₀³

Mumps virus¹ RIT 4385 strain, derived from Jeryl Lynn strain (live, attenuated)
not less than $10^{4.4}$ CCID₅₀³

Rubella virus² Wistar RA 27/3 strain (live, attenuated)
not less than $10^{3.0}$ CCID₅₀³

Varicella virus² Oka strain (live, attenuated) not less than $10^{3.3}$ PFU⁴

¹ produced on chick embryo cells

² produced on human diploid (MRC-5) cells

³ Cell culture infective dose 50%

⁴ Plaque forming units

This vaccine contains a trace amount of neomycin. See section 4.3.

Excipients with known effect

The vaccine contains 14 mg of sorbitol, 6.5 nanograms of para-aminobenzoic acid, 583 micrograms of phenylalanine, 0.65 micrograms of polysorbate 80 (E 433) and 709 micrograms of proline per dose (see section 4.4).

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Powder and solvent for solution for injection in pre-filled syringe.

Before reconstitution, the powder is a whitish to slightly pink coloured cake, a portion of which may be yellowish. The solvent is a clear colourless liquid.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Priorix-Tetra is indicated for active immunisation of individuals from the age of 11 months against measles, mumps, rubella and varicella.

Use in infants aged 9-10 months could be considered under special circumstances. See section 4.2.

The use of Priorix-Tetra should be based on official recommendations.

4.2 Posology and method of administration

Posology

Individuals from 11 months of age

Individuals aged from 11 months should receive two doses (each of 0.5 ml) of Priorix-Tetra. The age at which individuals may receive Priorix-Tetra should reflect applicable official recommendations*, which vary according to the epidemiology of these diseases.

The dose interval should preferably* be between 6 weeks and 3 months. When the first dose is administered at 11 months of age, the second dose should be administered within 3 months. Under no circumstances should the dose interval be less than 4 weeks. See section 5.1.

Alternatively, and in accordance with applicable official recommendations*:

- A single dose of Priorix-Tetra may be administered to individuals who have already received a single dose of another measles, mumps and rubella (MMR) vaccine and/or a single dose of another varicella vaccine
- A single dose of Priorix-Tetra may be administered followed by a single dose of another measles, mumps and rubella (MMR) vaccine and/or a single dose of another varicella vaccine.

* *Applicable official recommendations may vary regarding the interval between doses and the need for two or one doses of measles, mumps and rubella and of varicella-containing vaccines.*

Children from 9 to 10 months of age

In case an epidemiological situation requires vaccinating infants less than 11 months of age, the first dose of Priorix-Tetra can be given as from 9 months of age. A second dose of Priorix-Tetra should be given three months after the first dose (see section 5.1).

Method of administration

The vaccine is to be injected subcutaneously (SC) or intramuscularly (IM) in the deltoid region of the upper arm or in the higher anterolateral area of the thigh.

The vaccine should preferably be administered subcutaneously in subjects with thrombocytopenia or any coagulation disorder (see section 4.4).

For instructions on reconstitution of the medicinal product before administration, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1 or neomycin. A history of contact dermatitis to neomycin is not a contraindication. For hypersensitivity reactions to egg proteins, see section 4.4.

Current or recent immunosuppressive therapy (including high doses of corticosteroids). Priorix is not contraindicated in individuals who are receiving topical or low-dose parenteral corticosteroids (e.g. for asthma prophylaxis or replacement therapy) (see section 4.4).

Severe humoral or cellular (primary or acquired) immunodeficiency, e.g. severe combined immunodeficiency, agammaglobulinemia and AIDS or symptomatic HIV infection or an age-specific CD4+ T-lymphocyte percentage in children below 12 months: CD4+ <25 %; children between 12-35 months: CD4+ < 20 %; children between 36-59 months: CD4+ < 15 % (see section 4.4).

Pregnancy. Furthermore, pregnancy should be avoided for 1 month following vaccination (see section 4.6).

As with other vaccines, the administration of Priorix should be postponed in subjects suffering from acute severe febrile illness. The presence of a minor infection, such as a cold, should not result in the deferral of vaccination.

4.4 Special warnings and precautions for use

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following the administration of the vaccine.

Alcohol and other disinfecting agents must be allowed to evaporate from the skin before injection of the vaccine since they can inactivate the attenuated viruses in the vaccine.

Infants in their first year of life may not respond sufficiently to the components of the vaccine, due to the possible interference with maternal antibodies (see sections 4.2 and 5.1).

Due caution should be employed in administration of Priorix to individuals with Central Nervous System (CNS) disorder, susceptibility to febrile convulsions or family history of convulsions. Vaccinees with a history of febrile convulsions should be closely followed-up.

The measles and mumps components of the vaccine are produced in chick embryo cell culture and may therefore contain traces of egg protein. Persons with a history of anaphylactic, anaphylactoid, or other immediate reactions (e.g. generalised urticaria, swelling of the mouth and throat, difficulty in breathing, hypotension, or shock) subsequent to egg ingestion may be at an enhanced risk of immediate-type hypersensitivity reactions after vaccination, although these types of reactions have been shown to be very rare. Individuals who have experienced anaphylaxis after egg ingestion should be vaccinated with extreme caution, with adequate treatment for anaphylaxis on hand should such a reaction occur.

Limited protection against measles may be obtained by vaccination up to 72 hours after exposure to natural measles.

Syncope (fainting) can occur following, or even before, any vaccination especially in adolescents as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints.

As with any vaccine, a protective immune response may not be elicited in all vaccinees.

priorix should under no circumstances be administered intravascularly.

Thrombocytopenia

Cases of worsening of thrombocytopenia and cases of recurrence of thrombocytopenia in subjects who suffered thrombocytopenia after the first dose have been reported following vaccination with live measles, mumps and rubella vaccines. MMR-associated thrombocytopenia is rare and generally self-limited. In patients with existing thrombocytopenia or a history of thrombocytopenia after measles, mumps or rubella vaccination the risk-benefit of administering Priorix should be carefully evaluated. These patients should be vaccinated with caution and preferably using subcutaneous route.

Immunocompromised patients

Vaccination may be considered in patients with selected immune deficiencies where the benefits outweigh the risks (e.g. asymptomatic HIV subjects, IgG subclass deficiencies, congenital neutropenia, chronic granulomatous disease and complement deficiency diseases).

Immunocompromised patients who have no contraindication for this vaccination (see section 4.3) may not respond as well as immunocompetent subjects, therefore some of these patients may acquire measles, mumps or rubella in case of contact, despite appropriate vaccine administration. These patients should be monitored carefully for signs of measles, parotitis and rubella.

Due to the potential risk of decreased vaccine response and/or disseminated diseases, consideration should be given to the time interval between Priorix vaccination and immunosuppressive therapy (see section 4.3).

Transmission

Transmission of measles and mumps virus from vaccinees to susceptible contacts has never been documented. Pharyngeal excretion of the rubella and measles virus is known to occur about 7 to 28 days after vaccination with peak excretion around the 11th day. However, there is no evidence of transmission of these excreted vaccine viruses to susceptible contacts. Transmission of the rubella vaccine virus to infants via breast milk as well as transplacental transmission has been documented without any evidence of clinical disease.

Excipients with known effect

The vaccine contains 9 mg of sorbitol per dose.

The vaccine contains para-aminobenzoic acid. It may cause allergic reactions (possibly delayed), and exceptionally, bronchospasm.

The vaccine contains 334 micrograms of phenylalanine per dose. Phenylalanine may be harmful for individuals with phenylketonuria (PKU).

The vaccine contains 0.65 micrograms of polysorbate 80 per dose. Polysorbates may cause allergic reactions.

The vaccine contains 407 micrograms of proline per dose. Proline may be harmful for patients with hyperprolinaemia.

The vaccine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

The vaccine contains potassium, less than 1 mmol (39 mg) per dose, i.e. essentially 'potassium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Clinical studies have demonstrated that PRIORIX can be given simultaneously with any of the following monovalent or combination vaccines [including hexavalent vaccines (DTPa-HBV-IPV/Hib)]: diphtheria-tetanus-acellular pertussis vaccine (DTPa), reduced antigen diphtheria-tetanus-acellular pertussis vaccine (dTpa), *Haemophilus influenzae* type b vaccine (Hib), inactivated polio vaccine (IPV), hepatitis B vaccine (HBV), hepatitis A vaccine (HAV), meningococcal serogroup B vaccine (MenB), meningococcal serogroup C conjugate vaccine (MenC), meningococcal serogroups A, C, W-135 and Y conjugate vaccine (MenACWY), varicella zoster vaccine (VZV), oral polio vaccine (OPV) and pneumococcal conjugate vaccine in accordance with local recommendations.

Due to an increased risk of fever, tenderness at the injection site, change in eating habits and irritability when Bexsero was co-administered with a combined measles-mumps-rubella-varicella (MMR-V) vaccine, separate vaccination with PRIORIX can be considered when possible.

There are no data to support the use of PRIORIX with any other vaccines.

If PRIORIX is to be given at the same time as another injectable vaccine, the vaccines should always be administered at different injection sites.

If not given at the same time, an interval of at least one month is recommended between administration of PRIORIX and other live attenuated vaccines.

If tuberculin testing has to be done it should be carried out before or simultaneously with vaccination since it has been reported that combined measles, mumps and rubella vaccines may cause a temporary depression of tuberculin skin sensitivity. As this anergy may last up to a maximum of 6 weeks, tuberculin testing should not be performed within that period after vaccination to avoid false negative results.

In subjects who have received human gammaglobulins or a blood transfusion, vaccination should be delayed for three months or longer (up to 11 months) depending on the dose of human globulins administered because of the likelihood of vaccine failure due to passively acquired measles, mumps and rubella antibodies.

4.6 Fertility, pregnancy and lactationFertility

Priorix-Tetra has not been evaluated in fertility studies.

Pregnancy

Pregnant women should not be vaccinated with Priorix-Tetra.

Studies have not been conducted with Priorix-Tetra in pregnant women.

In a review of more than 3 500 susceptible women who were unknowingly in early stages of pregnancy when vaccinated with a rubella-containing vaccine, no cases of congenital rubella syndrome were reported. Subsequent post-marketing surveillance identified congenital rubella syndrome associated with a rubella vaccine strain (Wistar RA 27/3) following inadvertent vaccination of a pregnant woman with a measles, mumps and rubella vaccine.

Foetal damage has not been documented when measles, mumps or varicella vaccines have been given to pregnant women. Pregnancy should be avoided for 1 month following vaccination. Women who intend to become pregnant should be advised to delay.

Breast-feeding

Adequate human data on the use of Priorix-Tetra during lactation are not available.

4.7 Effects on ability to drive and use machines

No studies on the effects of Priorix-Tetra on the ability to drive and use machines have been performed. Priorix-Tetra is expected to have no or negligible influence on the ability to drive and use machines.

4.8 Undesirable effects

Summary of the safety profile

The safety profile presented below is based on data from clinical trials in which more than 6,700 doses of Priorix-Tetra were administered subcutaneously to more than 4,000 children from 9 to 27 months of age. Events were recorded for up to 42 days after vaccination.

The most common adverse reactions following Priorix-Tetra administration were pain and redness at the injection site as well as fever $\geq 38^{\circ}\text{C}$ (rectal) or $\geq 37.5^{\circ}\text{C}$ (axillary/oral).

Tabulated list of adverse reactions

Adverse reactions reported are listed according to the following frequency:

Very common: ($\geq 1/10$)

Common: ($\geq 1/100$ to $< 1/10$)

Uncommon: ($\geq 1/1\,000$ to $< 1/100$)

Rare: ($\geq 1/10\,000$ to $< 1/1\,000$)

Very rare: ($< 1/10\,000$)

Clinical trial data

System Organ Class	Frequency	Adverse reactions
Infections and infestations	Uncommon	upper respiratory tract infection
	Rare	otitis media
Blood and lymphatic system disorders	Uncommon	lymphadenopathy
Metabolism and nutrition disorders	Uncommon	anorexia
Psychiatric disorders	Common	irritability
	Uncommon	crying, nervousness, insomnia
Nervous system disorders	Rare	febrile convulsions*
Respiratory, thoracic and mediastinal disorders	Uncommon	rhinitis
	Rare	cough, bronchitis
Gastrointestinal disorders	Uncommon	parotid gland enlargement, diarrhoea, vomiting
Skin and subcutaneous tissue disorders	Common	rash
General disorders and administration site conditions	Very common	pain and redness at the injection site, fever (rectal $\geq 38^{\circ}\text{C}$ - $\leq 39.5^{\circ}\text{C}$; axillary/oral: $\geq 37.5^{\circ}\text{C}$ - $\leq 39^{\circ}\text{C}$)**
	Common	swelling at the injection site, fever (rectal $> 39.5^{\circ}\text{C}$; axillary/oral $> 39^{\circ}\text{C}$)**
	Uncommon	lethargy, malaise, fatigue

*The risk of febrile convulsions following the first dose vaccination of children aged 9 to 30 months with Priorix-Tetra compared with MMR or simultaneous, but separate MMR and varicella vaccination was assessed in a retrospective database analysis.

The study included 82 656 children immunized with MMRV, 149 259 with MMR and 39 203 with separate MMR and varicella vaccines.

Depending on the case definition used to identify febrile convulsions in the main risk period 5 to 12 days following the first dose, incidences of febrile convulsions were 2.18 (95% CI: 1.38; 3.45) or 6.19 (95% CI: 4.71; 8.13) per 10 000 subjects for the MMRV group and 0.49 (95% CI: 0.19; 1.25) or 2.55 (95% CI: 1.67; 3.89) per 10 000 subjects for the matched control cohorts. These data suggest one additional case of febrile convulsion per 5 882 or 2 747 subjects vaccinated with Priorix-Tetra compared to matched control cohorts who received MMR or simultaneous, but separate MMR and varicella vaccination (attributable risk of 1.70 (95% CI: -1.86; 3.46) and 3.64 (95% CI: -6.11; 8.30) per 10 000 subjects, respectively) – see section 5.1.

**Following the administration of the first dose of the combined measles-mumps-rubella-varicella vaccine, higher incidences of fever (approximately 1.5 fold) were observed when compared to the concomitant administration of measles-mumps-rubella and varicella vaccines at separate injection sites.

No clinical studies with Priorix-Tetra (MMRV) have been conducted in subjects > 6 years of age. The safety profile of Priorix-Tetra in subjects > 6 years of age is extrapolated from the available data with GlaxoSmithKline's MMR vaccine (Priorix) and monovalent Oka varicella vaccine (Varilrix). The spectrum of adverse reactions such as fever, rash, injection site pain, injection site swelling and injection site redness in subjects > 6 years of age who received Priorix or Varilrix was comparable to that observed in children < 6 years of age who received Priorix-Tetra. In these clinical studies evidence has been obtained to conclude that the second dose of the MMR vaccine is better tolerated in terms of fever than the first dose, whereas reactogenicity of the varicella vaccine tends to remain similar irrespective of the dose applied. Injection site swelling is "commonly" reported in children who received Priorix-Tetra, while it is "very commonly" reported in Varilrix studies in adolescents and adults.

Post-marketing surveillance data

The following additional adverse reactions have been identified in rare occasions during post-marketing surveillance. Because they are reported voluntarily from a population of unknown size, a true estimate of frequency cannot be provided.

System Organ Class	Adverse reactions
Infections and infestations	meningitis, herpes zoster*, measles-like syndrome, mumps-like syndrome (including orchitis, epididymitis and parotitis)
Blood and lymphatic system disorders	thrombocytopenia, thrombocytopenic purpura
Immune system disorders	allergic reactions (including anaphylactic and anaphylactoid reactions)
Nervous system disorders	encephalitis, cerebellitis, cerebrovascular accident, Guillain Barré syndrome, transverse myelitis, peripheral neuritis, cerebellitis like symptoms (including transient gait disturbance and transient ataxia)
Vascular disorders	vasculitis
Skin and subcutaneous tissue disorders	erythema multiforme, varicella-like rash
Musculoskeletal and connective tissue disorders	arthralgia, arthritis

*This adverse drug reaction reported after vaccination is also a consequence of wild-type varicella infection. There is no indication of an increased risk of herpes zoster occurrence following vaccination compared with wild-type disease.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRC Pharmacovigilance, Website: www.hpra.ie.

4.9 Overdose

No case of overdose has been reported.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmaco-therapeutic group: Vaccines, Viral vaccine, ATC code J07BD54

Efficacy

The efficacy of GlaxoSmithKline (GSK)'s monovalent Oka varicella (Varilrix) and Priorix-Tetra vaccines in preventing varicella disease has been evaluated in a large randomised multicountry clinical trial, which included GSK combined measles-mumps-rubella vaccine (Priorix) as active control. The trial has been conducted in Europe where no routine varicella vaccination was implemented at that time. Children aged 12-22 months received two doses of Priorix-Tetra six weeks apart or one dose of Varilrix. Vaccine efficacy against epidemiologically confirmed or PCR (Polymerase Chain Reaction) confirmed varicella of any severity (defined using a prespecified scale) and against moderate or severe confirmed varicella was demonstrated after a primary follow-up period of 2 years (median duration 3.2 years). Persistent efficacy was observed in the same study during the long term follow-up periods of 6 years (median duration 6.4 years) and 10 years (median duration 9.8 years). The data are presented in the table below.

Group	Timing	Efficacy against confirmed varicella of any severity	Efficacy against moderate or severe confirmed varicella
Priorix-Tetra (2 doses) N = 2,489	Year 2	94.9% (97.5% CI: 92.4;96.6)	99.5% (97.5% CI: 97.5;99.9)
	Year 6 ⁽¹⁾	95.0% (95% CI: 93.6;96.2)	99.0% (95% CI: 97.7;99.6)
	Year 10 ⁽¹⁾	95.4% (95% CI: 94.0;96.4)	99.1% (95% CI: 97.9;99.6)
Varilrix (1 dose) N = 2,487	Year 2	65.4 % (97.5% CI: 57.2;72.1)	90.7% (97.5% CI: 85.9;93.9)
	Year 6 ⁽¹⁾	67.0% (95% CI: 61.8;71.4)	90.3% (95% CI: 86.9;92.8)
	Year 10 ⁽¹⁾	67.2% (95% CI: 62.3;71.5)	89.5% (95% CI: 86.1;92.1)

N = number of subjects enrolled and vaccinated

(1) descriptive analysis

Effectiveness

Effectiveness data suggest a higher level of protection and a decrease in breakthrough varicella following two doses of varicella-containing vaccine than following one dose.

The effectiveness of two doses of Priorix-Tetra during varicella outbreaks in day care centres in Germany, where routine varicella vaccination is recommended for children as of 11 months of age, was 91% (95% CI: 65;98) against any disease and 94% (95% CI: 54;99) against moderate disease.

The effectiveness of one dose of Varilrix was estimated in different settings (outbreaks, case-control and database studies) and ranged from 20%-92% against any varicella disease and from 86%-100% against moderate or severe disease.

Immune response

Several clinical studies evaluated the immune response elicited by Priorix-Tetra administered subcutaneously. Anti-measles, anti-mumps and anti-rubella antibody titres were determined using commercially available enzyme-linked immunosorbent assays (ELISA). In addition, anti-mumps antibodies were titrated using a plaque-reduction neutralisation assay. These serological parameters are widely accepted as surrogate markers for immune protection. A modified commercial indirect immunofluorescence assay (IFA, meanwhile discontinued) and a commercial ELISA were used to compare the immune response against varicella elicited by Priorix-Tetra to that observed with GSK varicella vaccine.

In three clinical trials conducted in Europe (Austria, Finland, Germany, Greece, Poland) approximately 2,000 previously unvaccinated children from 11 to 23 months of age received 2 doses of Priorix-Tetra with an interval between doses of 6 weeks. Seroconversion rates and geometric mean antibody concentrations/titres (GMC/GMT) are summarized in the table below.

Antibody Test (cut-off)	Post dose 1		Post dose 2	
	Seroconversion rates (95 % CI)	GMC/GMT (95 % CI)	Seroconversion rates (95 % CI)	GMC/GMT (95 % CI)
Measles ELISA (150 mIU/ml)	96.4% (CI: 95.5;97.2)	3184.5 (CI: 3046.5;3328.7)	99.1% (CI: 98.6;99.5)	4828.6 (CI: 4644.3;5020.1)
Mumps				

ELISA (231 U/ml)	91.3% (CI: 90.0;92.5)	976.7 (CI: 934.8;1020.5)	98.8% (CI: 98.2;99.2)	1564.4 (CI: 1514.6;1615.8)
Neutralisation (1:28)	95.4% (CI: 94.3;96.3)	147.0 (CI: 138.6;155.8)	99.4% (CI: 98.9;99.7)	478.4 (CI: 455.1;503.0)
Rubella ELISA (4 IU/ml)	99.7% (CI: 99.4;99.9)	62.2 (CI: 60.0;64.5)	99.9% (CI: 99.6;100)	119.7 (CI: 116.4;123.1)
Varicella IFA (1:4)	97.2% (CI: 96.3;97.9)	97.5 (CI: 92.2;103.1)	99.8% (CI: 99.5;100)	2587.8 (CI: 2454.0;2728.9)
ELISA (50 mIU/ml)	89.4% (CI: 87.8;90.8)	112.0 (CI: 93.5;134.0)	99.2% (CI: 98.5;99.6)	2403.9 (CI: 1962.4;2944.6)

Seroconversion rates and geometric mean antibody concentrations/titres were similar to those observed after separate vaccination with Varilrix and Priorix.

In infants vaccinated at 11 months of age, the proportion of infants with protective measles titers (i.e., ≥ 150 mIU/mL) after the first dose is 91-92%, lower than the proportion observed when the first dose is administered since the age of 12 months.

The second dose of Priorix-Tetra induced an increase in seroconversion rates and/or antibody levels for the measles, mumps and rubella vaccine components. Therefore, to avoid infection during the interval between doses it is preferred that the second dose be administered within three months following the first dose.

Data suggest a higher efficacy and a decrease in breakthrough varicella following two doses of vaccine with respect to one dose. This correlates with an increase in anti-varicella antibodies elicited by the second dose, which suggests that the second dose of varicella antigen acts as a booster.

The immune response of Priorix-Tetra administered as a second dose of MMR vaccine in children 24 months to 6 years of age was evaluated in 2 clinical studies. Children were previously primed with respectively an MMR vaccine or with an MMR vaccine co-administered with a live attenuated varicella vaccine. Seropositivity rates for anti-varicella antibodies were 98.1% (IFA) in children previously vaccinated with MMR and 100% in children previously vaccinated with an MMR vaccine co-administered with a live attenuated varicella vaccine. Seropositivity rates were 100% for anti-measles, mumps, rubella antibodies in both studies.

Immune response in subjects > 6 years of age

No clinical studies to evaluate the immunogenicity of Priorix-Tetra in subjects > 6 years of age have been conducted. The immunogenicity of Priorix-Tetra in subjects > 6 years is extrapolated from available data with Priorix and Varilrix.

Immune response in children aged 9 to 10 months

A clinical trial conducted in Asia (Singapore) enrolled 300 healthy children 9 to 10 months of age at the time of first vaccine dose. Of these, 153 subjects received 2 doses of Priorix-Tetra with an interval between doses of 3 months and 147 subjects received Priorix and Varilrix. Seroconversion rates and geometric mean antibody concentrations/titres were similar to those observed after separate vaccination with Varilrix and Priorix. Seroconversion rates after a first dose of Priorix-Tetra were comparable for all antigens except measles to those seen in 12-24 months old children in other clinical studies. The seroconversion rate reported for measles in infants 9 to 10 months of age following 1 dose of Priorix-Tetra was 93.3% (95% CI: 87.6;96.9). Infants in their first year of life may not respond sufficiently to the components of the vaccine due to the possible interference with maternal antibodies. Therefore a second dose of Priorix-Tetra should be given three months after the first dose.

The immunogenicity and safety of Priorix-Tetra administered intramuscularly was evaluated in one comparative study conducted in 328 children aged 11-21 months who received 2 doses of Priorix-Tetra either by intramuscular or subcutaneous route. The study demonstrated comparable immunogenicity and safety profiles for both administration routes.

Persistence of measles, mumps and rubella immune response

In a clinical trial in which children aged 12-22 months received two doses of Priorix-Tetra (N = 2,489), the seropositivity rates for anti-measles, mumps and rubella antibodies, in terms of subjects with an antibody concentration equal to or above defined threshold, observed after follow-up periods of 2, 6 and 10 years are presented in the table below:

Timing	Antibody Test (cut-off)		
	Measles ELISA (150 mIU/ml)	Mumps ELISA (231 U/ml)	Rubella ELISA (4 IU/ml)
Year 2	99.1%	90.5%	100%
Year 6	99.0%	90.5%	99.8%
Year 10	98.5%	90.0%	97.7%

ELISA: Enzyme Linked Immuno Sorbent Assay

As robust efficacy data against varicella disease up to 10 years are provided above (see subsection "Efficacy") and since there is no threshold for protection against varicella disease established with the obtained immunological data, anti-varicella antibody persistence data are not provided.

Post-Marketing Observational Safety Surveillance Study

The risk of febrile convulsions following the first dose of Priorix-Tetra was assessed in a retrospective database analysis in children aged 9 to 30 months (see section 4.8).

5.2 Pharmacokinetic properties

Not applicable.

5.3 Preclinical safety data

A repeated dose toxicity study in animals did not reveal any local or systemic toxicity of the vaccine.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Powder:
 Amino acids (containing phenylalanine and proline)
 Lactose anhydrous
 Mannitol (E 421)
 Sorbitol (E 420)
 Medium 199 (containing phenylalanine, para-aminobenzoic acid, polysorbate 80 (E 433), proline, sodium and potassium)

Solvent:
 Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

18 months.
 After reconstitution, use the vaccine promptly. If this is not possible, it must be stored at 2°C – 8°C and used within 24 hours of reconstitution.

6.4 Special precautions for storage

Store and transport refrigerated (2°C – 8°C).

Do not freeze.

Store in the original packaging in order to protect from light.

For storage conditions after reconstitution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

Priorix is presented as:

- Powder for 1 dose in a vial (type I glass) with a stopper (butyl rubber).
- Solvent for 1 dose in pre-filled syringe (type I glass) with a plunger stopper (butyl rubber) and a rubber tip cap.

The tip cap and rubber plunger stopper of the pre-filled syringe and the stopper of the vial are made with synthetic rubber.

Pack sizes of 1 or 10, with 2 separate needles or without needles.

Not all pack sizes may be marketed.


6.6 Special precautions for disposal and other handling

The solvent and the reconstituted vaccine should be inspected visually for any foreign particulate matter and/or variation of physical aspects prior to reconstitution or administration. In the event of either being observed, do not use the solvent or the reconstituted vaccine.

Due to minor variation of its pH, the reconstituted vaccine may vary in colour from clear peach to fuchsia pink without deterioration of the vaccine potency.


The vaccine must be reconstituted by adding the entire contents of the pre-filled syringe of solvent to the vial containing the powder using a suitable needle (21G to 25G).

To attach the needle to the syringe, carefully read the instructions.



Hold the syringe by the barrel, not by the plunger or by the Luer Lock Adaptor (LLA).

Unscrew the syringe cap by twisting it anticlockwise.



To attach the needle, gently connect the hub to the LLA and rotate a quarter turn clockwise until you feel it lock.

Maintain the needle in the axis of the syringe. Failure to do this may cause the LLA to become distorted and leak.

During assembly of the syringe, if the LLA comes off, a new vaccine dose (new syringe and vial) should be used.

Reconstitute the vaccine as described below.

Do not pull the syringe plunger out of the barrel. If it happens, do not administer the vaccine.

1. Add the solvent to the powder. Shake well until the powder is completely dissolved in the solvent. After reconstitution, use the vaccine promptly.
2. Withdraw the entire contents of the vial.
3. A new needle should be used to administer the vaccine. Unscrew the needle from the syringe and attach the injection needle by repeating steps above.

Contact with disinfectants should be avoided (see section 4.4).

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

GlaxoSmithKline (Ireland) Limited
12 Riverwalk
Citywest Business Campus
Dublin 24
Ireland

8 MARKETING AUTHORISATION NUMBER

PA1077/117/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

The date of first authorisation: 10th August 2007
Date of last renewal: 26th July 2011

10 DATE OF REVISION OF THE TEXT

May 2026