

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Remifentanil Noridem 5 mg powder for concentrate for solution for injection or infusion

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains remifentanil hydrochloride equivalent to 5 mg of remifentanil.
After reconstitution as directed, each ml of solution contains 1 mg of remifentanil.
For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Powder for concentrate for solution for injection or infusion.
White to off white, lyophilised powder.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Remifentanil Noridem is indicated as an analgesic agent for use during induction and/or maintenance of general anaesthesia.

Remifentanil Noridem is indicated for provision of analgesia in mechanically ventilated intensive care patients 18 years of age and over.

4.2 Posology and method of administration

Remifentanil Noridem should be administered only in a setting fully equipped for the monitoring and support of respiratory and cardiovascular function, and by persons specifically trained in the use of anaesthetic drugs and the recognition and management of the expected adverse effects of potent opioids, including respiratory and cardiac resuscitation. Such training must include the establishment and maintenance of a patent airway and assisted ventilation.

Continuous infusions of Remifentanil Noridem must be administered by a calibrated infusion device into a fast flowing IV line or via a dedicated IV line. This infusion line should be connected at, or close to, the venous cannula and primed, to minimise the potential dead space (see section 6.6).

Remifentanil Noridem may also be given by target-controlled infusion (TCI) with an approved infusion device incorporating the Minto pharmacokinetic model with covariates for age and lean body mass (LBM) (Anesthesiology 1997; 86: 10-23).

Care should be taken to avoid obstruction or disconnection of infusion lines and to adequately clear the lines to remove residual Remifentanil Noridem after use (see section 4.4).

Remifentanil Noridem is for intravenous use only and must not be administered by epidural or intrathecal injection (see section 4.3).

Dilution

Remifentanil Noridem may be further diluted after reconstitution. For instructions on dilution of the medicinal product before administration, see section 6.6.

For manually-controlled infusion Remifentanil Noridem can be diluted to concentrations of 20 to 250 micrograms/mL (50 micrograms/mL is the recommended dilution for adults and 20 to 25 micrograms/mL for paediatric patients aged 1 year and over).

For TCI the recommended dilution of Remifentanil Noridem is 20 to 50 micrograms/mL.

General Anaesthesia

The administration of Remifentanil Noridem must be individualised based on the patient's response.

Adults**Administration by Manually-Controlled Infusion**

Table 1 summarises the starting injection/infusion rates and dose range:

Table 1. Dosing Guidelines for Adults

INDICATION	BOLUS INJECTION (micrograms/kg)	CONTINUOUS INFUSION (micrograms/kg/min)	
		Starting Rate	Range
Induction of anaesthesia	1 (give over not less than 30 seconds)	0.5 to 1	–
Maintenance of anaesthesia in ventilated patients			
• Nitrous oxide (66%)	0.5 to 1	0.4	0.1 to 2
• Isoflurane (starting dose 0.5 MAC)	0.5 to 1	0.25	0.05 to 2
• Propofol (starting dose 100 micrograms/kg/min)	0.5 to 1	0.25	0.05 to 2

When given by slow bolus injection Remifentanyl Noridem should be administered over not less than 30 seconds.

At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended above to avoid an increase of haemodynamic effects such as hypotension and bradycardia (see this section *Concomitant medication*).

No data are available for dosage recommendations for simultaneous use of other hypnotics other than those listed in Table 1 with remifentanyl.

Induction of anaesthesia: Remifentanyl Noridem should be administered with a standard dose of hypnotic agent, such as propofol, thiopental, or isoflurane, for the induction of anaesthesia. Remifentanyl Noridem can be administered at an infusion rate of 0.5 to 1 micrograms/kg/min, with or without an initial slow bolus injection of 1 microgram/kg given over not less than 30 seconds. If endotracheal intubation is to occur more than 8 to 10 minutes after the start of the infusion of Remifentanyl Noridem, then a bolus injection is not necessary.

Maintenance of anaesthesia in ventilated patients: After endotracheal intubation, the infusion rate of Remifentanyl Noridem should be decreased, according to anaesthetic technique, as indicated in Table 1. Due to the fast onset and short duration of action of Remifentanyl Noridem, the rate of administration during anaesthesia can be titrated upward in 25% to 100% increments or downward in 25% to 50% decrements, every 2 to 5 minutes to attain the desired level of μ -opioid response. In response to light anaesthesia, supplemental slow bolus injections may be administered every 2 to 5 minutes.

Anaesthesia in spontaneously breathing anaesthetised patients with a secured airway (e.g. laryngeal mask anaesthesia):In spontaneously breathing anaesthetised patients with a secured airway respiratory depression is likely to occur. Special care is needed to adjust the dose to the patient requirements and ventilatory support may be required. The recommended starting infusion rate for supplemental analgesia in spontaneously breathing anaesthetised patients is 0.04 micrograms/kg/min with titration to effect. A range of infusion rates from 0.025 to 0.1 micrograms/kg/min has been studied. Bolus injections are not recommended in spontaneously breathing anaesthetised patients.

Remifentanyl Noridem should not be used as an analgesic in procedures where patients remain conscious or do not receive any airway support during the procedure.

Concomitant medication: Remifentanyl Noridem decreases the amounts or doses of inhaled anaesthetics, hypnotics and benzodiazepines required for anaesthesia (see section 4.5).

Doses of the following agents used in anaesthesia: isoflurane, thiopentone, propofol and temazepam have been reduced by up to 75% when used concurrently with remifentanyl.

Guidelines for discontinuation/continuation into the immediate post-operative period: Due to the very rapid offset of action of Remifentanyl Noridem no residual opioid activity will be present within 5 to 10 minutes after discontinuation. For those patients undergoing surgical procedures where post-operative pain is anticipated, analgesics should be administered prior to discontinuation of Remifentanyl Noridem. Sufficient time must be allowed to reach the maximum effect of the longer acting analgesic. The choice of analgesic should be appropriate for the patient's surgical procedure and the level of post-operative care.

In the event that longer acting analgesia has not been established prior to the end of surgery, Remifentanyl Noridem may need to be continued to maintain analgesia during the immediate post-operative period until longer acting analgesia has reached its maximum effect.

Guidance on use in mechanically ventilated intensive care patients is provided in this section *Use in intensive care*.

In patients who are breathing spontaneously, the infusion rate of Remifentanyl Noridem should initially be decreased to a rate of 0.1 micrograms/kg/min. The infusion rate may then be increased or decreased by not greater than 0.025 micrograms/kg/min every five minutes, to balance the patient's level of analgesia and respiratory rate. Remifentanyl Noridem should only be used in a setting fully equipped for the monitoring and support of respiratory and cardiovascular function, under the close supervision of persons specifically trained in the recognition and management of the respiratory effects of potent opioids.

The use of bolus injections of Remifentanyl Noridem to treat pain during the post-operative period is not recommended in patients who are breathing spontaneously.

Administration by Target-Controlled Infusion

Induction and maintenance of anaesthesia in ventilated patients: Remifentanyl Noridem TCI should be used in association with an intravenous or inhalational hypnotic agent during the induction and maintenance of anaesthesia in ventilated adult patients (see Table 1, in this section *General anaesthesia*).

In association with these agents, adequate analgesia for induction of anaesthesia and surgery can generally be achieved with target blood remifentanyl concentrations ranging from 3 to 8 nanograms/mL. Remifentanyl Noridem should be titrated to individual patient response. For particularly stimulating surgical procedures target blood concentrations up to 15 nanograms/mL may be required.

At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended above to avoid an increase of haemodynamic effects such as hypotension and bradycardia (see Table 1 and this section *Concomitant medication*).

For information on blood remifentanyl concentrations achieved with manually-controlled infusion see section 6.6, Table 11.

As there are insufficient data, the administration of Remifentanyl Noridem by TCI for spontaneous ventilation anaesthesia is not recommended.

Guidelines for discontinuation/continuation into the immediate post-operative period: At the end of surgery when the TCI infusion is stopped or the target concentration reduced, spontaneous respiration is likely to return at calculated remifentanyl concentrations in the region of 1 to 2 nanograms/mL. As with manually-controlled infusion, post-operative analgesia should be established before the end of surgery with longer acting analgesics (see this section *Administration by manually-controlled infusion – Guidelines for discontinuation*).

As there are insufficient data, the administration of Remifentanyl Noridem by TCI for the management of post-operative analgesia is not recommended.

Paediatric patients (1 to 12 years of age)

Co-administration of remifentanyl and an intravenous anaesthetic agent for induction of anaesthesia has not been studied in detail and is therefore not recommended.

Remifentanyl Noridem TCI has not been studied in paediatric patients and therefore administration of Remifentanyl Noridem by TCI is not recommended in these patients. The following doses of Remifentanyl Noridem are recommended for maintenance of anaesthesia:

Table 2. Dosing Guidelines for Paediatric Patients (1 to 12 years of age)

CONCOMITANT ANAESTHETIC AGENT*	BOLUS INJECTION (micrograms/kg)	CONTINUOUS INFUSION (micrograms/kg/min)	
		Starting Rate	Typical Maintenance Rates
Halothane (starting dose 0.3 MAC)	1	0.25	0.05 to 1.3
Sevoflurane (starting dose 0.3 MAC)	1	0.25	0.05 to 0.9

Isoflurane (starting dose 0.5 MAC)	1	0.25	0.06 to 0.9
------------------------------------	---	------	-------------

*co-administered with nitrous oxide/oxygen in a ratio of 2:1

When given by bolus injection Remifentanyl Noridem should be administered over not less than 30 seconds. Surgery should not commence until at least 5 minutes after the start of the Remifentanyl Noridem infusion, if a simultaneous bolus dose has not been given. For sole administration of nitrous oxide (70%) with Remifentanyl Noridem, typical maintenance infusion rates should be between 0.4 and 3 micrograms/kg/min, and although not specifically studied, adult data suggest that 0.4 micrograms/kg/min is an appropriate starting rate. Paediatric patients should be monitored and the dose titrated to the depth of analgesia appropriate for the surgical procedure.

Concomitant medication: At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane, halothane and sevoflurane should be administered as recommended above to avoid an increase of haemodynamic effects such as hypotension and bradycardia. No data are available for dosage recommendations for simultaneous use of other hypnotics other than those listed in the table with remifentanyl (see this section *Adults-Concomitant medication*).

Guidelines for patient management in the immediate post-operative period/ Establishment of alternative analgesia prior to discontinuation of Remifentanyl Noridem: Due to the very rapid offset of action of Remifentanyl Noridem, no residual activity will be present within 5 to 10 minutes after discontinuation. For those patients undergoing surgical procedures where post-operative pain is anticipated, analgesics should be administered prior to discontinuation of Remifentanyl Noridem. Sufficient time must be allowed to reach the therapeutic effect of the longer acting analgesic. The choice of agent(s), the dose and the time of administration should be planned in advance and individually tailored to be appropriate for the patient's surgical procedure and the level of post-operative care anticipated (see section 4.4).

Neonates/infants (aged less than 1 year)

There is limited clinical trial experience of remifentanyl in neonates and infants (aged under 1 year old; see section 5.1). The pharmacokinetic profile of remifentanyl in neonates/infants (aged less than 1 year) is comparable to that seen in adults after correction for body weight differences (see section 5.2). However, because there are insufficient clinical data, the administration of remifentanyl is not recommended for this age group.

Use for Total Intravenous anaesthesia (TIVA): There is limited clinical trial experience of remifentanyl for TIVA in infants (see section 5.1). However, there are insufficient clinical data to make dosage recommendations.

Cardiac anaesthesia

Administration by Manually-Controlled Infusion

Table 3. Dosing Guidelines for Cardiac Anaesthesia

INDICATION	BOLUS INJECTION (micrograms/kg)	CONTINUOUS INFUSION (micrograms/kg/min)	
		Starting Rate	Typical Infusion Rates
Intubation	Not recommended	1	–
Maintenance of anaesthesia			
Isoflurane (starting dose 0.4 MAC)	0.5 to 1	1	0.003 to 4
Propofol (starting dose 50 micrograms/kg/min)	0.5 to 1	1	0.01 to 4.3
Continuation of post-operative analgesia, prior to extubation	Not recommended	1	0 to 1

Induction period of anaesthesia: After administration of hypnotic to achieve loss of consciousness, Remifentanyl Noridem should be administered at an initial infusion rate of 1 microgram/kg/min. The use of bolus injections of Remifentanyl Noridem during induction in cardiac surgical patients is not recommended. Endotracheal intubation should not occur until at least 5 minutes after the start of the infusion.

Maintenance period of anaesthesia: After endotracheal intubation the infusion rate of Remifentanyl Noridem should be titrated according to patient need. Supplemental slow bolus doses may also be given as required. High-risk cardiac patients, such as those with poor ventricular function or undergoing valve surgery, should be administered a maximum bolus dose of 0.5 micrograms/kg. These dosing recommendations also apply during hypothermic cardiopulmonary bypass (see section 5.2 – *Cardiac anaesthesia*).

Concomitant medication: At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended above to avoid

an increase of haemodynamic effects such as hypotension and bradycardia. No data are available for dosage recommendations for simultaneous use of other hypnotics other than those listed in the table with remifentanyl (see this section *Adults - Concomitant medication*).

Guidelines for post-operative patient management

Continuation of Remifentanyl Noridem post-operatively to provide analgesia prior to weaning for extubation: It is recommended that the infusion of Remifentanyl Noridem should be maintained at the final intra-operative rate during transfer of patients to the post-operative care area. Upon arrival into this area, the patient's level of analgesia and sedation should be closely monitored and the Remifentanyl Noridem infusion rate adjusted to meet the individual patient's requirements (see this section *Use in intensive care*, for further information on management of intensive care patients).

Establishment of alternative analgesia prior to discontinuation of Remifentanyl Noridem: Due to the very rapid offset of action of Remifentanyl Noridem, no residual opioid activity will be present within 5 to 10 minutes after discontinuation. Prior to discontinuation of Remifentanyl Noridem, patients must be given alternative analgesic and sedative agents at a sufficient time in advance to allow the therapeutic effects of these agents to become established. It is therefore recommended that the choice of agent(s), the dose and the time of administration are planned, before weaning the patient from the ventilator.

Guidelines for discontinuation of Remifentanyl Noridem: Due to the very rapid offset of action of Remifentanyl Noridem, hypertension, shivering and aches have been reported in cardiac patients immediately following discontinuation of Remifentanyl Noridem (see section 4.8). To minimise the risk of these occurring, adequate alternative analgesia must be established (as described above), before the Remifentanyl Noridem infusion is discontinued. The infusion rate should be reduced by 25% decrements in at least 10-minute intervals until the infusion is discontinued.

During weaning from the ventilator the Remifentanyl Noridem infusion should not be increased and only down titration should occur, supplemented as required with alternative analgesics. Haemodynamic changes such as hypertension and tachycardia should be treated with alternative agents as appropriate.

When other opioid agents are administered as part of the regimen for transition to alternative analgesia, the patient must be carefully monitored. The benefit of providing adequate post-operative analgesia must always be balanced against the potential risk of respiratory depression with these agents.

Administration by Target-Controlled Infusion

Induction and maintenance of anaesthesia: Remifentanyl Noridem TCI should be used in association with an intravenous or inhalational hypnotic agent during the induction and maintenance of anaesthesia in ventilated adult patients (see Table 3). In association with these agents, adequate analgesia for cardiac surgery is generally achieved at the higher end of the range of target blood remifentanyl concentrations used for general surgical procedures. Following titration of remifentanyl to individual patient response, blood concentrations as high as 20 nanograms/mL have been used in clinical studies. At the doses recommended above, remifentanyl significantly reduces the amount of hypnotic agent required to maintain anaesthesia. Therefore, isoflurane and propofol should be administered as recommended above to avoid an increase of haemodynamic effects such as hypotension and bradycardia (see Table 3 and this section *Concomitant medication*).

For information on blood remifentanyl concentrations achieved with manually-controlled infusion see section 6.6, Table 11.

Guidelines for discontinuation/continuation into the immediate post-operative period: At the end of surgery when the TCI infusion is stopped or the target concentration reduced, spontaneous respiration is likely to return at calculated remifentanyl concentrations in the region of 1 to 2 nanograms/mL. As with manually-controlled infusion, post-operative analgesia should be established before the end of surgery with longer acting analgesics (see this section *Administration by manually-controlled infusion – Guidelines for discontinuation*).

As there are insufficient data, the administration of Remifentanyl Noridem by TCI for the management of post-operative analgesia is not recommended.

Paediatric patients (1 to 12 years of age)

There are insufficient data to make a dosage recommendation for use during cardiac surgery.

Use in intensive care

Adults

Remifentanyl Noridem can be used for the provision of analgesia in mechanically ventilated intensive care patients. Sedative agents should be added as appropriate.

The safety and efficacy from well-controlled clinical trials of remifentanyl in mechanically ventilated intensive care patients has been established for durations up to 3 days (see this section *Renally-impaired intensive care patients* and section 5.2). Therefore, the use of Remifentanyl Noridem is not recommended for a duration of treatment greater than 3 days.

Remifentanyl TCI has not been studied in intensive care patients and therefore administration of Remifentanyl Noridem by TCI is not recommended in these patients.

In adults, it is recommended that Remifentanyl Noridem is initiated at an infusion rate of 0.1 micrograms/kg/min (6 micrograms/kg/h) to 0.15 micrograms/kg/min (9 micrograms/kg/h). The infusion rate should be titrated in increments of 0.025 micrograms/kg/min (1.5 micrograms/kg/h) to achieve the desired level of analgesia. A period of at least 5 minutes should be allowed between dose adjustments. The patient should be regularly assessed and the Remifentanyl Noridem infusion rate adjusted accordingly. If an infusion rate of 0.2 micrograms/kg/min (12 micrograms/kg/h) is reached and sedation is required, it is recommended that dosing with an appropriate sedative agent is initiated (see below). The dose of sedative agent should be titrated to obtain the desired level of sedation. Further increases to the Remifentanyl Noridem infusion rate in increments of 0.025 micrograms/kg/min (1.5 micrograms/kg/h) may be made if additional analgesia is required.

Table 4 summarises the starting infusion rates and typical dose range for provision of analgesia in individual patients:

Table 4. Dosing Guidelines for use of Remifentanyl Noridem within the Intensive Care Setting

CONTINUOUS INFUSION micrograms/kg/min (micrograms/kg/h)	
Starting Rate	Range
0.1 (6) to 0.15 (9)	0.006 (0.38) to 0.74 (44.6)

Bolus doses of Remifentanyl Noridem are not recommended in the intensive care setting.

The use of Remifentanyl Noridem will reduce the dosage requirement of any concomitant sedative agents. Typical starting doses for sedative agents, if required, are given in Table 5.

Table 5. Recommended starting dose of sedative agents, if required:

Sedative Agents	Bolus (mg/kg)	Infusion (mg/kg/h)
Propofol	Up to 0.5	0.5
Midazolam	Up to 0.03	0.03

To allow separate titration of the respective agents, sedative agents should not be prepared as one mixture in the same infusion bag.

Additional analgesia for ventilated patients undergoing stimulating procedures: An increase in the existing Remifentanyl Noridem infusion rate may be required to provide additional analgesic cover for ventilated patients undergoing stimulating and/or painful procedures such as endotracheal suctioning, wound dressing and physiotherapy. It is recommended that a Remifentanyl Noridem infusion rate of at least 0.1 micrograms/kg/min (6 micrograms/kg/h) should be maintained for at least 5 minutes prior to the start of the stimulating procedure. Further dose adjustments may be made every 2 to 5 minutes in increments of 25% to 50% in anticipation of, or in response to, additional requirement for analgesia. A mean infusion rate of 0.25 micrograms/kg/min (15 micrograms/kg/h), maximum 0.74 micrograms/kg/min (45 micrograms/kg/h), has been administered for provision of additional analgesia during stimulating procedures.

Establishment of alternative analgesia prior to discontinuation of Remifentanyl Noridem: Due to the very rapid offset of action of Remifentanyl Noridem, no residual opioid activity will be present within 5 to 10 minutes after discontinuation regardless of the duration of infusion. Following administration of Remifentanyl Noridem, the possibility of tolerance, hyperalgesia and associated haemodynamic changes should be considered when used in Intensive Care Unit (see section 4.4 *Special warnings and precautions for use*). Therefore, prior to discontinuation of Remifentanyl Noridem, patients must be given alternative analgesic and sedative agents to prevent hyperalgesia and associated haemodynamic changes. These agents must be given at a sufficient time in advance to allow the therapeutic effects of these agents to become established. The range of options for analgesia includes long acting oral, intravenous, or regional analgesics controlled by the nurse or the patient. These techniques should always be titrated to individual patient needs as the infusion of Remifentanyl Noridem is reduced. It is recommended that the choice of agent(s), the dose, and the time of administration are planned prior to discontinuation of Remifentanyl Noridem.

There is a potential for the development of tolerance with time during prolonged administration of μ -opioid agonists.

Guidelines for extubation and discontinuation of Remifentanyl Noridem: In order to ensure a smooth emergence from a Remifentanyl Noridem-based regimen it is recommended that the infusion rate of Remifentanyl Noridem is titrated in stages to 0.1 micrograms/kg/min (6 micrograms/kg/h) over a period up to 1 hour prior to extubation.

Following extubation, the infusion rate should be reduced by 25% decrements in at least 10-minute intervals until the infusion is discontinued. During weaning from the ventilator the Remifentanyl Noridem infusion should not be increased and only down titration should occur, supplemented as required with alternative analgesics.

Upon discontinuation of Remifentanyl Noridem, the IV cannula should be cleared or removed to prevent subsequent inadvertent administration.

When other opioid agents are administered as part of the regimen for transition to alternative analgesia, the patient must be carefully monitored. The benefit of providing adequate analgesia must always be balanced against the potential risk of respiratory depression with these agents.

Paediatric intensive care patients

There are no data available on use in paediatric patients.

Renally-impaired intensive care patients

No adjustments to the doses recommended above are necessary in renally-impaired patients, including those undergoing renal replacement therapy; however, the clearance of the carboxylic acid metabolite is reduced in patients with renal impairment (see section 5.2).

Special patient populations

Elderly (over 65 years of age)

General anaesthesia: The initial starting dose of remifentanyl administered to patients over 65 should be half the recommended adult dose and then shall be titrated to individual patient need as an increased sensitivity to the pharmacological effects of remifentanyl has been seen in this patient population. This dose adjustment applies to use in all phases of anaesthesia including induction, maintenance, and immediate post-operative analgesia.

Because of the increased sensitivity of elderly patients to remifentanyl, when administering Remifentanyl Noridem by TCI in this population the initial target concentration should be 1.5 to 4 nanograms/mL with subsequent titration to response.

Cardiac anaesthesia: No initial dose reduction is required (see this section *Cardiac anaesthesia*).

Intensive Care: No initial dose reduction is required (see this section *Use in intensive care*).

Obese patients

For manually-controlled infusion it is recommended that for obese patients the dosage of Remifentanyl Noridem should be reduced and based upon ideal body weight as the clearance and volume of distribution of remifentanyl are better correlated with ideal body weight than actual body weight.

With the calculation of lean body mass (LBM) used in the Minto model, LBM is likely to be underestimated in female patients with a body mass index (BMI) greater than 35 kg/m² and in male patients with BMI greater than 40 kg/m². To avoid underdosing in these patients, remifentanyl TCI should be titrated carefully to individual response.

Renal impairment

On the basis of investigations carried out to date, a dose adjustment in patients with impaired renal function, including intensive care patients, is not necessary.

Hepatic impairment

Studies carried out with a limited number of patients with impaired liver function, do not justify any special dosage recommendations. However, patients with severe hepatic impairment may be slightly more sensitive to the respiratory depressant effects of remifentanyl (see section 4.4). These patients shall be closely monitored and the dose of remifentanyl shall be titrated to individual patient need.

Neurosurgery

Limited clinical experience in patients undergoing neurosurgery has shown that no special dosage recommendations are required.

ASA III/IV patients

General anaesthesia: As the haemodynamic effects of potent opioids can be expected to be more pronounced in ASA III/IV patients, caution should be exercised in the administration of Remifentanyl Noridem in this population. Initial dosage reduction and subsequent titration to effect is therefore recommended. In paediatric patients, there are insufficient data to make a dosage recommendation.

For TCI, a lower initial target of 1.5 to 4 nanograms/mL should be used in ASA III or IV patients and subsequently titrated to response.

Cardiac anaesthesia: No initial dose reduction is required (see this section *Cardiac anaesthesia*).

4.3 Contraindications

As glycine is present in the formulation, Remifentanyl Noridem is contra-indicated for epidural and intrathecal use (see section 5.3).

Remifentanyl Noridem is contra-indicated in patients with hypersensitivity to the active substance or to other fentanyl analogues or to any of the excipients listed in section 6.1.

Remifentanyl Noridem is contra-indicated for use as the sole agent for induction of anaesthesia.

4.4 Special warnings and precautions for use

Remifentanyl Noridem should be administered only in a setting fully equipped for the monitoring and support of respiratory and cardiovascular function, and by persons specifically trained in the use of anaesthetic drugs and the recognition and management of the expected adverse effects of potent opioids, including respiratory and cardiac resuscitation. Such training must include the establishment and maintenance of a patent airway and assisted ventilation. The use of Remifentanyl Noridem in mechanically ventilated intensive care patients is not recommended for a duration of treatment greater than 3 days.

Patients with known hypersensitivity to opioids of a different class may exhibit a hypersensitivity reaction following administration of Remifentanyl Noridem. Caution should be exercised before using Remifentanyl Noridem in these patients (see section 4.3).

Rapid offset of action/Transition to alternative analgesia

Due to the very rapid offset of action of Remifentanyl Noridem, no residual opioid activity will be present within 5 to 10 minutes after the discontinuation of Remifentanyl Noridem. For those patients undergoing surgical procedures where post-operative pain is anticipated, analgesics should be administered prior to discontinuation of Remifentanyl Noridem. The possibility of tolerance, hyperalgesia and associated haemodynamic changes should be considered when used in Intensive Care Unit (see section 4.2 *Posology and method of administration*). Prior to discontinuation of Remifentanyl Noridem, patients must be given alternative analgesic and sedative agents. Sufficient time must be allowed to reach the therapeutic effect of the longer acting analgesic. The choice of agent(s), the dose and the time of administration should be planned in advance and individually tailored to be appropriate for the patient's surgical procedure and the level of post-operative care anticipated. When other opioid agents are administered as part of the regimen for transition to alternative analgesia, the benefit of providing adequate post-operative analgesia must always be balanced against the potential risk of respiratory depression with these agents.

Risk from concomitant use of sedative medicines such as benzodiazepines or related drugs

Concomitant use of Remifentanyl Noridem and sedative medicines such as benzodiazepines or related drugs may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing with these sedative medicines should be reserved for patients for whom alternative treatment options are not possible. If a decision is made to prescribe Remifentanyl Noridem concomitantly with sedative medicines, the lowest effective dose should be used, and the duration of treatment should be as short as possible.

The patients should be followed closely for signs and symptoms of respiratory depression and sedation. In this respect, it is strongly recommended to inform patients and their caregivers to be aware of these symptoms (see section 4.5).

Discontinuation of treatment and withdrawal syndrome

Repeated administration at short term intervals for prolonged periods may result in the development of withdrawal syndrome after cessation of therapy. Symptoms following withdrawal of remifentanyl including tachycardia, hypertension and agitation

have been reported infrequently upon abrupt cessation, particularly after prolonged administration of more than 3 days. Where reported, re-introduction and tapering of the infusion has been beneficial. The use of remifentanyl in mechanically ventilated intensive care patients is not recommended for duration of treatment greater than 3 days.

Muscle rigidity - prevention and management

At the doses recommended muscle rigidity may occur. As with other opioids, the incidence of muscle rigidity is related to the dose and rate of administration. Therefore, slow bolus injections should be administered over not less than 30 seconds.

Muscle rigidity induced by remifentanyl must be treated in the context of the patient's clinical condition with appropriate supporting measures. Excessive muscle rigidity occurring during the induction of anaesthesia should be treated by the administration of a neuromuscular blocking agent and/or additional hypnotic agents. Muscle rigidity seen during the use of remifentanyl as an analgesic may be treated by stopping or decreasing the rate of administration of remifentanyl. Resolution of muscle rigidity after discontinuing the infusion of remifentanyl occurs within minutes. Alternatively, an opioid antagonist may be administered, however this may reverse or attenuate the analgesic effect of remifentanyl.

Respiratory depression – prevention and management

As with all potent opioids, profound analgesia is accompanied by marked respiratory depression. Therefore, remifentanyl should only be used in areas where facilities for monitoring and dealing with respiratory depression are available. Special care should be taken in patients with respiratory dysfunction. The appearance of respiratory depression should be managed appropriately, including decreasing the rate of infusion by 50%, or a temporary discontinuation of the infusion. Unlike other fentanyl analogues, remifentanyl has not been shown to cause recurrent respiratory depression even after prolonged administration. However, as many factors may affect post-operative recovery it is important to ensure that full consciousness and adequate spontaneous ventilation are achieved before the patient is discharged from the recovery area.

Cardiovascular effects

The risk of cardiovascular effects such as hypotension and bradycardia, which may rarely lead to asystole/cardiac arrest (see sections 4.5 and 4.8) may be reduced by lowering the rate of infusion of Remifentanyl Noridem or the dose of concurrent anaesthetics or by using IV fluids, vasopressor or anticholinergic agents as appropriate.

Debilitated, hypovolaemic, hypotensive and elderly patients may be more sensitive to the cardiovascular effects of remifentanyl.

Inadvertent administration

A sufficient amount of Remifentanyl Noridem may be present in the dead space of the IV line and/or cannula to cause respiratory depression, apnoea and/or muscle rigidity if the line is flushed with IV fluids or other drugs. This may be avoided by administering Remifentanyl Noridem into a fast-flowing IV line or via a dedicated IV line, which is removed when Remifentanyl Noridem is discontinued.

Neonates/infants

There is limited data available on use in neonates/infants under 1 year of age (see sections 4.2 *Neonates/infants (aged less than 1 year)* and 5.1).

Tolerance and opioid use disorder (abuse and dependence)

Tolerance, physical and psychological dependence, and opioid use disorder (OUD) may develop upon repeated administration of opioids. Abuse or intentional misuse of opioids may result in overdose and/or death. The risk of developing OUD is increased in patients with a personal or a family history (parents or siblings) of substance use disorders (including alcohol use disorder), in current tobacco users or in patients with a personal history of other mental health disorders (e.g. major depression, anxiety and personality disorders).

4.5 Interaction with other medicinal products and other forms of interaction

Remifentanyl is not metabolised by plasmacholinesterase, therefore, interactions with drugs metabolised by this enzyme are not anticipated.

As with other opioids, remifentanyl, whether given by manually-controlled infusion or TCI, decreases the doses of inhaled and IV anaesthetics, and benzodiazepines required for anaesthesia (see section 4.2). If doses of concomitantly administered CNS depressant drugs are not reduced, patients may experience an increased incidence of adverse effects associated with these agents.

Sedative medicines such as benzodiazepines or related drugs: The concomitant use of opioids with sedative medicines such as benzodiazepines or related drugs increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. The dose and duration of concomitant use should be limited (see section 4.4). The concomitant use of opioids and gabapentinoids (gabapentin and pregabalin) increases the risk of opioid overdose, respiratory depression and death.

Co-administration of remifentanyl with a serotonergic agent, such as Selective Serotonin Reuptake Inhibitors (SSRIs), Serotonin Norepinephrine Reuptake Inhibitors (SNRIs) or Monoamine Oxidase Inhibitors (MAOIs) may increase the risk of serotonin syndrome, a potentially life-threatening condition. Caution should be exercised with concomitant use of MAOIs. Irreversible MAOIs should be discontinued at least 2 weeks prior to remifentanyl use.

The cardiovascular effects of remifentanyl (hypotension and bradycardia – see section 4.4 and 4.8) may be exacerbated in patients receiving concomitant cardiac depressant drugs, such as beta-blockers and calcium channel blocking agents.

After receiving remifentanyl, it is advisable that alcoholic drink is avoided.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate and well-controlled studies in pregnant women. Remifentanyl Noridem should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus.

Breast-feeding

It is not known whether remifentanyl is excreted in human milk. However, because fentanyl analogues are excreted in human milk and remifentanyl-related material was found in rat milk after dosing with remifentanyl, nursing mothers should be advised to discontinue breast feeding for 24 hours following administration of remifentanyl.

Labour and delivery

There are insufficient data to recommend remifentanyl for use during labour and Caesarean section. It is known that remifentanyl crosses the placental barrier and fentanyl analogues can cause respiratory depression in the child. In case remifentanyl is administered nevertheless, the patient and the neonate must be monitored for signs of excess sedation or respiratory depression (see section 4.4).

4.7 Effects on ability to drive and use machines

After anaesthesia with remifentanyl the patient should not drive or operate machinery. The physician should decide when these activities may be resumed. It is advisable that the patient is accompanied when returning home.

4.8 Undesirable effects

The most common undesirable effects associated with remifentanyl are direct extensions of μ -opioid agonist pharmacology. These adverse events resolve within minutes of discontinuing or decreasing the rate of remifentanyl administration. The frequencies below are defined as very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1\,000$ to $< 1/100$), rare ($\geq 1/10\,000$ to $< 1/1\,000$), very rare ($< 1/10\,000$), and not known (cannot be estimated from the available data).

System Organ Class	Frequency	Adverse reactions
Immune System Disorders	Rare:	Allergic reactions including anaphylaxis have been reported in patients receiving remifentanyl in conjunction with one or more anaesthetic agents.
	Not known	Anaphylactic shock
Psychiatric Disorders	Not known:	Drug dependence, Withdrawal syndrome
Nervous System Disorders	Very common:	Skeletal muscle rigidity
	Rare:	Sedation (during recovery from general anaesthesia)
	Not known:	Convulsions
Cardiac Disorders	Common:	Bradycardia
	Rare:	Asystole/cardiac arrest, usually preceded by bradycardia, has been reported in patients receiving remifentanyl in conjunction with other anaesthetic agents.

	Not known:	Atrioventricular block, Arrhythmia
Vascular Disorders	Very common:	Hypotension
	Common:	Post-operative hypertension
Respiratory, Thoracic and Mediastinal Disorders	Common:	Acute respiratory depression, apnoea, Cough
	Uncommon:	Hypoxia
Gastrointestinal Disorders	Very common:	Nausea, vomiting
	Uncommon:	Constipation
Skin and Subcutaneous Tissue Disorders	Common:	Pruritus
General Disorders and Administration Site Conditions	Common:	Post-operative shivering
	Uncommon:	Post-operative aches
	Not Know:	Drug tolerance

Discontinuation of treatment

Symptoms following withdrawal of remifentanil including tachycardia, hypertension and agitation have been reported infrequently upon abrupt cessation, particularly after prolonged administration of more than 3 days (see section 4.4).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRa Pharmacovigilance, Website: www.hpra.ie.

4.9 Overdose

As with all potent opioid analgesics, overdose would be manifested by an extension of the pharmacologically predictable actions of remifentanil. Due to the very short duration of action of Remifentanil Noridem, the potential for deleterious effects due to overdose is limited to the immediate time period following drug administration. Response to discontinuation of the drug is rapid, with return to baseline within ten minutes.

In the event of overdose, or suspected overdose, take the following actions: discontinue administration of Remifentanil Noridem, maintain a patent airway, initiate assisted or controlled ventilation with oxygen, and maintain adequate cardiovascular function. If depressed respiration is associated with muscle rigidity, a neuromuscular blocking agent may be required to facilitate assisted or controlled respiration. Intravenous fluids and vasopressor for the treatment of hypotension and other supportive measures may be employed.

Intravenous administration of an opioid antagonist such as naloxone may be given as a specific antidote to manage severe respiratory depression and muscle rigidity. The duration of respiratory depression following overdose with Remifentanil Noridem is unlikely to exceed the duration of action of the opioid antagonist.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Opioid anaesthetics, ATC code: N01AH06.

Mechanism of actions

Remifentanil is a selective μ -opioid agonist with a rapid onset and very short duration of action. The μ -opioid activity of remifentanil is antagonised by narcotic antagonists, such as naloxone.

Pharmacodynamic effects

Assays of histamine in patients and normal volunteers have shown no elevation in histamine levels after administration of remifentanil in bolus doses up to 30 micrograms/kg.

Neonates/infants (aged less than 1 year):

In a randomised (ratio of 2:1, remifentanil:halothane), open label, parallel group, multicentre study in 60 young infants and neonates $\leq$ 8 weeks of age (mean 5.5 weeks) with an ASA physical status of I-II who were undergoing pyloromyotomy, the efficacy and safety of remifentanil (given as a 0.4 μ g/kg/min initial continuous infusion plus supplemental doses or infusion rate changes as needed) was compared with halothane (given at 0.4% with supplemental increases as needed). Maintenance of

anaesthesia was achieved by the additional administration of 70% nitrous oxide (N₂O) plus 30% oxygen. Recovery times were superior in the remifentanyl relative to the halothane groups (not significant).

Use for Total Intravenous anaesthesia (TIVA) - children aged 6 months to 16 years

TIVA with remifentanyl in paediatric surgery was compared to inhalation anaesthesia in three randomised, open-label studies. The results are summarised in the table below.

Surgical intervention	Age (y), (N)	Study condition (maintenance)	Extubation (min) (mean (SD))
Lower abdominal/urological surgery	0.5-16 (120)	TIVA: propofol (5 - 10 mg/kg/h) + remifentanyl (0.125 - 1.0 µg/kg/min)	11.8 (4.2)
		Inhalation anaesthesia: sevoflurane (1.0 - 1.5 MAC) and remifentanyl (0.125 - 1.0 µg/kg/min)	15.0 (5.6) (p<0.05)
ENT-surgery	4-11 (50)	TIVA: propofol (3 mg/kg/h) + remifentanyl (0.5 µg/kg/min)	11 (3.7)
		Inhalation anaesthesia: desflurane (1.3 MAC) and N ₂ O mixture	9.4 (2.9) Not significant
General or ENT surgery	2-12 (153)	TIVA: remifentanyl (0.2 - 0.5 µg/kg/min) + propofol (100 - 200 µg/kg/min)	Comparable extubation times (based on limited data)
		Inhalation anaesthesia: sevoflurane (1 - 1.5 MAC) + N ₂ O mixture	

In the study in lower abdominal/urological surgery comparing remifentanyl/propofol with remifentanyl/sevoflurane, hypotension occurred significantly more often under remifentanyl/sevoflurane, and bradycardia occurred significantly more often under remifentanyl/propofol. In the study in ENT surgery comparing remifentanyl/propofol with desflurane/nitrous oxide, a significantly higher heart rate was seen in subjects receiving desflurane/nitrous oxide compared with remifentanyl/propofol and with baseline values.

5.2 Pharmacokinetic properties

Elimination

Following administration of the recommended doses of remifentanyl, the effective half-life is 3 to 10 minutes. The average clearance of remifentanyl in young healthy adults is 40 mL/min/kg, the central volume of distribution is 100 mL/kg, and the steady-state volume of distribution is 350 mL/kg.

Absorption

Blood concentrations of remifentanyl are proportional to the dose administered throughout the recommended dose range. For every 0.1 micrograms/kg/min increase in infusion rate, the blood concentration of remifentanyl will rise 2.5 nanograms/mL. Remifentanyl is approximately 70% bound to plasma proteins.

Biotransformation

Remifentanyl is an esterase metabolised opioid that is susceptible to metabolism by non-specific blood and tissue esterases. The metabolism of remifentanyl results in the formation of a carboxylic acid metabolite, which in dogs is 1/4600th as potent as remifentanyl. Studies in man indicate that all pharmacological activity is associated with the parent compound. The activity of this metabolite is therefore not of any clinical consequence. The half-life of the metabolite in healthy adults is 2 hours. In patients with normal renal function, the time for 95% elimination of the primary metabolite of remifentanyl by the kidneys is approximately 7 to 10 hours. Remifentanyl is not a substrate for plasma cholinesterase.

Placental and milk transfer

Placental transfer studies in rats and rabbits showed that pups are exposed to remifentanyl and/or its metabolites during growth and development. Remifentanyl-related material is transferred to the milk of lactating rats. In a human clinical trial, the concentration of remifentanyl in foetal blood was approximately 50% of that in maternal blood. The foetal arterio-venous ratio of remifentanyl concentrations was approximately 30%, suggesting metabolism of remifentanyl in the neonate.

Cardiac anaesthesia

The clearance of remifentanyl is reduced by approximately 20% during hypothermic (28°C) cardiopulmonary bypass. A decrease in body temperature lowers elimination clearance by 3% per degree centigrade.

Renal impairment

The rapid recovery from remifentanyl-based sedation and analgesia is unaffected by renal status.

The pharmacokinetics of remifentanyl are not significantly changed in patients with varying degrees of renal impairment even after administration for up to 3 days in the intensive care setting.

The clearance of the carboxylic acid metabolite is reduced in patients with renal impairment. In intensive care patients with moderate/severe renal impairment, the concentration of the carboxylic acid metabolite may exceed 250-fold the level of remifentanyl at steady-state in some patients. The available clinical data demonstrates that accumulation of the metabolite does not result in clinically relevant μ -opioid effects even after administration of remifentanyl infusions for up to 3 days in these patients. There are insufficient data available on the safety and pharmacokinetic profile of the metabolite following infusions of Remifentanyl Noridem for durations greater than 3 days.

There is no evidence that remifentanyl is extracted during renal replacement therapy.

The carboxylic acid metabolite is extracted during haemodialysis by at least 30%.

Hepatic impairment

The pharmacokinetics of remifentanyl are not changed in patients with severe hepatic impairment awaiting liver transplant, or during the anhepatic phase of liver transplant surgery. Patients with severe hepatic impairment may be slightly more sensitive to the respiratory depressant effects of remifentanyl. These patients should be closely monitored and the dose of remifentanyl should be titrated to the individual patient need.

Paediatric patients

The average clearance and steady state volume of distribution of remifentanyl are increased in younger children and decline to young healthy adult values by age 17. The elimination half-life of remifentanyl in neonates is not significantly different from that of young healthy adults. Changes in analgesic effect after changes in infusion rate of remifentanyl should be rapid and similar to those seen in young healthy adults. The pharmacokinetics of the carboxylic acid metabolite in paediatric patients 2 to 17 years of age are similar to those seen in adults after correcting for differences in body weight.

Elderly

The clearance of remifentanyl is slightly reduced in elderly patients (>65 years) compared to young patients. The pharmacodynamic activity of remifentanyl increases with increasing age. Elderly patients have a remifentanyl EC₅₀ for formation of delta waves on the electroencephalogram (EEG) that is 50% lower than young patients; therefore, the initial dose of remifentanyl should be reduced by 50% in elderly patients and then carefully titrated to meet the individual patient need.

5.3 Preclinical safety data

Remifentanyl, like some other fentanyl analogues, produced increases in action potential duration (APD) in dog isolated Purkinje fibres. There were no effects at a concentration of 0.1 micromolar (38 nanograms/ml). Effects were seen at a concentration of 1 micromolar (377 nanograms/ml), and were statistically significant at a concentration of 10 micromolar (3770 nanograms/mL). These concentrations are 12-fold and 119-fold respectively the highest likely free concentrations (or 3-fold and 36-fold respectively, the highest likely whole blood concentrations) following the maximum recommended therapeutic dose.

Acute toxicity

Expected signs of μ -opioid intoxication were observed in non-ventilated mice, rats, and dogs after large single bolus intravenous doses of remifentanyl. In these studies, the most sensitive species, the male rat, survived following administration of 5 mg/kg. Hypoxia-induced brain microhaemorrhages observed in dogs were reversed within 14 days after completion of dosing.

Repeat dose toxicity

Bolus doses of remifentanyl administered to non-ventilated rats and dogs resulted in respiratory depression in all dose groups, and in reversible brain microhaemorrhages in dogs. Subsequent investigations showed that the microhaemorrhages resulted

from hypoxia and were not specific to remifentanyl. Brain microhaemorrhages were not observed in infusion studies in non-ventilated rats and dogs because these studies were conducted at doses that did not cause severe respiratory depression.

It is to be derived from preclinical studies that respiratory depression and associated sequelae are the most likely cause of potentially serious adverse events in humans.

Intrathecal administration to dogs of the glycine formulation alone (i.e. without remifentanyl) caused agitation, pain, and hind limb dysfunction and incoordination. These effects are believed to be secondary to the glycine excipient. Because of the better buffering properties of blood, the more rapid dilution, and the low glycine concentration of the remifentanyl formulation, this finding has no clinical relevance for intravenous administration of remifentanyl.

Reproductive toxicity studies

Remifentanyl reduced fertility in male rats after daily injection for at least 70 days. A no-effect dose was not demonstrated. Fertility was not affected in female rats. Teratogenic effects were not seen in rats or rabbits.

Administration of remifentanyl to rats throughout late gestation and lactation did not significantly affect the survival, development, or reproductive performance of the F₁ generation.

Genotoxicity

Remifentanyl did not yield positive findings in a series of *in vitro* and *in vivo* genotoxicity tests, except in the *in vitro* mouse lymphoma tk assay, which gave a positive result with metabolic activation. Since the mouse lymphoma results could not be confirmed in further *in vitro* and *in vivo* tests, treatment with remifentanyl is not considered to pose a genotoxic hazard to patients.

Carcinogenicity

Long-term carcinogenicity studies were not performed.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Glycine
Hydrochloric acid (*for pH adjustment*)

6.2 Incompatibilities

Remifentanyl Noridem should only be reconstituted and diluted with those infusion solutions recommended (see section 6.6).

It should not be reconstituted, diluted or mixed with Lactated Ringer's Injection or Lactated Ringer's and 5% Dextrose Injection.

Remifentanyl Noridem should not be mixed with propofol in the same infusion bag prior to administration.

For compatibility when given into a running IV catheter, please see section 6.6.

Administration of Remifentanyl Noridem into the same intravenous line with blood/serum/plasma is not recommended. Non-specific esterases in blood products may lead to the hydrolysis of remifentanyl to its inactive metabolite.

Remifentanyl Noridem should not be mixed with other therapeutic agents prior to administration.

6.3 Shelf life

Vials	Remifentanyl Noridem 5 mg Powder for Concentrate for Solution for Injection or Infusion: 3 years
After reconstitution/dilution	Chemical and physical in-use stability has been demonstrated for 24 hours at 25 °C. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8 °C, unless reconstitution/dilution has taken place in controlled and validated aseptic conditions.

6.4 Special precautions for storage

Do not store above 25°C. Keep the vial in the outer carton in order to protect from light.
For storage conditions of the reconstituted / diluted medicinal product, see section 6.3.

6.5 Nature and contents of container

Glass vial with bromobutyl rubber stopper and aluminium seal with a dark blue plastic flip-off cap.

5 or 10 vials.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Remifentanil Noridem should be prepared for intravenous use by adding, as appropriate 1, 2, or 5 mL of diluent to give a reconstituted solution with a concentration of 1 mg/mL remifentanil. The reconstituted solution is clear, colourless, and practically free from particulate material. After reconstitution, visually inspect the product (where the container permits) for particulate material, discolouration or damage of container. Discard any solution where such defects are observed. Reconstituted product is for single use only. Any unused material should be discarded.

Remifentanil Noridem should not be administered by manually-controlled infusion without further dilution to concentrations of 20 to 250 micrograms/mL (50 micrograms/mL is the recommended dilution for adults and 20 to 25 micrograms/mL for paediatric patients aged 1 year and over).

Remifentanil Noridem should not be administered by TCI without further dilution (20 to 50 micrograms/mL is the recommended dilution for TCI).

The dilution is dependent upon the technical capability of the infusion device and the anticipated requirements of the patient.

One of the following IV fluids listed below should be used for dilution:

Water for Injections

Glucose 5% solution for injection

Glucose 5% and Sodium Chloride 0.9% solution for injection

Sodium Chloride 0.9% solution for injection

Sodium Chloride 0.45% solution for injection

After dilution, visually inspect the product to ensure it is clear, colourless, practically free from particulate matter and the container is undamaged. Discard any solution where such defects are observed.

Remifentanil Noridem has been shown to be compatible with the following intravenous fluids when administered into a running IV catheter:

Lactated Ringer's solution for injection

Lactated Ringer's and Glucose 5% solution for injection

Remifentanil Noridem has been shown to be compatible with propofol when administered into a running IV catheter.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Tables 6-11 give guidelines for infusion rates of Remifentanil Noridem for manually-controlled infusion:

Table 6. Infusion Rates (mL/kg/h) of Remifentanil Noridem Powder for Concentrate for Solution for Injection or Infusion

Drug Delivery Rate (micrograms/kg/min)	Infusion Delivery Rate (mL/kg/h) for Solution Concentrations of			
	20 micrograms /mL 1 mg/50 mL	25 micrograms /mL 1 mg/40 mL	50 micrograms /mL 1 mg/20 mL	250 micrograms /mL 10 mg/40 mL
0.0125	0.038	0.03	0.015	Not recommended
0.025	0.075	0.06	0.03	Not recommended
0.05	0.15	0.12	0.06	0.012

0.075	0.23	0.18	0.09	0.018
0.1	0.3	0.24	0.12	0.024
0.15	0.45	0.36	0.18	0.036
0.2	0.6	0.48	0.24	0.048
0.25	0.75	0.6	0.3	0.06
0.5	1.5	1.2	0.6	0.12
0.75	2.25	1.8	0.9	0.18
1.0	3.0	2.4	1.2	0.24
1.25	3.75	3.0	1.5	0.3
1.5	4.5	3.6	1.8	0.36
1.75	5.25	4.2	2.1	0.42
2.0	6.0	4.8	2.4	0.48

Table 7. Infusion Rates (mL/h) of Remifentanil Noridem Powder for Concentrate for Solution for Injection or Infusion for a 20 micrograms/mL Solution

Infusion Rate (micrograms/kg/min)	Patient Weight (kg)						
	5	10	20	30	40	50	60
0.0125	0.188	0.375	0.75	1.125	1.5	1.875	2.25
0.025	0.375	0.75	1.5	2.25	3.0	3.75	4.5
0.05	0.75	1.5	3.0	4.5	6.0	7.5	9.0
0.075	1.125	2.25	4.5	6.75	9.0	11.25	13.5
0.1	1.5	3.0	6.0	9.0	12.0	15.0	18.0
0.15	2.25	4.5	9.0	13.5	18.0	22.5	27.0
0.2	3.0	6.0	12.0	18.0	24.0	30.0	36.0
0.25	3.75	7.5	15.0	22.5	30.0	37.5	45.0
0.3	4.5	9.0	18.0	27.0	36.0	45.0	54.0
0.35	5.25	10.5	21.0	31.5	42.0	52.5	63.0
0.4	6.0	12.0	24.0	36.0	48.0	60.0	72.0

Table 8. Infusion Rates (mL/h) of Remifentanil Noridem Powder for Concentrate for Solution for Injection or Infusion for a 25 micrograms/mL Solution

Infusion Rate (micrograms/kg/min)	Patient Weight (kg)									
	10	20	30	40	50	60	70	80	90	100
0.0125	0.3	0.6	0.9	1.2	1.5	1.8	2.1	2.4	2.7	3.0
0.025	0.6	1.2	1.8	2.4	3.0	3.6	4.2	4.8	5.4	6.0
0.05	1.2	2.4	3.6	4.8	6.0	7.2	8.4	9.6	10.8	12.0
0.075	1.8	3.6	5.4	7.2	9.0	10.8	12.6	14.4	16.2	18.0
0.1	2.4	4.8	7.2	9.6	12.0	14.4	16.8	19.2	21.6	24.0
0.15	3.6	7.2	10.8	14.4	18.0	21.6	25.2	28.8	32.4	36.0
0.2	4.8	9.6	14.4	19.2	24.0	28.8	33.6	38.4	43.2	48.0

Table 9. Infusion Rates (mL/h) of Remifentanil Noridem Powder for Concentrate for Solution for Injection or Infusion for a 50 micrograms/mL Solution

Infusion Rate (micrograms/kg/min)	Patient Weight (kg)							
	30	40	50	60	70	80	90	100
0.025	0.9	1.2	1.5	1.8	2.1	2.4	2.7	3.0
0.05	1.8	2.4	3.0	3.6	4.2	4.8	5.4	6.0
0.075	2.7	3.6	4.5	5.4	6.3	7.2	8.1	9.0
0.1	3.6	4.8	6.0	7.2	8.4	9.6	10.8	12.0
0.15	5.4	7.2	9.0	10.8	12.6	14.4	16.2	18.0
0.2	7.2	9.6	12.0	14.4	16.8	19.2	21.6	24.0
0.25	9.0	12.0	15.0	18.0	21.0	24.0	27.0	30.0
0.5	18.0	24.0	30.0	36.0	42.0	48.0	54.0	60.0
0.75	27.0	36.0	45.0	54.0	63.0	72.0	81.0	90.0

1.0	36.0	48.0	60.0	72.0	84.0	96.0	108.0	120.0
1.25	45.0	60.0	75.0	90.0	105.0	120.0	135.0	150.0
1.5	54.0	72.0	90.0	108.0	126.0	144.0	162.0	180.0
1.75	63.0	84.0	105.0	126.0	147.0	168.0	189.0	210.0
2.0	72.0	96.0	120.0	144.0	168.0	192.0	216.0	240.0

Table 10. Infusion Rates (mL/h) of Remifentanil Noridem Powder for Concentrate for Solution for Injection or Infusion for a 250 micrograms/mL Solution

Infusion Rate (micrograms/kg/min)	Patient Weight (kg)							
	30	40	50	60	70	80	90	100
0.1	0.72	0.96	1.20	1.44	1.68	1.92	2.16	2.40
0.15	1.08	1.44	1.80	2.16	2.52	2.88	3.24	3.60
0.2	1.44	1.92	2.40	2.88	3.36	3.84	4.32	4.80
0.25	1.80	2.40	3.00	3.60	4.20	4.80	5.40	6.00
0.5	3.60	4.80	6.00	7.20	8.40	9.60	10.80	12.00
0.75	5.40	7.20	9.00	10.80	12.60	14.40	16.20	18.00
1.0	7.20	9.60	12.00	14.40	16.80	19.20	21.60	24.00
1.25	9.00	12.00	15.00	18.00	21.00	24.00	27.00	30.00
1.5	10.80	14.40	18.00	21.60	25.20	28.80	32.40	36.00
1.75	12.60	16.80	21.00	25.20	29.40	33.60	37.80	42.00
2.0	14.40	19.20	24.00	28.80	33.60	38.40	43.20	48.00

Table 11 provides the equivalent blood remifentanil concentration using a TCI approach for various manually-controlled infusion rates at steady state:

Table 11. Remifentanil Blood Concentrations (nanograms/mL) estimated using the Minto (1997) Pharmacokinetic Model in a 70 kg, 170 cm, 40 Year Old Male Patient for Various Manually-Controlled Infusion rates (micrograms /kg/min) at Steady State.

Remifentanil Noridem Infusion Rate (micrograms /kg/min)	Remifentanil Blood Concentration (nanograms/mL)
0.05	1.3
0.10	2.6
0.25	6.3
0.40	10.4
0.50	12.6
1.0	25.2
2.0	50.5

7 MARKETING AUTHORISATION HOLDER

Noridem Enterprises Ltd
Evagorou & Makariou
Mitsi Building 3, Office 115
1065 Nicosia
Cyprus

8 MARKETING AUTHORISATION NUMBER

PA1122/014/003

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of First Authorisation: 21st December 2011
Date of last Renewal: 31st January 2015

10 DATE OF REVISION OF THE TEXT

May 2025
01 May 2025