

**IRISH MEDICINES BOARD ACTS 1995 AND 2006**

**MEDICINAL PRODUCTS(CONTROL OF PLACING ON THE MARKET)REGULATIONS,2007**

**(S.I. No.540 of 2007)**

**PA1285/016/001**

Case No: 2039802

The Irish Medicines Board in exercise of the powers conferred on it by the above mentioned Regulations hereby grants to

**Medis ehf**

**Medis ehf, Reykjavikurvegur 78, 220 Hafnarfjorour, Iceland**

an authorisation, subject to the provisions of the said Regulations, in respect of the product

**Glimepiride 1 mg Tablets**

The particulars of which are set out in Part I and Part II of the attached Schedule. The authorisation is also subject to the general conditions as may be specified in the said Regulations as listed on the reverse of this document.

This authorisation, unless previously revoked, shall continue in force from **13/09/2007** until **21/07/2010**.

Signed on behalf of the Irish Medicines Board this

\_\_\_\_\_

A person authorised in that behalf by the said Board.

## Part II

### Summary of Product Characteristics

#### 1 NAME OF THE MEDICINAL PRODUCT

Glimepiride 1 mg Tablets

#### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 1 mg of glimepiride.

For excipients, see 6.1.

#### 3 PHARMACEUTICAL FORM

Tablet.

Pink, scored, oblong, flat with bevelled edges.

#### 4 CLINICAL PARTICULARS

##### 4.1 Therapeutic Indications

Glimepiride is indicated for the treatment of type 2 diabetes mellitus, when diet, physical exercise and weight reduction alone are not adequate.

##### 4.2 Posology and method of administration

The basis for successful treatment of diabetes is a good diet, regular physical activity, as well as routine checks of blood and urine. Tablets or insulin can not compensate if the patient does not keep to the recommended diet.

Dosage is determined by the results of blood and urinary glucose determinations.

The starting dose is 1 mg glimepiride per day. If good control is achieved this dosage should be used for maintenance therapy.

If control is unsatisfactory the dosage should be increased, based on the glycaemic control, in a stepwise manner with an interval of about 1 to 2 weeks between each step, to 2, 3 or 4 mg glimepiride per day.

A dosage of more than 4 mg glimepiride per day gives better results only in exceptional cases. The maximum recommended dose is 6 mg glimepiride per day.

In patients not adequately controlled with the maximum daily dose of glimepiride, concomitant insulin therapy can be initiated if necessary. While maintaining the glimepiride dose, insulin treatment is started at low dose and titrated up depending on the desired level of metabolic control. The combination therapy should be initiated under close medical supervision.

Normally a single daily dose of glimepiride is sufficient. It is recommended that this dose be taken shortly before or during a substantial breakfast or - if none is taken - shortly before or during the first main meal.

If a dose is forgotten, this should not be corrected by increasing the next dose.  
Tablets should be swallowed whole with some liquid.

If a patient has a hypoglycaemic reaction on 1 mg glimepiride daily, this indicates that they can be controlled by diet alone.

In the course of treatment, as an improvement in control of diabetes is associated with higher insulin sensitivity, glimepiride requirements may fall. To avoid hypoglycaemia timely dose reduction or cessation of therapy must therefore be considered. Change in dosage may also be necessary, if there are changes in weight or life style of the patient, or other factors that increase the risk of hypo- or hyperglycaemia.

**Switch over from other oral hypoglycaemic agents to glimepiride:**

A switch over from other oral hypoglycaemic agents to glimepiride can generally be done. For the switch over to glimepiride the strength and the half life of the previous medication has to be taken into account. In some cases, especially in antidiabetics with a long half life (e.g. chlorpropamide), a wash out period of a few days is advisable in order to minimise the risk of hypoglycaemic reactions due to the additive effect. The recommended starting dose is 1 mg glimepiride per day. Based on the response the glimepiride dosage may be increased stepwise, as indicated earlier.

**Switch over from Insulin to glimepiride:**

In exceptional cases, where type 2 diabetic patients are regulated on insulin, a changeover to glimepiride may be indicated. The changeover should be undertaken under close medical supervision.

**Use in renal or hepatic impairment:**

See section 4.3 Contraindications.

### 4.3 Contraindications

Glimepiride should not be used in the following cases: insulin dependent diabetes, diabetic coma, ketoacidosis, severe renal or hepatic function disorders, hypersensitivity to glimepiride, other sulphonylureas or sulphonamides or excipients in the tablet.

In case of severe renal or hepatic function disorders, a change over to insulin is required.

Glimepiride is contra-indicated in pregnancy and lactation.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactose deficiency or glucose-galactose malabsorption should not take this medicine.

### 4.4 Special warnings and precautions for use

Glimepiride must be taken shortly before or during a meal.

When meals are taken at irregular hours or skipped altogether, treatment with glimepiride may lead to hypoglycaemia. Possible symptoms of hypoglycaemia include: headache, ravenous hunger, nausea, vomiting, lassitude, sleepiness, disordered sleep, restlessness, aggressiveness, impaired concentration, impaired alertness and reaction time, depression, confusion, speech and visual disorders, aphasia, tremor, paresis, sensory disturbances, dizziness, helplessness, loss of self-control, delirium, cerebral convulsions, somnolence and loss of consciousness up to and including coma, shallow respiration and bradycardia.

In addition, signs of adrenergic counter-regulation may be present such as sweating, clammy skin, anxiety, tachycardia, hypertension, palpitations, angina pectoris and cardiac arrhythmias.

The clinical picture of a severe hypoglycaemic attack may resemble that of a stroke. Symptoms can almost always be promptly controlled by immediate intake of carbohydrates (sugar). Artificial sweeteners have no effect.

It is known from other sulphonylureas that, despite initially successful countermeasures, hypoglycaemia may recur.

Severe hypoglycaemia or prolonged hypoglycaemia, only temporarily controlled by the usual amounts of sugar, require immediate medical treatment and occasionally hospitalisation.

Factors favouring hypoglycaemia include:

- unwillingness or (more commonly in older patients) incapacity of the patient to cooperate,
- undernutrition, irregular mealtimes or missed meals or periods of fasting,
- alterations in diet,
- imbalance between physical exertion and carbohydrate intake,
- consumption of alcohol, especially in combination with skipped meals,
- impaired renal function,
- serious liver dysfunction,
- overdosage with glimepiride,
- certain uncompensated disorders of the endocrine system affecting carbohydrate metabolism or counterregulation of hypoglycaemia (as for example in certain disorders of thyroid function and in anterior pituitary or adrenocortical insufficiency),
- concurrent administration of certain other medicines (see “Interactions”).

Treatment with glimepiride requires regular monitoring of glucose levels in blood and urine. In addition determination of the proportion of glycosylated haemoglobin is recommended.

Regular hepatic and haematological monitoring (especially leucocytes and thrombocytes) are required during treatment with glimepiride.

In stress-situations (e.g. accidents, acute operations, infections with fever, etc.) a temporary switch to insulin may be indicated.

No experience has been gained concerning the use of glimepiride in patients with severe impairment of liver function or dialysis patients. In patients with severe impairment of renal or liver function change over to insulin is indicated.

Treatment of patients with G6PD-deficiency with sulfonylurea agents can lead to haemolytic anaemia. Since Glimepiride belongs to the class of sulfonylurea agents, caution should be used in patients with G6PD-deficiency and a non-sulfonylurea alternative should be considered.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

If glimepiride is taken simultaneously with certain other medicines, both undesired increases and decreases in the hypoglycaemic action of glimepiride can occur. For this reason, other medicines should only be taken with the knowledge (or at the prescription) of the doctor.

Based on the experience with glimepiride and with other sulphonylureas the following interactions have to be mentioned.

Potential of the blood-glucose-lowering effect and, thus, in some instances hypoglycaemia may occur when one of the following drugs is taken, for example:

- phenylbutazone, azapropazone and oxyfenbutazone
- insulin and oral antidiabetic products
- metformin
- salicylates and p-amino-salicylic acid
- anabolic steroids and male sex hormones
- chloramphenicol
- coumarin anticoagulants
- fenfluramine
- fibrates
- ACE inhibitors
- fluoxetine
- allopurinol

- sympatholytics
- cyclo-, tro- and phosphamides
- sulphinpyrazone
- certain long acting sulphonamides
- tetracyclines
- MAO-inhibitors
- quinolone antibiotics
- probenecid
- miconazole
- pentoxifylline (high dose parenteral)
- tritoqualine

Weakening of the blood-glucose-lowering effect and, thus raised blood glucose levels may occur when one of the following drugs is taken, for example:

- oestrogens and progestagens
- saluretics, thiazide diuretics
- thyroid stimulating agents, glucocorticoids
- phenothiazine derivatives, chlorpromazine
- adrenaline and sympathicomimetics
- nicotinic acid (high dosages) and nicotinic acid derivatives
- laxatives (long term use)
- phenytoin, diazoxide
- glucagon, barbiturates and rifampicin
- acetazolamide

H<sub>2</sub> antagonists, betablockers, clonidine and reserpine may lead to either potentiation or weakening of the blood glucose lowering effect.

Under the influence of sympatholytic drugs such as betablockers, clonidine, guanethidine and reserpine, the signs of adrenergic counterregulation to hypoglycaemia may be reduced or absent.

Alcohol intake may potentiate or weaken the hypoglycaemic action of glimepiride in an unpredictable fashion.

Glimepiride may either potentiate or weaken the effects of coumarin derivatives.

## 4.6 Pregnancy and lactation

### Pregnancy

Glimepiride is contra-indicated during pregnancy. The use of insulin is required under such circumstances. Patients who consider pregnancy should inform their physician.

### Lactation

Because sulphonylurea derivatives like glimepiride pass into the breast milk, glimepiride must not be taken by breast-feeding women.

## 4.7 Effects on ability to drive and use machines

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia or hyperglycaemia or, for example, as a result of visual impairment. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery).

Patients should be advised to take precautions to avoid hypoglycaemia whilst driving. This is particularly important in those who have reduced or absent awareness of the warning symptoms of hypoglycaemia or have frequent episodes of hypoglycaemia. It should be considered whether it is advisable to drive or operate machinery in these circumstances.

## 4.8 Undesirable effects

Based on experience with glimepiride and with other sulphonylureas the following side effects have to be mentioned.

### Immune system disorders

In very rare cases mild hypersensitivity reactions may develop into serious reactions with dyspnoea, fall in blood pressure and sometimes shock. Allergic vasculitis is possible in very rare cases.

Cross allergenicity with sulphonylureas, sulphonamides or related substances is possible.

### Blood and lymphatic system disorders

Changes in haematology are rare during glimepiride treatment. Moderate to severe thrombocytopenia, leucopenia, erythrocytopenia, granulocytopenia, agranulocytosis, haemolytic anaemia and pancytopenia may occur.

These are in general reversible upon discontinuation of medication.

### Metabolism and nutrition disorders

In rare cases hypoglycaemic reactions have been observed after administration of glimepiride. These reactions mostly occur immediately, may be severe and are not always easy to correct. The occurrence of such reactions depends, as with other hypoglycaemic therapies, on individual factors such as dietary habits and the dosage (see further under "Special warnings and special precautions for use").

### Eye disorders

Transient visual disturbances may occur especially on initiation of treatment, due to changes in blood glucose levels.

### Gastrointestinal disorders

Gastrointestinal complaints like nausea, vomiting and diarrhoea, pressure or a feeling of fullness in the stomach and abdominal pain are very rare and seldom lead to discontinuation of therapy.

### Hepato-biliary disorders

Elevation of liver enzymes may occur. In very rare cases, impairment of liver function (e.g. with cholestasis and jaundice) may develop, as well as hepatitis which may progress to liver failure.

### Skin and subcutaneous tissue disorders

Hypersensitivity reactions of the skin may occur as itching, rash and urticaria.

In very rare cases hypersensitivity to light may occur.

### Investigations

In very rare cases, a decrease in the sodium serum concentrations may occur.

## 4.9 Overdose

After ingestion of an overdose hypoglycaemia may occur, lasting from 12 to 72 hours, and may recur after an initial recovery. Symptoms may not be present for up to 24 hours after ingestion. In general observation in hospital is recommended. Nausea, vomiting and epigastric pain may occur. The hypoglycaemia may in general be accompanied by neurological symptoms like restlessness, tremor, visual disturbances, co-ordination problems, sleepiness, coma and convulsions.

Treatment primarily consists of preventing absorption by inducing vomiting and then drinking water or lemonade with activated charcoal (adsorbent) and sodium-sulphate (laxative). If large quantities have been ingested, gastric lavage is indicated, followed by activated charcoal and sodium-sulphate. In case of (severe) overdose hospitalisation in an intensive care department is indicated. Start the administration of glucose as soon as possible, if necessary by a bolus intravenous injection of 50 ml of a 50% solution, followed by an infusion of a 10% solution with strict monitoring of blood glucose. Further treatment should be symptomatic.

In particular when treating hypoglycaemia due to accidental intake of glimepiride in infants and young children, the

dose of glucose given must be carefully controlled to avoid the possibility of producing dangerous hyperglycaemia. Blood glucose should be closely monitored.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Oral blood glucose lowering drugs: Sulfonamides, urea derivatives.  
ATC Code: A10B B12.

Glimepiride is an orally active hypoglycaemic substance belonging to the sulphonylurea group. It may be used in non-insulin dependent diabetes mellitus.

Glimepiride acts mainly by stimulating insulin release from pancreatic beta cells.

As with other sulphonylureas this effect is based on an increase of responsiveness of the pancreatic beta cells to the physiological glucose stimulus. In addition, glimepiride seems to have pronounced extrapancreatic effects also postulated for other sulphonylureas.

#### **Insulin release:**

Sulphonylureas regulate insulin secretion by closing the ATP-sensitive potassium channel in the beta cell membrane. Closing the potassium channel induces depolarisation of the beta cell and results - by opening of calcium channels - in an increased influx of calcium into the cell.

This leads to insulin release through exocytosis.

Glimepiride binds with a high exchange rate to a beta cell membrane protein which is associated with the ATP-sensitive potassium channel but which is different from the usual sulphonylurea binding site.

#### **Extrapancreatic activity:**

The extrapancreatic effects are for example an improvement of the sensitivity of the peripheral tissue for insulin and a decrease of the insulin uptake by the liver.

The uptake of glucose from blood into peripheral muscle and fat tissues occurs via special transport proteins, located in the cell membrane. The transport of glucose in these tissues is the rate limiting step in the use of glucose. Glimepiride increases very rapidly the number of active glucose transport molecules in the plasma membranes of muscle and fat cells, resulting in stimulated glucose uptake.

Glimepiride increases the activity of the glycosyl-phosphatidylinositol-specific phospholipase C which may be correlated with the drug-induced lipogenesis and glycogenesis in isolated fat and muscle cells.

Glimepiride inhibits the glucose production in the liver by increasing the intracellular concentration of fructose-2,6-bisphosphate, which in its turn inhibits the gluconeogenesis

#### **General**

In healthy persons, the minimum effective oral dose is approximately 0.6 mg. The effect of glimepiride is dose-dependent and reproducible. The physiological response to acute physical exercise, reduction of insulin secretion, is still present under glimepiride.

There was no significant difference in effect regardless of whether the drug was given 30 minutes or immediately before a meal. In diabetic patients, good metabolic control over 24 hours can be achieved with a single daily dose.

Although the hydroxy metabolite of glimepiride caused a small but significant decrease in serum glucose in healthy persons, it accounts for only a minor part of the total drug effect.

### **Combination therapy with insulin**

Data for combination therapy with insulin are limited. In patients not adequately controlled with the maximum dosage of glimepiride, concomitant insulin therapy can be initiated. In two studies, the combination achieved the same improvement in metabolic control as insulin alone; however, a lower average dose of insulin was required in combination therapy.

## **5.2 Pharmacokinetic properties**

**Absorption:** The bioavailability of glimepiride after oral administration is complete. Food intake has no relevant influence on absorption, only absorption rate is slightly diminished. Maximum serum concentrations ( $C_{max}$ ) are reached approx. 2.5 hours after oral intake (mean 0.3 µg/ml during multiple dosing of 4 mg daily) and there is a linear relationship between dose and both  $C_{max}$  and AUC (area under the time/concentration curve).

**Distribution:** Glimepiride has a very low distribution volume (approx. 8.8 litres) which is roughly equal to the albumin distribution space, high protein binding (>99%), and a low clearance (approx. 48 ml/min).

In animals, glimepiride is excreted in milk. Glimepiride is transferred to the placenta. Passage through the blood brain barrier is low.

**Biotransformation and elimination:** Mean dominant serum half-life, which is of relevance for the serum concentrations under multiple-dose conditions, is about 5 to 8 hours. After high doses, slightly longer half-lives were noted.

After a single dose of radiolabelled glimepiride, 58% of the radioactivity was recovered in the urine, and 35% in the faeces. No unchanged substance was detected in the urine. Two metabolites - most probably resulting from hepatic metabolism - were identified both in urine and faeces: the hydroxy derivative and the carboxy derivative. After oral administration of glimepiride, the terminal half-lives of these metabolites were 3 to 6 and 5 to 6 hours respectively.

Comparison of single and multiple once-daily dosing revealed no significant differences in pharmacokinetics, and the intraindividual variability was very low. There was no relevant accumulation.

Pharmacokinetics were similar in males and females, as well as in young and elderly (above 65 years) patients. In patients with low creatinine clearance, there was a tendency for glimepiride clearance to increase and for average serum concentrations to decrease, most probably resulting from a more rapid elimination because of lower protein binding. Renal elimination of the two metabolites was impaired. Overall no additional risk of accumulation is to be assumed in such patients.

Pharmacokinetics in five non-diabetic patients after bile duct surgery were similar to those in healthy persons.

## **5.3 Preclinical safety data**

Preclinical effects observed occurred at exposures sufficiently in excess of the maximum human exposure as to indicate little relevance to clinical use, or were due to the pharmacodynamic action (hypoglycaemia) of the compound. This finding is based on conventional safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenicity, and reproduction toxicity studies. In the latter (covering embryotoxicity, teratogenicity and developmental toxicity), adverse effects observed were considered to be secondary to the hypoglycaemic effects induced by the compound in dams and in offspring.

# **6 PHARMACEUTICAL PARTICULARS**

## **6.1 List of excipients**

Lactose monohydrate  
Microcrystalline cellulose  
Sodium starch glycolate Type A  
Povidone

Iron oxide red (E172)  
Magnesium stearate

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf Life**

2 years.

## **6.4 Special precautions for storage**

Do not store above 30°C.

## **6.5 Nature and contents of container**

PVC/Al Blister Strip or PP Tablet Container with LDPE Cap.

Pack sizes: 7, 10, 14, 20, 21, 28, 30, 50, 56, 60, 84, 90, 100, 120 or 500 tablets.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product**

No special requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Medis ehf,  
Reykjavíkurvegur 78,  
220 Hafnarfjörður,  
Iceland

## **8 MARKETING AUTHORISATION NUMBER**

PA 1285/16/1

## **9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 22<sup>nd</sup> July 2005

## **10 DATE OF REVISION OF THE TEXT**

September 2007