

Package leaflet: Information for the user

RIZATRIPTAN MSD 5 mg Oral Lyophilisate rizatriptan RIZATRIPTAN MSD 10 mg Oral Lyophilisate rizatriptan

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What RIZATRIPTAN MSD is and what it is used for
2. What you need to know before you take RIZATRIPTAN MSD
3. How to take RIZATRIPTAN MSD
4. Possible side effects
5. How to store RIZATRIPTAN MSD
6. Contents of the pack and other information

1. What RIZATRIPTAN MSD is and what it is used for

RIZATRIPTAN MSD belongs to a class of medicines called selective serotonin 5-HT_{1B/1D} receptor agonists.

RIZATRIPTAN MSD is used to treat the headache phase of the migraine attack in adults.

Treatment with RIZATRIPTAN MSD:

Reduces swelling of blood vessels surrounding the brain. This swelling results in the headache pain of a migraine attack.

2. What you need to know before you take RIZATRIPTAN MSD

Do not take RIZATRIPTAN MSD if:

- you are allergic to rizatriptan benzoate or any of the other ingredients of this medicine (listed in section 6)
- you have moderately severe or severe high blood pressure, or mild high blood pressure that is not controlled by medication
- you have or have ever had heart problems including heart attack or pain on the chest (angina) or you have experienced heart disease related signs
- you have severe liver or severe kidney problems
- you have had a stroke (cerebrovascular accident CVA) or mini stroke (transient ischaemic attack TIA)
- you have blockage problems with your arteries (peripheral vascular disease)
- you are taking monoamine oxidase (MAO) inhibitors such as moclobemide, phenelzine, tranylcypromine, or pargyline (drugs against depression), or linezolid (an antibiotic), or if it has been less than two weeks since you stopped taking MAO inhibitors
- you are now taking ergotamine-type medications, such as ergotamine or dihydro-ergotamine to treat your migraine or methysergide to prevent a migraine attack

- you are taking any other drug in the same class, such as sumatriptan, naratriptan, or zolmitriptan to treat your migraine (see **Other medicines and RIZATRIPTAN MSD** below).

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking RIZATRIPTAN MSD.

Warnings and precautions

Talk to your doctor or pharmacist before taking RIZATRIPTAN MSD, if:

- you have any of the following risk factors for heart disease: high blood pressure, diabetes, you smoke or you are using nicotine substitution, your family has a history of heart disease, you are a man over 40 years of age, or you are a postmenopausal woman
- you have kidney or liver problems
- you have a particular problem with the way your heart beats (bundle branch block)
- you have or have had any allergies
- your headache is associated with dizziness, difficulty in walking, lack of coordination or weakness in the leg and arm
- you use herbal preparation containing St. John's wort
- you have had allergic reaction like swelling of face, lips, tongue and/or throat which may cause difficulty breathing and/or swallowing (angioedema).
- you are taking selective serotonin reuptake inhibitors (SSRIs) such as sertraline, escitalopram oxalate, and fluoxetine or serotonin norepinephrine reuptake inhibitors (SNRIs) such as venlafaxine, and duloxetine for depression.
- you have had short lived symptoms including chest pain and tightness.

If you take RIZATRIPTAN MSD too often this may result in you getting a chronic headache. In such cases you should contact your doctor as you may have to stop taking RIZATRIPTAN MSD.

Tell your doctor or pharmacist about your symptoms. Your doctor will decide if you have migraine. You should take RIZATRIPTAN MSD only for a migraine attack. RIZATRIPTAN MSD should not be used to treat headaches that might be caused by other, more serious conditions.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription. This includes herbal medicines and those you normally take for a migraine. This is because RIZATRIPTAN MSD can affect the way some medicines work. Also, other medicines can affect RIZATRIPTAN MSD.

Other medicines and RIZATRIPTAN MSD

Do not take RIZATRIPTAN MSD:

- if you are already taking a 5-HT_{1B/1D} agonist (sometimes referred to as ‘triptans’), such as sumatriptan, naratriptan or zolmitriptan.
- if you are taking a monoamine oxidase (MAO) inhibitor such as moclobemide, phenelzine, tranylcypromine, linezolid, or pargyline or if it has been less than two weeks since you stopped taking an MAO inhibitor.
- if you use ergotamine-type medications such as ergotamine or dihydro-ergotamine to treat your migraine
- if you use methysergide to prevent a migraine attack.

The above listed medicines when taken with RIZATRIPTAN MSD may increase the risk of side effects.

You should wait at least 6 hours after taking RIZATRIPTAN MSD before you take ergotamine-type medications such as ergotamine or dihydro-ergotamine or methysergide.

You should wait at least 24 hours after taking ergotamine- type medications before taking RIZATRIPTAN MSD.

Ask your doctor for instructions and the risks about taking RIZATRIPTAN MSD

- if you are taking propranolol (see section 3: **How to take RIZATRIPTAN MSD**).

- if you are taking SSRIs such as sertraline, escitalopram oxalate, and fluoxetine or SNRIs such as venlafaxine, and duloxetine for depression.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

RIZATRIPTAN MSD with food and drink

RIZATRIPTAN MSD can take longer to work if it is taken after food. Although it is better to take it on an empty stomach, you can still take it if you have eaten.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Available data on the safety of rizatriptan when used during the first 3 months of pregnancy do not suggest an increased risk of birth defects. It is not known whether RIZATRIPTAN MSD is harmful to an unborn baby when taken by a pregnant woman after the first 3 months of pregnancy.

If you are breastfeeding, you may postpone breastfeeding for 12 hours after treatment to avoid exposure in your baby.

Children and adolescents

The use of RIZATRIPTAN MSD oral lyophilisates in children under 18 years of age is not recommended.

Use in patients older than 65 years

There have been no full studies to look at how safe and effective RIZATRIPTAN MSD is amongst patients older than 65 years.

Driving or using machines

You may feel sleepy or dizzy while taking RIZATRIPTAN MSD. If this happens, do not drive or use any tools or machines.

RIZATRIPTAN MSD contains aspartame

This medicine contains 1.88 mg aspartame in each 5 mg oral lyophilisate which is equivalent to 1.1 mg phenylalanine. This medicine contains 3.75 mg aspartame in each 10 mg oral lyophilisate which is equivalent to 2.1 mg phenylalanine.

Aspartame is a source of phenylalanine. It may be harmful if you have phenylketonuria (PKU), a rare genetic disorder in which phenylalanine builds up because the body cannot remove it properly.

3. How to take RIZATRIPTAN MSD

RIZATRIPTAN MSD is used to treat migraine attacks. Take RIZATRIPTAN MSD as soon as possible after your migraine headache has started. Do not use it to prevent an attack.

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual dose is 10 mg.

If you are currently taking propranolol or have kidney or liver problems you should use the 5-mg dose of RIZATRIPTAN MSD. You should leave at least 2 hours between taking propranolol and RIZATRIPTAN MSD up to a maximum of 2 doses in a 24-hour period.

If migraine returns within 24 hours

In some patients, migraine symptoms can return within a 24-hour period. If your migraine does return you can take an additional dose of RIZATRIPTAN MSD. You should always wait at least 2 hours between doses.

If after 2 hours you still have a migraine

If you do not respond to the first dose of RIZATRIPTAN MSD during an attack, you should not take a second dose of RIZATRIPTAN MSD for treatment of the same attack. It is still likely, however, that you will respond to RIZATRIPTAN MSD during the next attack.

Do not take more than 2 doses of RIZATRIPTAN MSD in a 24-hour period, (for example, do not take more than two 10-mg or 5-mg oral lyophilisates or tablets in a 24-hour period). You should always wait at least 2 hours between doses.

If your condition worsens, seek medical attention.

How to administer RIZATRIPTAN MSD oral lyophilisates

- RIZATRIPTAN MSD (rizatriptan benzoate) is available as a 5- or 10-mg oral lyophilisate that dissolves in the mouth.
- Open the 'RIZATRIPTAN MSD' oral lyophilisate blister pack with dry hands.
- The oral lyophilisate should be placed on your tongue, where it dissolves and can be swallowed with the saliva.
- The oral lyophilisate can be used in situations in which liquids are not available, or to avoid the nausea and vomiting that may accompany the ingestion of tablets with liquids.

RIZATRIPTAN MSD is also available as a tablet to be taken with liquids.

If you take more RIZATRIPTAN MSD than you should:

If you take more RIZATRIPTAN MSD than you should, talk to your doctor or pharmacist straight away. Take the medicine pack with you.

Signs of overdosage can include dizziness, drowsiness, vomiting, fainting and slow heart rate.

If you have any further questions on the use of this product ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects may happen with this medicine.

In adult studies, the most common side effects reported were dizziness, sleepiness and tiredness.

Common (affects 1 to 10 users in 100)

- tingling (paraesthesia), headache, decreased sensitivity of skin (hypoesthesia), decreased mental sharpness, insomnia.
- fast or irregular heart beat (palpitation).
- flushing (redness of the face lasting a short time).
- throat discomfort.
- feeling sick (nausea), dry mouth, vomiting, diarrhoea, indigestion (dyspepsia).
- feeling of heaviness in parts of the body, neck pain, stiffness.

- pain in abdomen or chest.

Uncommon (affects 1 to 10 users in 1000)

- bad taste in your mouth.
- unsteadiness when walking (ataxia), dizziness (vertigo), blurred vision, tremor, fainting (syncope).
- confusion, nervousness.
- high blood pressure (hypertension); thirst, hot flushes, sweating.
- rash; itching and lumpy rash (hives); swelling of face, lips, tongue and/or throat which may cause difficulty breathing and/or swallowing (angioedema), difficulty breathing (dyspnoea).
- feeling of tightness in parts of the body, muscle weakness.
- changes in the rhythm or rate of the heartbeat (arrhythmia); abnormalities of the electrocardiogram (a test that records the electrical activity of your heart), very fast heartbeat (tachycardia).
- facial pain; muscle pain.

Rare (affects 1 to 10 users in 10,000)

- wheezing.
- allergic reaction (hypersensitivity); sudden life-threatening allergic reaction (anaphylaxis).
- stroke (this generally occurs in patients with risk factors for heart and blood vessel disease (high blood pressure, diabetes, smoking, use of nicotine substitution, family history of heart disease or stroke, man over 40 years of age, postmenopausal women, particular problem with the way your heart beats (bundle branch block)).
- slow heartbeat (bradycardia).

Not known (frequency cannot be estimated from the available data):

- heart attack, spasm of the blood vessels of the heart (these generally occur in patients with risk factors for heart and blood vessel disease (high blood pressure, diabetes, smoking, use of nicotine substitution, family history of heart disease or stroke, man over 40 years of age, postmenopausal women, particular problem with the way your heart beats (bundle branch block)).
- a syndrome called "serotonin syndrome" that may cause side effects like coma, unstable blood pressure, extremely high body temperature, lack of muscle coordination, agitation, and hallucinations.
- severe shedding of the skin with or without fever (toxic epidermal necrolysis).
- seizure (convulsions/fits).
- spasm of blood vessels of the extremities including coldness and numbness of the hands or feet.
- spasm of the blood vessels of the colon (large bowel), which can cause abdominal pain.

Tell your doctor right away if you have symptoms of allergic reactions, serotonin syndrome, heart attack or stroke.

In addition, tell your doctor if you experience any symptoms that suggest an allergic reaction (such as a rash or itching) after taking RIZATRIPTAN MSD.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, website: www.HPRA.ie. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store RIZATRIPTAN MSD

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton/blister after "EXP". The expiry date refers to the last day of the month.

Do not store RIZATRIPTAN MSD oral lyophilisates above 30°C.

Store in the original packaging in order to protect from moisture.

Do not remove the oral lyophilisate blister from the outer aluminium sachet until you are ready to take the medicine inside. Do not use the medicine if you notice that the aluminium sachet is damaged.

Always keep the aluminium sachets in the carrying case.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What RIZATRIPTAN MSD contains

RIZATRIPTAN MSD 5 mg oral lyophilisates

The active substance of RIZATRIPTAN MSD is rizatriptan. One oral lyophilisate contains 5 mg rizatriptan as 7.265 mg of rizatriptan benzoate.

RIZATRIPTAN MSD 10 mg oral lyophilisates

The active substance of RIZATRIPTAN MSD is rizatriptan. One oral lyophilisate contains 10 mg rizatriptan as 14.53 mg of rizatriptan benzoate.

The other ingredients of RIZATRIPTAN MSD oral lyophilisates are: gelatin, mannitol (E421), glycine, aspartame (E951), and peppermint flavour (composed of peppermint oil, maltodextrin, and dextrin).

What RIZATRIPTAN MSD looks like and contents of pack

RIZATRIPTAN MSD 5 mg oral lyophilisates

5 mg oral lyophilisates are white to off-white, round with a modified triangle on one side, with a peppermint flavour.

RIZATRIPTAN MSD 10 mg oral lyophilisates

10 mg oral lyophilisates are white to off-white, round with a modified square on one side, with a peppermint flavour.

Pack sizes: Packs with 2, 3, 6, 12 or 18 oral lyophilisates

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Organon Pharma (Ireland) Limited
2 Dublin Landings
North Wall Quay - North Dock
Dublin, D01 V4A3
Ireland.

Manufacturer

MERCK SHARP & DOHME B.V.
Waarderweg 39
2031 BN Haarlem
The Netherlands

N.V. Organon
Kloosterstraat 6
5349 AB Oss
The Netherlands

This medicinal product is authorised in the Member States of the EEA under the following names:

MAXALT 5 mg Oral Lyophilisate

Austria, Finland, Sweden
Belgium, Luxemburg
Denmark
Germany
Greece
Iceland
Ireland
Italy
Norway
Romania
The Netherlands

MAXALT RAPITAB
MAXALT LYO 5 mg
MAXALT Smelt, smeltetabletter
MAXALT lingua 5 mg Schmelztabletten
MAXALT Rapid Sol. Tab.
MAXALT SMELT 5 mg frostþurrkaðar töflur
Rizatriptan MSD 5 mg oral lyophilisates
MAXALT RPD 5 mg liofilizzato orale
Maxalt Rapitab
Maxalt 5mg liofilizat oral
MAXALT SMELT

MAXALT 10 mg Oral Lyophilisate

Austria, Finland, Sweden
Belgium, Luxemburg
Denmark
France
Germany
Greece
Iceland
Ireland
Italy
Lithuania
Norway
Romania
Spain
The Netherlands
United Kingdom

MAXALT RAPITAB
MAXALT LYO 10 mg
MAXALT Smelt, smeltetabletter
MAXALTLYO
MAXALT lingua 10 mg Schmelztabletten
MAXALT Rapid Sol. Tab.
MAXALT SMELT 10 mg frostþurrkaðar töflur
Rizatriptan MSD 10 mg oral lyophilisates
MAXALT RPD 10 mg liofilizzato orale
MAXALT 10 mg geriamieji liofilizatai
Maxalt Rapitab
Maxalt 10mg liofilizat oral
MAXALT MAX 10 mg
MAXALT SMELT
MAXALT MELT

This leaflet was last revised in June 2025

©2025 Organon group of companies. . All rights reserved.