

PACKAGE LEAFLET: INFORMATION FOR THE USER

FINASTERIDE® 5 mg Film-coated Tablets (finasteride)

This medicine is for use in men only

Read all of this leaflet carefully before you start to take this medicine.

- Keep this leaflet. You may want to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Finasteride is and what it is used for
2. Before you take Finasteride
3. How to take Finasteride
4. Possible side effects
5. How to store Finasteride
6. Further information

1. What Finasteride is and what it is used for

Finasteride contains a medicine called finasteride. This belongs to a group of medicines called '5-alpha reductase inhibitors'.

Finasteride shrinks the prostate gland in men when it is swollen. The prostate gland is found underneath the bladder (but only in men). It produces the fluid found in semen. A swollen prostate gland can lead to a condition called 'benign prostatic hyperplasia' or BPH.

What is BPH?

If you have BPH it means that your prostate gland is swollen. It can press on the tube that urine passes through, on its way out of your body.

This can lead to problems such as:

- feeling like you need to pass urine more often, especially at night
- feeling that you must pass urine right away
- finding it difficult to start passing urine
- when you pass urine the flow of urine is weak
- when you pass urine the flow stops and starts
- feeling that you cannot empty your bladder completely.

In some men, BPH can lead to more serious problems, such as:

- urinary tract infections
- a sudden inability to pass urine
- the need for surgery.

What else should you know about BPH?

- BPH is **not** cancer and does not lead to cancer, but the two conditions can be present at the same time.
- Before you start Finasteride, your doctor will do some simple tests to check whether you have prostate cancer.

Talk to your doctor if you have any questions about this.

2. Before you take Finasteride**Do not take Finasteride if:**

- you are a woman (because this medicine is for men)
- you are allergic (hypersensitive) to finasteride or any of the other ingredients (listed in Section 6).

Do not take Finasteride if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist.

Finasteride should not be used in children.

Take special care with Finasteride

Check with your doctor or pharmacist before taking your medicine if:

- your partner is pregnant or planning to become pregnant. You should use a condom or other barrier method of contraception when taking Finasteride. This is because your semen could contain a tiny amount of the drug and may affect normal development of the baby's sex organs.
- you are going to have a blood test called PSA. This is because Finasteride can affect the results of this test.

If you are not sure, talk to your doctor or pharmacist before taking Finasteride.

Taking other medicines

Finasteride does not usually affect other medicines. Always tell your doctor about all medicines you are taking or planning to take, including any obtained without a prescription.

Taking Finasteride with food and drink

Finasteride can be taken with or without food.

Pregnancy and breast feeding

- Finasteride should not be taken by women.
- Do not touch crushed or broken Finasteride tablets if you are a woman who is pregnant or planning to become pregnant (whole tablets are coated to stop contact with the medicine during normal use). This is because this medicine may affect the normal development of the baby's sex organs.
- If a woman who is pregnant comes into contact with crushed or broken Finasteride tablets, speak to your doctor.

Driving and using machines

Finasteride is not likely to affect you being able to drive, use tools or machines.

Important information about some of the ingredients of Finasteride

Finasteride contains lactose which is a type of sugar. If you have ever been told by your doctor that you cannot tolerate or digest some sugars (have an intolerance to some sugars), talk to your doctor before taking this medicine.

3. How to take Finasteride

You should take your tablets exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Taking this medicine

- The usual dose is one tablet each day.
- Take this medicine by mouth.
- Your doctor may prescribe Finasteride along with another medicine (called doxazosin) to help control your BPH.

If you take more Finasteride than you should

If you take too many tablets by mistake, contact your doctor immediately.

If you forget to take Finasteride

- If you forget to take a tablet, skip the missed dose.
- Take the next dose as usual.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Finasteride

Your condition may show an early improvement after taking Finasteride. However, it may take at least six months for the full effect to develop. It is important to keep taking Finasteride for as long as your doctor tells you, even if you do not feel any benefit straight away.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Finasteride can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

Allergic reactions

If you have an allergic reaction, stop taking Finasteride and see your doctor straight away. The signs may include:

- Skin rash, itching, or lumps under the skin (hives)
- Swelling of your lips, tongue, throat and face

Other side effects may include:

- you may be unable to have an erection (impotence)
- you may have less desire to have sex (decreased libido)
- you may have problems with ejaculation, for example a decrease in the amount of semen released during sex. This decrease in the amount of semen does not appear to affect normal sexual function.

Other side effects reported in some men are:

- Breast swelling or tenderness
- Heart palpitations
- Pain in your testicles
- An inability to have an erection which may continue after stopping the medication
- Problems with ejaculation that continued after stopping medication.
- Male infertility and/or poor quality of semen. Improvement in the quality of semen has been reported after stopping the medication
- Decrease in sex drive that continued after stopping the medication
- Depression

You should promptly report to your doctor any changes in your breast tissue such as lumps, pain, enlargement or nipple discharge as these may be signs of a serious condition, such as breast cancer.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects, you can help provide more information on the safety of this medicine.

IMB Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2

Tel: +353 1 6764971. Fax: +353 1 6762517, Website: www.imb.ie

e-mail: imbpharmacovigilance@imb.ie

5. How to store Finasteride

Keep your tablets out of the sight and reach of children.

Do not store your tablets above 30°C. Keep them in the original package.

If you have been given a calendar pack, do not remove the tablets from the blister until you are ready to take the medicine.

Do not use Finasteride after the expiry date which is stated on the carton after the letters EXP. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information

What Finasteride contains

- The active ingredient in Finasteride Tablets is finasteride. Each tablet contains 5 mg finasteride.
- The other ingredients in Finasteride tablets are: Docusate sodium, Hydroxypropyl cellulose E463, Hypromellose E464, Indigo carmine aluminium lake E132, Lactose monohydrate, Magnesium stearate E572, Microcrystalline cellulose E460, Pregelatinised maize starch, Sodium starch glycollate Type A, Talc, Titanium dioxide E171, Yellow iron oxide E172.

What Finasteride looks like and the content of the pack

Finasteride is available as blue, apple-shaped tablets. Finasteride Tablets are supplied in blister packs of 28 tablets.

This leaflet gives you some important patient information about Finasteride. If you have questions after you read it ask your doctor or pharmacist, who will give you further information.

Marketing Authorisation Holder and Manufacturer

The Marketing Authorisation Holder

Merck Sharp & Dohme Ireland (Human Health) Limited
Red Oak North
South County Business Park
Leopardstown
Dublin 18
Ireland

Manufacturer

Merck Sharp & Dohme Limited
Shotton Lane
Cramlington

Northumberland NE23 3JU
United Kingdom

This leaflet was last approved in

© Merck Sharp & Dohme Limited 2014. All rights reserved.

MSD Logo