

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Clorom XL 500 mg Prolonged-Release Tablets

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated prolonged release tablet contains;  
Clarithromycin citrate equivalent to clarithromycin 500 mg.

### Excipient with known effect:

Each tablet contains 298 mg of lactose monohydrate

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Prolonged-release tablet.

Yellow coloured, film-coated, oblong shaped, biconvex tablet, with both sides plain.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

Clorom XL 500 mg Tablets are indicated in adults and children 12 years and older.

Clorom XL 500 mg Tablets are indicated for treatment of infections caused by susceptible organisms. Indications include: Lower respiratory tract infections for example, acute and chronic bronchitis, and pneumonia (see section 4.4 and 5.1 regarding Sensitivity Testing).

Upper respiratory tract infections for example, sinusitis and pharyngitis.

Clorom XL 500 mg Tablets are also indicated in skin and soft tissue infections of mild to moderate severity, for example folliculitis, cellulitis and erysipelas (see section 4.4 and 5.1 regarding Sensitivity Testing).

### 4.2 Posology and method of administration

#### Posology

##### *Adults*

The usual recommended dosage of Clorom XL Tablets in adults is one 500 mg modified-release tablet daily to be taken with food. In more severe infections, the dosage can be increased to two 500 mg modified-release tablets daily. The usual duration of treatment is 6 to 14 days.

##### *Children older than 12 years*

As for adults.

Do not crush or chew Clorom XL 500 mg Tablets.

Clorom XL 500 mg should not be used in patients with significant renal impairment (creatinine clearance less than 30 mL/min), as appropriate clarithromycin dosage reduction is not possible when administering this product. Clarithromycin immediate release tablets may be utilized in this patient population (see section 4.3). The dosage of clarithromycin should be reduced by one-half, i.e. 250 mg once daily, or 250 mg twice daily in more severe infections. Treatment should not be continued beyond 14 days in these patients. For patients with moderate renal function (creatinine clearance 30 to 60 ml/min), a 50% dosage reduction should be implemented resulting in a maximum dose of one Clorom XL 500 mg Tablets per day.

##### *Children younger than 12 years*

Use of Clorom XL Tablets is not recommended for children younger than 12 years. Clinical trials have been conducted using clarithromycin paediatric suspension in children 6 months to 12 years of age. Therefore, children under 12 years of age should use clarithromycin paediatric suspension (granules for oral suspension).

#### Method of administration

For oral use.

### **4.3 Contraindications**

Hypersensitivity to macrolide antibiotic drugs or any of the excipients listed in section 6.1.

Concomitant administration of clarithromycin and any of the following drugs is contraindicated: astemizole, cisapride, domperidone, pimozone and terfenadine as this may result in QT prolongation and cardiac arrhythmias, including ventricular tachycardia, ventricular fibrillation, and torsades de pointe (see sections 4.4 and 4.5).

Concomitant administration with ticagrelor, ivabradine or ranolazine is contraindicated.

Concomitant administration of clarithromycin and ergot alkaloids (e.g. ergotamine or dihydroergotamine) is contraindicated, as this may result in ergot toxicity (see section 4.5).

Concomitant administration of clarithromycin and oral midazolam is contraindicated (see section 4.5).

Concomitant administration of clarithromycin and lomitapide is contraindicated (see section 4.5).

Clarithromycin must not be given to patients with electrolyte disturbances (hypokalaemia or hypomagnesaemia, due to the risk of prolongation of the QT interval) (see section 4.4).

Clarithromycin must not be given to patients with history of QT prolongation (congenital or documented acquired QT prolongation) or ventricular cardiac arrhythmia, including torsades de pointe (see sections 4.4 and 4.5).

Clarithromycin must not be used concomitantly with HMG-CoA reductase inhibitors (statins), that are extensively metabolised by CYP3A4 (lovastatin or simvastatin), due to the increased risk of myopathy including rhabdomyolysis (see section 4.4 and 4.5).

Clarithromycin is contraindicated in patients who suffer from severe hepatic failure in combination with renal impairment.

As with other strong CYP3A4 inhibitors, Clarithromycin must not be used in patients taking colchicine (see sections 4.4 and 4.5).

### **4.4 Special warnings and precautions for use**

The physician should not prescribe clarithromycin to pregnant women without carefully weighing the benefits against risk; particularly during the first three months of pregnancy (see section 4.6).

Caution is advised in patients with severe renal insufficiency (see section 4.2).

Clarithromycin is principally metabolised by the liver. Therefore, caution should be exercised in administering this antibiotic to patients with impaired hepatic function.

Caution should also be exercised when administering clarithromycin to patients with moderate to severe renal impairment (see also section 4.3).

Hepatic dysfunction, including increased liver enzymes, and hepatocellular and/or cholestatic hepatitis, with or without jaundice, has been reported with clarithromycin. This hepatic dysfunction may be severe and is usually reversible. Cases of fatal hepatic failure (see section 4.8) have been reported. Some patients may have had pre-existing hepatic disease or may have been taking other hepatotoxic medicinal products. Patients should be advised to stop treatment and contact their doctor if signs and symptoms of hepatic disease develop, such as anorexia, jaundice, dark urine, pruritus, or tender abdomen.

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including macrolides, and may range in severity from mild to life-threatening.

*Clostridium difficile*- associated diarrhoea (CDAD) has been reported with use of nearly all antibacterial agents including clarithromycin, and may range in severity from mild diarrhoea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon, which may lead to overgrowth of *C. difficile*. CDAD must be considered in all patients who present with diarrhoea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents. Therefore, discontinuation of clarithromycin therapy should be considered regardless of the indication.

Microbial testing should be performed and adequate treatment initiated. Drugs inhibiting peristalsis should be avoided.

There have been post-marketing reports of colchicine toxicity with concomitant use of clarithromycin and colchicine, especially in elderly and/or patients with renal insufficiency, some with a fatal outcome (see sections 4.5 and 4.8). Concomitant administration of clarithromycin and colchicine is contraindicated (see section 4.3).

Caution is advised regarding concomitant administration of clarithromycin and triazolobenzodiazepines, such as triazolam, and intravenous or buccal (oromucosal) midazolam (see section 4.5).

Caution is advised regarding concomitant administration of clarithromycin with other ototoxic drugs, especially with aminoglycosides. Monitoring of vestibular and auditory function should be carried out during and after treatment.

#### Cardiovascular Events:

Prolongation of the QT interval, reflecting effects on cardiac repolarisation imparting a risk of developing cardiac arrhythmia and torsades de pointes, have been seen in patients treated with macrolides including clarithromycin (see section 4.8). Due to increased risk of QT prolongation and ventricular arrhythmias (including torsades de pointes), the use of clarithromycin is contraindicated: in patients taking any of astemizole, cisapride, domperidone, pimozone and terfenadine; in patients with electrolyte disturbances such as hypomagnesaemia or hypokalaemia; and in patients with a history of QT prolongation or ventricular cardiac arrhythmia (see section 4.3).

Furthermore, clarithromycin should be used with caution in the following:

- Patients with coronary artery disease, severe cardiac insufficiency, conduction disturbances or clinically relevant bradycardia:
- Patients concomitantly taking other medicinal products associated with QT prolongation other than those which are contraindicated

Epidemiological studies investigating the risk of adverse cardiovascular outcomes with macrolides have shown variable results. Some observational studies have identified a rare short-term risk of arrhythmia, myocardial infarction and cardiovascular mortality associated with macrolides including clarithromycin. Consideration of these findings should be balanced with treatment benefits when prescribing clarithromycin.

#### Pneumonia:

In view of the emerging resistance of *Streptococcus pneumoniae* to macrolides, it is important that sensitivity testing be performed when prescribing clarithromycin for community-acquired pneumonia. In hospital-acquired pneumonia, clarithromycin should be used in combination with additional appropriate antibiotics.

#### Skin and soft tissue infections of mild to moderate severity:

These infections are most often caused by *Staphylococcus aureus* and *Streptococcus pyogenes*, both of which may be resistant to macrolides. Therefore, it is important that sensitivity testing be performed. In cases where *beta*-lactam antibiotics cannot be used (e.g. allergy), other antibiotics, such as clindamycin, may be the drug of first choice. Currently, macrolides are only considered to play a role in some skin and soft tissue infections, such as those caused by *Corynebacterium minutissimum*, acne vulgaris, and erysipelas and in situations where penicillin treatment cannot be used.

In the event of severe acute hypersensitivity reactions, such as anaphylaxis, severe cutaneous adverse reactions (SCAR) (e.g. acute generalised exanthematous pustulosis (AGEP), Stevens-Johnson Syndrome, toxic epidermal necrolysis, and drug reaction with eosinophilia and systemic symptoms (DRESS)), clarithromycin therapy should be discontinued immediately and appropriate treatment should be urgently initiated.

Clarithromycin should be used with caution when administered concurrently with medications that induce the cytochrome CYP3A4 enzyme (see section 4.5).

#### HMG-CoA Reductase Inhibitors (statins):

Concomitant use of clarithromycin with lovastatin or simvastatin is contraindicated (see section 4.3). Caution should be exercised when prescribing clarithromycin with other statins.

Rhabdomyolysis has been reported in patients taking clarithromycin and statins. Patients should be monitored for signs and symptoms of myopathy.

In situations where the concomitant use of clarithromycin with statins cannot be avoided, it is recommended to prescribe the lowest registered dose of the statin. Use of a statin that is not dependent on CYP3A metabolism (e.g. fluvastatin) can be considered (see section 4.5).

Oral hypoglycaemic agents/Insulin:

The concomitant use of clarithromycin and oral hypoglycaemic agents (such as sulphonylurias) and/or insulin can result in significant hypoglycaemia. Careful monitoring of glucose is recommended (see section 4.5).

Oral anticoagulants:

There is a risk of serious haemorrhage and significant elevations in International Normalized Ratio (INR) and prothrombin time when clarithromycin is co-administered with warfarin (see section 4.5). INR and prothrombin times should be frequently monitored while patients are receiving clarithromycin and oral anticoagulants concurrently.

Caution should be exercised when clarithromycin is co-administered with direct acting oral anticoagulants such as dabigatran, rivaroxaban apixaban and edoxaban, particularly to patients at high risk of bleeding (see section 4.5).

Long-term use may, as with other antibiotics, result in colonisation with increased numbers of non-susceptible bacteria and fungi. If superinfections occur, appropriate therapy should be instituted.

Attention should also be paid to the possibility of cross resistance between clarithromycin and other macrolide drugs, as well as lincomycin and clindamycin.

Excipients:

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

**4.5 Interaction with other medicinal products and other forms of interaction**

**The use of the following drugs is strictly contraindicated due to the potential for severe drug interaction effects:**

Astemizole, Cisapride, domperidone, pimozone and terfenadine:

Elevated cisapride levels have been reported in patients receiving clarithromycin and cisapride concomitantly. This may result in QT prolongation and cardiac arrhythmias including ventricular tachycardia, ventricular fibrillation and torsades de pointes. Similar effects have been observed in patients taking clarithromycin and pimozone concomitantly (see section 4.3).

Macrolides have been reported to alter the metabolism of terfenadine resulting in increased levels of terfenadine which has occasionally been associated with cardiac arrhythmias, such as QT prolongation, ventricular tachycardia, ventricular fibrillation and torsades de pointes (see section 4.3). In one study in 14 healthy volunteers, the concomitant administration of clarithromycin and terfenadine resulted in 2- to 3-fold increase in the serum level of the acid metabolite of terfenadine and in prolongation of the QT interval which did not lead to any clinically detectable effect. Similar effects have been observed with concomitant administration of astemizole and other macrolides.

Ergot alkaloids:

Post-marketing reports indicate that co-administration of clarithromycin with ergotamine or dihydroergotamine has been associated with acute ergot toxicity characterized by vasospasm, and ischaemia of the extremities and other tissues including the central nervous system. Concomitant administration of clarithromycin and these medicinal products is contraindicated (see section 4.3).

Oral Midazolam

When midazolam was co-administered with clarithromycin tablets (500 mg twice daily), midazolam AUC was increased 7-fold after oral administration of midazolam. Concomitant administration of oral midazolam and clarithromycin is contraindicated (see section 4.3).

HMG-CoA Reductase Inhibitors (statins)

Concomitant use of clarithromycin with lovastatin or simvastatin is contraindicated (see 4.3) as these statins are extensively metabolized by CYP3A4 and concomitant treatment with clarithromycin increases their plasma concentration, which increases the risk of myopathy, including rhabdomyolysis. Reports of rhabdomyolysis have been received for patients taking clarithromycin concomitantly with these statins. If treatment with clarithromycin cannot be avoided, therapy with lovastatin or simvastatin must be suspended during the course of treatment.

Caution should be exercised when prescribing clarithromycin with statins. In situations where the concomitant use of clarithromycin with statins cannot be avoided, it is recommended to prescribe the lowest registered dose of the statin. Use of a statin that is not dependent on CYP3A metabolism (e.g. fluvastatin) can be considered. Patients should be monitored for signs and symptoms of myopathy.

### **Effects of Other Medicinal Products on Clarithromycin**

Drugs that are inducers of CYP3A (e.g. rifampicin, phenytoin, carbamazepine, phenobarbital, St John's wort) may induce the metabolism of clarithromycin. This may result in sub-therapeutic levels of clarithromycin leading to reduced efficacy. Furthermore, it might be necessary to monitor the plasma levels of the CYP3A inducer, which could be increased owing to the inhibition of CYP3A by clarithromycin (see also the relevant product information for the CYP3A4 inhibitor administered). Concomitant administration of rifabutin and clarithromycin resulted in an increase in rifabutin and decrease in clarithromycin serum levels together with an increased risk of uveitis.

The following drugs are known or suspected to affect circulating concentrations of clarithromycin; clarithromycin dosage adjustment or consideration of alternative treatments may be required.

#### Fluconazole

Concomitant administration of fluconazole 200 mg daily and clarithromycin 500 mg twice daily to 21 healthy volunteers led to increases in the mean steady-state minimum clarithromycin concentration ( $C_{min}$ ) and area under the curve (AUC) of 33% and 18% respectively. Steady state concentrations of the active metabolite 14-OH-clarithromycin were not significantly affected by concomitant administration of fluconazole. No clarithromycin dose adjustment is necessary.

#### Ritonavir

A pharmacokinetic study demonstrated that the concomitant administration of ritonavir 200 mg every eight hours and clarithromycin 500 mg every 12 hours resulted in a marked inhibition of the metabolism of clarithromycin. The clarithromycin  $C_{max}$  increased by 31%,  $C_{min}$  increased 182% and AUC increased by 77% with concomitant administration of ritonavir. An essentially complete inhibition of the formation of 14-OH-clarithromycin was noted. Because of the large therapeutic window for clarithromycin, no dosage reduction should be necessary in patients with normal renal function. However, for patients with renal impairment, the following dosage adjustments should be considered: For patients with  $CL_{CR}$  30 to 60 mL/min the dose of clarithromycin should be reduced by 50%. For patients with  $CL_{CR} < 30$  mL/min the dose of clarithromycin should be decreased by 75%. Doses of clarithromycin greater than 1 gm/day should not be co-administered with protease inhibitors.

Similar dose adjustments should be considered in patients with reduced renal function when ritonavir is used as a pharmacokinetic enhancer with other HIV protease inhibitors including atazanavir and saquinavir (see section below, bidirectional pharmacokinetic interactions).

#### Efavirenz, nevirapine, rifampicin, rifabutin and rifapentine

Strong inducers of the cytochrome P450 metabolism system such as efavirenz, nevirapine, rifampicin, rifabutin, and rifapentine may accelerate the metabolism of clarithromycin and thus lower the plasma levels of clarithromycin, while increasing those of 14-OH-clarithromycin, a metabolite that is also microbiologically active. Since the microbiological activities of clarithromycin and 14-OH-clarithromycin are different for different bacteria, the intended therapeutic effect could be impaired during concomitant administration of clarithromycin and enzyme inducers.

#### Etravirine

Clarithromycin exposure was decreased by etravirine; however, concentrations of the active metabolite, 14-OH-clarithromycin, were increased. Because 14-OH-clarithromycin has reduced activity against Mycobacterium avium complex (MAC), overall activity against this pathogen may be altered; therefore alternatives to clarithromycin should be considered for the treatment of MAC.

### **Effect of Clarithromycin on Other Medicinal Products**

#### CYP3A-based interactions

Co-administration of clarithromycin, known to inhibit CYP3A, and a drug primarily metabolised by CYP3A may be associated with elevations in drug concentrations that could increase or prolong both therapeutic and adverse effects of the concomitant drug.

The use of clarithromycin is contraindicated in patients receiving the CYP3A substrates astemizole, cisapride, domperidone, pimozide and terfenadine due to the risk of QT prolongation and cardiac arrhythmias, including ventricular tachycardia, ventricular fibrillation, and torsades de pointes (see sections 4.3 and 4.4).

The use of clarithromycin is also contraindicated with ergot alkaloids, oral midazolam, HMG CoA reductase inhibitors metabolised mainly by CYP3A4 (e.g. lovastatin and simvastatin), colchicine, ticagrelor, ivabradine and ranolazine (see section 4.3).

Concomitant administration of clarithromycin with lomitapide is contraindicated due the potential for markedly increased transaminases (see section 4.3).

Caution is required if clarithromycin is co-administered with other drugs known to be CYP3A enzyme substrates, especially if the CYP3A substrate has a narrow safety margin (e.g. carbamazepine) and/or the substrate is extensively metabolised by this enzyme. Dosage adjustments may be considered, and when possible, serum concentrations of drugs primarily metabolised by CYP3A should be monitored closely in patients concurrently receiving clarithromycin. Drugs or drug classes that are known or suspected to be metabolised by the same CYP3A isozyme include (but this list is not comprehensive) alprazolam, carbamazepine, cilostazole, ciclosporin, disopyramide, ibrutinib, methylprednisolone, midazolam (intravenous), omeprazole, oral anticoagulants (e.g. warfarin, rivaroxaban, apixaban), atypical antipsychotics (e.g. quetiapine), quinidine, rifabutin, sildenafil, sirolimus, tacrolimus, triazolam and vinblastine.

Drugs interacting by similar mechanisms through other isozymes within the cytochrome P450 system include phenytoin, theophylline and valproate.

#### Direct acting oral anticoagulants (DOACs)

The DOACs dabigatran and edoxaban are substrates for the efflux transporter P-gp. Rivaroxaban and apixaban are metabolised via CYP3A4 and are also substrates for P-gp. Caution should be exercised when clarithromycin is co-administered with these agents particularly to patients at high risk of bleeding (see section 4.4).

#### Antiarrhythmics

There have been post-marketing reports of torsade de pointes occurring with concurrent use of clarithromycin and quinidine or disopyramide. Electrocardiograms should be monitored for QT prolongation during coadministration of clarithromycin with these drugs. Serum concentrations of these medications should also be monitored.

There have been post marketing reports of hypoglycemia with the concomitant administration of clarithromycin and disopyramide. Therefore blood glucose levels should be monitored during concomitant administration of clarithromycin and disopyramide.

#### Oral hypoglycemic agents/Insulin

With certain hypoglycemic drugs such as nateglinide, and repaglinide, inhibition of CYP3A enzyme by clarithromycin may be involved and could cause hypoglycemia when used concomitantly. Careful monitoring of glucose is recommended.

#### Omeprazole

Clarithromycin (500 mg every 8 hours) was given in combination with omeprazole (40 mg daily) to healthy adult subjects. The steady-state plasma concentrations of omeprazole were increased ( $C_{max}$ ,  $AUC_{0-24}$ , and  $t_{1/2}$  increased by 30%, 89%, and 34%, respectively), by the concomitant administration of clarithromycin. The mean 24-hour gastric pH value was 5.2 when omeprazole was administered alone and 5.7 when omeprazole was co-administered with clarithromycin.

#### Sildenafil, tadalafil and vardenafil

Each of these phosphodiesterase inhibitors is metabolised, at least in part, by CYP3A, and CYP3A may be inhibited by concomitantly administered clarithromycin. Co-administration of clarithromycin with sildenafil, tadalafil or vardenafil would likely result in increased phosphodiesterase inhibitor exposure. Reduction of sildenafil, tadalafil and vardenafil dosages should be considered when these drugs are co-administered with clarithromycin.

#### Theophylline, carbamazepine

Results of clinical studies indicate that there was a modest but statistically significant ( $p \leq 0.05$ ) increase of circulating theophylline or carbamazepine levels when either of these drugs were administered concomitantly with clarithromycin. Dose reduction may need to be considered.

#### Oral anticoagulants (e.g., warfarin, acenocoumarol)

In isolated cases, patients receiving combination therapy with clarithromycin and oral anticoagulants may experience increased pharmacologic effects and even toxic effects of these drugs. International normalized ratio (INR) or Prothrombin times should be carefully monitored while patients are simultaneously receiving clarithromycin and oral anticoagulants.

#### Tolterodine

The primary route of metabolism for tolterodine is via the 2D6 isoform of cytochrome P450 (CYP2D6). However, in a subset of the population devoid of CYP2D6, the identified pathway of metabolism is via CYP3A. In this population subset, inhibition of CYP3A results in significantly higher serum concentrations of tolterodine. A reduction in tolterodine dosage may be necessary in the presence of CYP3A inhibitors, such as clarithromycin in the CYP2D6 poor metaboliser population.

#### Triazolobenzodiazepines (e.g., alprazolam, midazolam, triazolam)

When midazolam was co-administered with clarithromycin tablets (500 mg twice daily), midazolam AUC was increased 2.7-fold after intravenous administration of midazolam. If intravenous midazolam is co-administered with clarithromycin, the patient must be closely monitored to allow dose adjustment. Drug delivery of midazolam via oromucosal route, which could bypass pre-systemic elimination of the drug, will likely result in a similar interaction to that observed after intravenous midazolam rather than oral administration. The same precautions should also apply to other benzodiazepines that are metabolised by CYP3A, including triazolam and alprazolam. For benzodiazepines which are not dependent on CYP3A for their elimination (temazepam, nitrazepam, lorazepam), a clinically important interaction with clarithromycin is unlikely.

There have been post-marketing reports of drug interactions and central nervous system (CNS) effects (e.g., somnolence and confusion) with the concomitant use of clarithromycin and triazolam. Monitoring the patient for increased CNS pharmacological effects is suggested.

There are no in-vivo human data available describing an interaction between clarithromycin and the following drugs: aprepitant, eletriptan, halofantrine, and ziprasidone. However, because in vitro data suggest these drugs are CYP3A substrates, caution should be used when they are co-administered with clarithromycin.

Eletriptan should not be coadministered with CYP3A inhibitors such as clarithromycin.

There have been spontaneous or published reports of drug interactions of CYP3A inhibitors, including clarithromycin, with cyclosporine, tacrolimus, methylprednisolone, vinblastine, and cilostazol.

#### **Other drug interactions**

##### Aminoglycosides

Caution is advised regarding concomitant administration of clarithromycin with other ototoxic drugs, especially with aminoglycosides (see section 4.4).

##### Colchicine

Colchicine is a substrate for both CYP3A and the efflux transporter, P-glycoprotein (Pgp). Clarithromycin and other macrolides are known to inhibit CYP3A and Pgp. When clarithromycin and colchicine are administered together, inhibition of Pgp and/or CYP3A by clarithromycin may lead to increased exposure to colchicine (see section 4.3 and 4.4).

##### Digoxin

Digoxin is thought to be a substrate for the efflux transporter, P-glycoprotein (Pgp). Clarithromycin is known to inhibit Pgp. When clarithromycin and digoxin are administered together, inhibition of Pgp by clarithromycin may lead to increased exposure to digoxin. Elevated digoxin serum concentrations in patients receiving clarithromycin and digoxin concomitantly have also been reported in post marketing surveillance. Some patients have shown clinical signs consistent with digoxin toxicity, including potentially fatal arrhythmias. Serum digoxin concentrations should be carefully monitored while patients are receiving digoxin and clarithromycin simultaneously.

##### Zidovudine

Simultaneous oral administration of clarithromycin tablets and zidovudine to HIV-infected adult patients may result in decreased steady-state zidovudine concentrations. Because clarithromycin appears to interfere with the absorption of simultaneously administered oral zidovudine, this interaction can be largely avoided by staggering the doses of clarithromycin and zidovudine to allow for a 4-hour interval between each medication. This interaction does not appear to occur in paediatric HIV-infected patients taking clarithromycin suspension with zidovudine or dideoxyinosine. This interaction is unlikely when clarithromycin is administered via intravenous infusion.

##### Phenytoin and Valproate

There have been spontaneous or published reports of interactions of CYP3A inhibitors, including clarithromycin with drugs not thought to be metabolised by CYP3A (e.g. phenytoin and valproate). Serum level determinations are recommended for these drugs when administered concomitantly with clarithromycin. Increased serum levels have been reported.

##### Hydroxychloroquine and chloroquine

Clarithromycin should be used with caution in patients receiving these medicines known to prolong the QT interval due to the potential to induce cardiac arrhythmia and serious adverse cardiovascular events.

### Corticosteroids

Caution should be exercised in concomitant use of clarithromycin with systemic and inhaled corticosteroids that are primarily metabolised by CYP3A due to the potential for increased systemic exposure to corticosteroids. If concomitant use occurs, patients should be closely monitored for systemic corticosteroid undesirable effects.

### **Bi-directional drug interactions**

#### Atazanavir

Both clarithromycin and atazanavir are substrates and inhibitors of CYP3A, and there is evidence of a bi-directional drug interaction. Co-administration of clarithromycin (500 mg twice daily) with atazanavir (400 mg once daily) resulted in a 2-fold increase in exposure to clarithromycin and a 70% decrease in exposure to 14-OH-clarithromycin, with a 28% increase in the AUC of atazanavir. Because of the large therapeutic window for clarithromycin, no dosage reduction should be necessary in patients with normal renal function. For patients with moderate renal function (creatinine clearance 30 to 60 mL/min), the dose of clarithromycin should be decreased by 50%. For patients with creatinine clearance <30 mL/min, the dose of clarithromycin should be decreased by 75% using an appropriate clarithromycin formulation. Doses of clarithromycin greater than 1000 mg per day should not be co-administered with protease inhibitors.

#### Calcium Channel Blockers

Caution is advised regarding the concomitant administration of clarithromycin and calcium channel blockers metabolized by CYP3A4 (e.g. verapamil, amlodipine, diltiazem) due to the risk of hypotension. Plasma concentrations of clarithromycin as well as calcium channel blockers may increase due to the interaction. Hypotension, bradyarrhythmias and lactic acidosis have been observed in patients taking clarithromycin and verapamil concomitantly.

#### Itraconazole

Both clarithromycin and itraconazole are substrates and inhibitors of CYP3A, leading to a bidirectional drug interaction. Clarithromycin may increase the plasma levels of itraconazole, while itraconazole may increase the plasma levels of clarithromycin. Patients taking itraconazole and clarithromycin concomitantly should be monitored closely for signs or symptoms of increased or prolonged pharmacologic effect.

#### Saquinavir

Both clarithromycin and saquinavir are substrates and inhibitors of CYP3A, and there is evidence of a bi-directional drug interaction. Concomitant administration of clarithromycin (500 mg twice daily) and saquinavir (soft gelatin capsules, 1200 mg three times daily) to 12 healthy volunteers resulted in steady-state AUC and  $C_{max}$  values of saquinavir which were 177% and 187% higher than those seen with saquinavir alone. Clarithromycin AUC and  $C_{max}$  values were approximately 40% higher than those seen with clarithromycin alone. No dose adjustment is required when the two drugs are co-administered for a limited time at the doses/formulations studied. Observations from drug interaction studies using the soft gelatin capsule formulation may not be representative of the effects seen using the saquinavir hard gelatin capsule. Observations from drug interaction studies performed with saquinavir alone may not be representative of the effects seen with saquinavir/ritonavir therapy. When saquinavir is co-administered with ritonavir, consideration should be given to the potential effects of ritonavir on clarithromycin (see section 4.5: Ritonavir).

Patients taking oral contraceptives should be warned that if diarrhoea, vomiting or breakthrough bleeding occur there is a possibility of contraceptive failure.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

The safety of clarithromycin for use during pregnancy has not been established. Based on variable results obtained from animal studies and experience in humans, the possibility of adverse effects on embryofoetal development cannot be excluded. Some observational studies evaluating exposure to clarithromycin during the first and second trimester have reported an increased risk of miscarriage compared to no antibiotic use or other antibiotic use during the same period. The available epidemiological studies on the risk of major congenital malformations with use of macrolides including clarithromycin during pregnancy provide conflicting results.

Therefore, use during pregnancy is not advised without carefully weighing the benefits against risk

### Breast-feeding

The safety of clarithromycin for using during breast-feeding of infants has not been established. Clarithromycin is excreted into human breast milk in small amounts. It has been estimated that an exclusively breastfed infant would receive about 1.7% of the maternal weight-adjusted dose of clarithromycin.

### Fertility

In the rat, fertility studies have not shown any evidence of harmful effects (see section 5.3).

#### 4.7 Effects on ability to drive and use machines

There are no data on the effect of clarithromycin on the ability to drive or use machines. The potential for dizziness, vertigo, confusion and disorientation, which may occur with the medication, should be taken into account before patients drive or use machines.

#### 4.8 Undesirable effects

##### a. Summary of the safety profile

The most frequent and common adverse reactions related to clarithromycin therapy for both adult and paediatric populations are abdominal pain, diarrhoea, nausea, vomiting and taste perversion. These adverse reactions are usually mild in intensity and are consistent with the known safety profile of macrolide antibiotics (see section b of section 4.8).

There was no significant difference in the incidence of these gastrointestinal adverse reactions during clinical trials between the patient population with or without pre-existing mycobacterial infections.

##### b. Tabulated summary of adverse reactions

The following table displays adverse reactions reported in clinical trials and from post-marketing experience with clarithromycin immediate-release tablets, granules for oral suspension, powder for solution for injection, extended-release tablets and modified-release tablets.

The reactions considered at least possibly related to clarithromycin are displayed by system organ class and frequency using the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ) and not known (adverse reactions from post-marketing experience; cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness when the seriousness could be assessed

System Organ Class	Very common $\geq 1/10$	Common $\geq 1/100$ to $< 1/10$	Uncommon $\geq 1/1,000$ to $< 1/100$	Not Known* (cannot be estimated from the available data)
Infections and infestations			Cellulitis <sup>1</sup> , candidiasis, gastroenteritis <sup>2</sup> , infection <sup>3</sup> , vaginal infection	Pseudomembranous colitis, erysipelas,
Blood and lymphatic system			Leukopenia, neutropenia <sup>4</sup> , thrombocythaemia <sup>3</sup> , eosinophilia <sup>4</sup>	Agranulocytosis, thrombocytopenia
Immune system disorders <sup>5</sup>			Anaphylactoid reaction <sup>1</sup> , hypersensitivity	Anaphylactic reaction. angioedema
Metabolism and nutrition disorders			Anorexia, decreased appetite	
Psychiatric disorders		Insomnia	Anxiety, nervousness <sup>3</sup>	Psychotic disorder, confusional state <sup>5</sup> , depersonalisation, depression, disorientation, hallucination, abnormal dreams, mania
Nervous system disorders		Dysgeusia, headache	Loss of consciousness <sup>1</sup> , dyskinesia <sup>1</sup> ,	Convulsion, ageusia, parosmia, anosmia,

			dizziness, somnolence <sup>5</sup> , tremor	paraesthesia
Ear and labyrinth disorders			Vertigo, hearing impaired, tinnitus	Deafness
Cardiac disorders			Cardiac arrest <sup>1</sup> , atrial fibrillation <sup>1</sup> , electrocardiogram QT prolonged, extrasystoles <sup>1</sup> , palpitations	Torsades de pointes, ventricular tachycardia, ventricular fibrillation
Vascular disorders		Vasodilation <sup>1</sup>		Haemorrhage
Respiratory, thoracic and mediastinal disorder			Asthma <sup>1</sup> , epistaxis <sup>2</sup> , pulmonary embolism <sup>1</sup>	
Gastrointestinal disorders		Diarrhoea, vomiting, dyspepsia, nausea, abdominal pain	Oesophagitis <sup>1</sup> , gastroesophageal reflux disease <sup>2</sup> , gastritis, proctalgia <sup>2</sup> , stomatitis, glossitis, abdominal distension <sup>4</sup> , constipation, dry mouth, eructation, flatulence	Pancreatitis acute, tongue discolouration, tooth discolouration
Hepatobiliary disorders		Liver function test abnormal	Cholestasis <sup>4</sup> , hepatitis <sup>4</sup> , alanine aminotransferase increased, aspartate aminotransferase increased, gamma- glutamyltransferase increased <sup>4</sup>	Hepatic failure, jaundice hepatocellular
Skin and subcutaneous tissue disorders		Rash, hyperhidrosis	Dermatitis bullous <sup>1</sup> , pruritus, urticaria, rash maculo- papular <sup>3</sup>	Severe cutaneous adverse reactions (SCAR) (e.g. Acute generalised exanthematous pustulosis (AGEP),Stevens- Johnson syndrome, toxic epidermal necrolysis, drug rash with eosinophilia and systemic symptoms (DRESS)), acne
Musculoskeletal and connective tissue disorders			Muscle spasms <sup>3</sup> , musculoskeletal stiffness <sup>1</sup> , myalgia <sup>2</sup>	Rhabdomyolysis <sup>2</sup> , 6, myopathy

Renal and urinary disorders			Blood creatinine increased <sup>1</sup> , blood urea increased <sup>1</sup>	Renal failure, nephritis interstitial
General disorders and administration site conditions	Injection site phlebitis <sup>1</sup>	Injection site pain <sup>1</sup> , injection site inflammation <sup>1</sup>	Malaise <sup>4</sup> , pyrexia <sup>3</sup> , asthenia, chest pain <sup>4</sup> , chills <sup>4</sup> , fatigue <sup>4</sup>	
Investigations			Albumin globulin ratio abnormal <sup>1</sup> , blood alkaline phosphatase increased <sup>4</sup> , blood lactate dehydrogenase increased <sup>4</sup>	International normalised ratio increased, prothrombin time prolonged, urine.

\* Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. Patient exposure is estimated to be greater than 1 billion patient treatment days for clarithromycin.

1 ADRs reported only for the Powder for Concentrate for Solution for Infusion formulation

2 ADRs reported only for the Extended-Release Tablets formulation

3 ADRs reported only for the Granules for Oral Suspension formulation

4 ADRs reported only for the Immediate-Release Tablets formulation

5, 6, See section c)

#### **c. Description of selected adverse reactions**

Injection site phlebitis, injection site pain, and injection site inflammation are specific to the clarithromycin intravenous formulation.

In some of the reports of rhabdomyolysis, clarithromycin was administered concomitantly with statins, fibrates, colchicine or allopurinol (see section 4.3 and 4.4).

There have been post-marketing reports of drug interactions and central nervous system (CNS) effects (e.g. somnolence and confusion) with the concomitant use of clarithromycin and triazolam. Monitoring the patient for increased CNS pharmacological effects is suggested (see section 4.5).

There have been rare reports of clarithromycin ER tablets in the stool, many of which have occurred in patients with anatomic (including ileostomy or colostomy) or functional gastrointestinal disorders with shortened GI transit times. In several reports, tablet residues have occurred in the context of diarrhoea. It is recommended that patients who experience tablet residue in the stool and no improvement in their condition should be switched to a different clarithromycin formulation (e.g. suspension) or another antibiotic.

Special population: Adverse Reactions in Immunocompromised Patients (see section e).

#### **d. Paediatric populations**

Clinical trials have been conducted using clarithromycin paediatric suspension in children 6 months to 12 years of age. Therefore, children under 12 years of age should use clarithromycin paediatric suspension.

Frequency, type and severity of adverse reactions in children are expected to be the same as in adults.

#### **e. Other special populations**

##### *Immunocompromised patients*

In AIDS and other immunocompromised patients treated with the higher doses of clarithromycin over long periods of time for mycobacterial infections, it was often difficult to distinguish adverse events possibly associated with clarithromycin administration from underlying signs of Human Immunodeficiency Virus (HIV) disease or intercurrent illness.

In adult patients, the most frequently reported adverse reactions by patients treated with total daily doses of 1000 mg of clarithromycin were: nausea, vomiting, taste perversion, abdominal pain, diarrhoea, rash, flatulence, headache, constipation, hearing disturbance, Serum Glutamic Oxaloacetic Transaminase (SGOT) and Serum Glutamic Pyruvate Transaminase (SGPT) elevations. Additional low-frequency events included dyspnoea, insomnia and dry mouth.

In these immunocompromised patients, evaluations of laboratory values were made by analysing those values outside the seriously abnormal level (i.e. the extreme high or low limit) for the specified test. On the basis of these criteria, about 2% to 3% of those patients who received 1000mg of clarithromycin daily had seriously abnormal elevated levels of SGOT and SGPT, and abnormally low white blood cell and platelet counts. A lower percentage of patients in these two dosage groups also had elevated Blood Urea Nitrogen levels.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via:

HPRA Pharmacovigilance  
Earlsfort Terrace  
IRL - Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.hpra.ie](http://www.hpra.ie)  
e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

### 4.9 Overdose

Reports indicate that the ingestion of large amounts of clarithromycin can be expected to produce gastro-intestinal symptoms. One patient who had a history of bipolar disorder ingested 8 grams of clarithromycin and showed altered mental status, paranoid behaviour, hypokalaemia and hypoxaemia.

Adverse reactions accompanying overdosage should be treated by the prompt elimination of unabsorbed drug and supportive measures. As with other macrolides, clarithromycin serum levels are not expected to be appreciably affected by haemodialysis or peritoneal dialysis.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

#### **General Properties**

##### ATC classification

Pharmacotherapeutic group: Antibacterial for systemic use, macrolide, ATC code: J01FA09.

##### Mechanism of Action

Clarithromycin is an antibiotic belonging to the macrolide antibiotics group. It exerts its antibacterial action by inhibiting the intracellular protein synthesis of susceptible bacteria. It selectively binds to the 50S subunit of bacterial ribosomes and thus prevents the translocation of activated amino acids.

Clarithromycin has relevant bactericidal activity against several bacterial strains.

The organisms include *H. influenzae*, *S. pneumoniae*, *S. pyogenes*, *S. aureus*, *M. catarrhalis*, *H. pylori*, *C. pneumoniae*, *M. pneumoniae*, *L. pneumophila*, *M. avium*, and *M. intracellulare*.

The 14(R)-hydroxy metabolite of clarithromycin, a product of parent drug metabolism in humans, also has antimicrobial activity. The metabolite is less active than the parent compound for most organisms, including *Mycobacterium* spp. An exception is *Haemophilus influenzae* against which the metabolite is 1 to 2 times more active than the parent compound. Clarithromycin combined with the metabolite showed a strain-dependent additive or synergistic effect both in vitro and in vivo.

##### PK/PD Relationship

Clarithromycin is extensively distributed in body tissues and fluids. Because of high tissue penetration, intracellular concentrations are higher than serum concentrations.

Clarithromycin concentrations in tonsil and whole lung tissue are 2- to 6-fold higher than those observed in the serum. Tissue and serum concentrations observed in Abbott studies with immediate-release (IR) tablets are presented below.

Mean Clarithromycin Concentration [250mg BID]		
Tissue Type	Tissue	Serum
Tonsil	1.6 µg/g	0.8 µg/ml
Lung	8.8 µg/g	1.7 µg/ml

The pharmacokinetics of orally administered modified-release (MR) clarithromycin tablets have been studied in adult humans (refer to section 5.2) and compared with clarithromycin 250 mg and 500 mg IR tablets. The extent of absorption – area under curve (AUC) – was found to be equivalent when equal total daily doses were administered. The equivalent AUCs would be expected to drive tissue levels equivalent to those observed for clarithromycin IR tablets.

In a study in healthy volunteers, it was shown that the concentrations of clarithromycin in epithelial lining fluid (ELF) following administration of the MR formulation remained above 1 µg/ml for 24 hours and above 10 µg/ml for up to 18 hours. In most subjects, the concentrations of clarithromycin in ELF were approximately 30 times greater than those in plasma, and the ratio appeared to be independent of formulation and time of assessment. A peak tissue concentration above 40 µg/ml was observed for the MR formulation, demonstrating extensive uptake of clarithromycin into lung tissue. This level is well above the minimum inhibitory concentration (MIC) values of all common community-acquired respiratory pathogens.

Clarithromycin accumulated extensively in the alveolar macrophages (AM), with AM levels approximately 100- to 600-fold higher than those in plasma and 4- to 18-fold higher than those in ELF for most subjects. While concentrations of 14(R)-hydroxy-clarithromycin in AM were not quantifiable in some subjects and were rather variable, the AM levels were generally similar for the MR and IR tablets. The concentrations in AM were greater than those in plasma, but accumulation was less for the metabolite than for parent clarithromycin.

### **Mechanism of Resistance**

Acquired macrolide resistance in *S. pneumoniae*, *S. pyogenes*, and *S. aureus* is mediated primarily by the presence of one of two mechanisms (i.e. *erm* and *mef* or *msr*).

Ribosomal binding of the antimicrobial is prevented through methylation of the ribosome by an enzyme (*erm*). Alternatively an efflux mechanism (*mef* or *msr*) can prevent the antimicrobial from reaching its ribosomal target by pumping the antimicrobial out of the cell. No acquired resistance mechanisms have been identified in *Moraxella* or *Haemophilus* spp. Macrolide resistance mechanisms are equally effective against 14- and 15-membered macrolides including erythromycin, clarithromycin, roxithromycin, and azithromycin. The mechanisms for penicillin resistance and macrolide resistance are unrelated. Attention should be paid to the *erm*-mediated cross-resistance between macrolides such as clarithromycin and lincosamides such as lincomycin and clindamycin.

Clarithromycin antagonises the bacterial effects of beta-lactam antibiotics. Also the effects of lincomycin and clindamycin are antagonised, at least in vitro.

### **Breakpoints**

The following breakpoints for clarithromycin, separating susceptible organisms from resistant organisms, have been established by the European Committee for Antimicrobial Susceptibility Testing (EUCAST).

Breakpoints (MIC, µg/ml)		
Microorganism	Susceptible (≤)	Resistant (>)
<i>Staphylococcus spp.</i> <sup>1</sup>	1 mg/ml	2 mg/ml
<i>Streptococcus spp.</i> <sup>1</sup> (groups A, B, C and G)	0.25 mg/ml	0.5 mg/ml
<i>Streptococcus pneumoniae</i> <sup>1</sup>	0.25 mg/ml	0.5 mg/ml
<i>Moraxella catarrhalis</i> <sup>1</sup>	0.25 mg/ml	0.5 mg/ml
<i>Helicobacter pylori</i>	0.25 mg/ml	0.5 mg/ml
Erythromycin can be used to determine susceptibility to clarithromycin.		

The prevalence of acquired resistance rates may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of an agent in at least some types of infections is questionable.

Clarithromycin has a pronounced effect against a wide variety of aerobic, anaerobic, Gram-positive, Gram-negative, and acid-resistant bacteria.

The activity of 14(R)-hydroxy-clarithromycin is greater than that of clarithromycin against *Haemophilus influenzae*. Studies done in vitro have suggested an additive activity of the 14(R)-hydroxy-clarithromycin and the parent molecule against *H. influenzae*.

Category 1: susceptible organisms		
Gram-positive	Gram-negative	Others
<i>Listeria monocytogenes</i>	<i>Bordetella pertussis</i>	<i>Borrelia burgdorferi</i>

Clostridium perfringens Peptococcus niger Propionibacterium acnes Streptococcus group F	Haemophilus influenzae§ Legionella pneumophila Moraxella catarrhalis Pasteurella multocida	Chlamydia pneumoniae (TWAR) Chlamydia trachomatis Mycobacterium avium Mycobacterium chelonae Mycobacterium fortuitum Mycobacterium intracellulare Mycobacterium kansasii Mycobacterium leprae Mycoplasma pneumoniae
Category 2: organisms for which acquired resistance may be problematic#		
Staphylococcus aureus (resistant or susceptible* to methicillin) + Staphylococcus coagulase negative + Streptococcus pneumoniae *+ Streptococcus pyogenes * Streptococcus group B, C, G Streptococcus spp.		
Category 3: intrinsic resistant organisms		
Enterobacteriaceae Non-lactose fermenting Gram negative rods		
* species against which efficacy has been demonstrated in clinical investigations (if susceptible)		
§ Breakpoints for macrolides and related antibiotics were set to categorise wild type H. influenzae as intermediate		
+ Indicates species for which a high rate of resistance (i.e. greater than 50%) has been observed in one or more area/country/region(s) of the EU		
# ≥ 10% resistance in at least one country of the European Union		

## 5.2 Pharmacokinetic properties

The kinetics of orally administered extended-release clarithromycin have been studied in adult humans and compared with clarithromycin 250 mg and 500 mg immediate release tablets. The extent of absorption was found to be equivalent when equal total daily doses were administered. The absolute bioavailability is approximately 50%. Little or no unpredicted accumulation was found and the metabolic disposition did not change in any species following multiple dosing. Based upon the finding of equivalent absorption the following *in vitro* and *in vivo* data are applicable to the extended -release formulation.

*In vitro*: Results of *in vitro* studies showed that the protein binding of clarithromycin in human plasma averaged about 70% at concentrations of 0.45 - 4.5 microgram/mL. A decrease in binding to 41% at 45.0 microgram/mL suggested that the binding sites might become saturated, but this only occurred at concentrations far in excess of therapeutic drug levels.

*In vivo*: Clarithromycin levels in all tissues, except the central nervous system, were several times higher than the circulating drug levels. The highest concentrations were found in the liver and lung tissue, where the tissue to plasma ratios reached 10 to 20.

The pharmacokinetic behaviour of clarithromycin is non-linear. In fed patients given 500 mg clarithromycin extended-release daily, the peak steady state plasma concentration of clarithromycin and 14 hydroxy clarithromycin were 1.3 and 0.48 microgram/mL, respectively. When the dosage was increased to 1000 mg daily, these steady-state values were 2.4 microgram/mL and 0.67 microgram/mL respectively. Elimination half-lives of the parent drug and metabolite were approximately 5.3 and 7.7 hours respectively. The apparent half-lives of both clarithromycin and its hydroxylated metabolite tended to be longer at higher doses.

Urinary excretion accounted for approximately 40% of the clarithromycin dose. Faecal elimination accounts for approximately 30%.

## 5.3 Preclinical safety data

In repeated dose studies, clarithromycin toxicity was related to dose and duration of treatment. The primary target organ was the liver in all species, with hepatic lesions seen after 14 days in dogs and monkeys. Systemic exposure levels associated with this toxicity are not known but toxic mg/kg doses were higher than the dose recommended for patient treatment.

#### Fertility, Reproduction and Teratogenicity

Studies performed in rats at oral doses up to 500 mg/kg/day (highest dose associated with overt renal toxicity) demonstrated no evidence for clarithromycin-related adverse effects on male fertility. This dose corresponds to a human equivalent dose (HED) of approximately 5 times the maximum recommended human dose (MRHD) on a mg/m<sup>2</sup> basis for a 60-kg individual. Fertility and reproduction studies in female rats have shown that a daily dosage of 150mg/kg/day (highest dose tested) caused no adverse effects on the oestrus cycle, fertility, parturition and number and viability of offspring. Oral teratogenicity studies in rats (Wistar and Sprague-Dawley), rabbits (New Zealand White) and cynomolgous monkeys failed to demonstrate any teratogenicity from clarithromycin at the highest doses tested up to 1.5, 2.4 and 1.5 times the MRHD on a mg/m<sup>2</sup> basis in the respective species. However, a similar study in Sprague-Dawley rats indicated a low (6%) incidence of cardiovascular abnormalities which appeared to be due to spontaneous expression of genetic changes. Two mouse studies revealed a variable incidence (3-30%) of cleft palate at ~5 times the MRHD on a mg/m<sup>2</sup> basis for a 60-kg individual. Embryonic loss was seen in monkeys but only at dose levels which were clearly toxic to the mothers.

No evidence of mutagenic potential of clarithromycin was seen during a range of in vitro and in vivo tests.

No other toxicological findings considered to be of relevance to the dose level recommended for patient treatment have been reported.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Lactose Monohydrate  
Hypromellose  
Hypromellose Phthalate  
Talc  
Magnesium Stearate

#### Film Coat

Hypromellose  
Lactose Monohydrate  
Macrogol  
Quinoline Yellow Aluminium Lake (E104)  
Talc  
Titanium Dioxide (E171)

### **6.2 Incompatibilities**

None known.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

Blister pack of Clear PVC/PVDC lidded with aluminium foil for blister pack of 7, 10, 14, 20, 28, 30, 56, 60, 84, 90, 100 and 112 film coated tablets.

Not all pack sizes may be marketed

#### **6.6 Special precautions for disposal and other handling**

No special requirements.

#### **7 MARKETING AUTHORISATION HOLDER**

Aspire Pharma (Malta) Limited  
Notabile Gardens No. 2 Level 3  
Trident Park  
Mdina Road  
Central Business District  
Birkirkara  
CBD 2010  
Malta

#### **8 MARKETING AUTHORISATION NUMBER**

PA23142/006/001

#### **9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of First Authorisation: 15<sup>th</sup> April 2011

Date of last renewal: 13<sup>th</sup> January 2016

#### **10 DATE OF REVISION OF THE TEXT**

June 2024