

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Bondulc 40 micrograms/ml eye drops, solution

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml of solution contains 40 micrograms of Travoprost.

Average active substance/drop: 0.97 - 1.4 µg

Excipients with known effect: benzalkonium chloride 0.150 mg/ml, macrogol glycerol hydroxy stearate 40 5 mg/ml (see section 4.4.).

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Eye drops, solution.

Clear, colourless solution.

pH: 5.5-7.0

Osmolality: 266-294 mOsm/Kg

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Decrease of elevated intraocular pressure in adult patients with ocular hypertension or open-angle glaucoma (see section 5.1).

Decrease of elevated intraocular pressure in paediatric patients aged 2 months to < 18 years with ocular hypertension or paediatric glaucoma (see section 5.1).

4.2 Posology and method of administration

Posology

Use in adults, including elderly population

The dose is one drop of Bondulc in the conjunctival sac of the affected eye(s) once daily. Optimal effect is obtained if the dose is administered in the evening.

Nasolacrimal occlusion or gently closing the eyelid after administration is recommended. This may reduce the systemic absorption of medicinal products administered via the ocular route and result in a decrease in systemic adverse reactions.

If more than one topical ophthalmic medicinal product is being used, the medicinal products must be administered at least 5 minutes apart (see section 4.5).

If a dose is missed, treatment should be continued with the next dose as planned. The dose should not exceed one drop in the affected eye(s) daily.

When substituting another ophthalmic antiglaucoma medicinal product with Bondulc, the other medicinal product should be discontinued and Travoprost should be started the following day.

Hepatic and renal impairment

Travoprost has been studied in patients with mild to severe hepatic impairment and in patients with mild to severe renal impairment (creatinine clearance as low as 14 ml/min). No dosage adjustment is necessary in these patients (see section 5.2).

Paediatric population

Bondulc can be used in paediatric patients from 2 months to < 18 years at the same posology as in adults. However, data in the age group 2 months to < 3 years (9 patients) is limited (see section 5.1).

The safety and efficacy of Bondulc in children below the age of 2 months have not been established. No data are available.

Method of administration

For ocular use

For patients who wear contact lenses, please refer to section 4.4.

The patient should remove the protective overwrap immediately prior to initial use. To prevent contamination of the dropper tip and solution, care must be taken not to touch the eyelids, surrounding areas or other surfaces with the dropper tip of the bottle.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Eye colour change

Travoprost may gradually change the eye colour by increasing the number of melanosomes (pigment granules) in melanocytes. Before treatment is instituted, patients must be informed of the possibility of a permanent change in eye colour. Unilateral treatment can result in permanent heterochromia. The long term effects on the melanocytes and any consequences thereof are currently unknown. The change in iris colour occurs slowly and may not be noticeable for months to years. The change in eye colour has predominantly been seen in patients with mixed coloured irides, i.e., blue-brown, grey-brown, yellow-brown and green-brown; however, it has also been observed in patients with brown eyes. Typically, the brown pigmentation around the pupil spreads concentrically towards the periphery in affected eyes, but the entire iris or parts of it may become more brownish. After discontinuation of therapy, no further increase in brown iris pigment has been observed.

Periorbital and eye lid changes

In controlled clinical trials, periorbital and/or eyelid skin darkening in association with the use of Travoprost has been reported in 0.4% of patients. Periorbital and lid changes including deepening of the eyelid sulcus have also been observed with prostaglandin analogues.

Travoprost may gradually change eyelashes in the treated eye(s); these changes were observed in about half of the patients in clinical trials and include: increased length, thickness, pigmentation, and/or number of lashes. The mechanism of eyelash changes and their long term consequences are currently unknown.

Travoprost has been shown to cause slight enlargement of the palpebral fissure in studies in the monkey. However, this effect was not observed during the clinical trials and is considered to be species specific.

There is no experience of Travoprost in inflammatory ocular conditions; nor in neovascular, angle-closure, narrow-angle or congenital glaucoma and only limited experience in thyroid eye disease, in open-angle glaucoma of pseudophakic patients and in pigmentary or pseudoexfoliative glaucoma.

Travoprost should therefore be used with caution in patients with active intraocular inflammation.

Aphakic patients

Macular oedema has been reported during treatment with prostaglandin F2a analogues. Caution is recommended when using Travoprost in aphakic patients, pseudophakic patients with a torn posterior lens capsule or anterior chamber lenses, or in patients with known risk factors for cystoid macular oedema.

Iritis/uveitis

In patients with known predisposing risk factors for iritis/uveitis, Travoprost should be used with caution.

Contact with the skin

Skin contact with Bondulc must be avoided as transdermal absorption of travoprost has been demonstrated in rabbits.

Prostaglandins and prostaglandin analogues are biologically active materials that may be absorbed through the skin. Women who are pregnant or attempting to become pregnant should exercise appropriate precautions to avoid direct exposure to the contents of the bottle. In the unlikely event of coming in contact with a substantial portion of the contents of the bottle, thoroughly cleanse the exposed area immediately.

Excipients

Bondulc contains the preservative benzalkonium chloride, which may cause eye irritation. Avoid contact with soft contact lenses.

Remove contact lenses prior to application and wait at least 15 minutes before reinsertion.

Known to discolour soft contact lenses.

Bondulc contains Macrogol glycerol hydroxy stearate 40 which may cause skin reactions.

Paediatric population

Efficacy and safety data in the age group 2 months to < 3 years (9 patients) is limited (see section 5.1). No data are available for children below the age of 2 months.

In children < 3 years old that mainly suffer from PCG (primary congenital glaucoma), surgery (e.g. trabeculotomy/goniotomy) remains the first line treatment.

No long-term safety data are available in the paediatric population.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed.

4.6 Fertility, pregnancy and lactation

Women of child-bearing potential/contraception

Travoprost must not be used in women of child bearing age/potential unless adequate contraceptive measures are in place (see section 5.3).

Pregnancy

Travoprost has harmful pharmacological effects on pregnancy and/or the foetus/new-born child. Travoprost should not be used during pregnancy unless clearly necessary.

Breastfeeding

It is unknown whether travoprost from the eye drops is excreted in human breast milk. Animal studies have shown excretion of travoprost and metabolites in breast milk. The use of Travoprost by breast-feeding mothers is not recommended.

Fertility

There are no data on the effects of travoprost on human fertility. Animal studies showed no effect of travoprost on fertility at doses more than 250 times the maximum recommended human ocular dose.

4.7 Effects on ability to drive and use machines

Bondulc has no or negligible influence on the ability to drive and use machines, however as with any eye drop, temporary blurred vision or other visual disturbances may affect the ability to drive or use machines. If blurred vision occurs at instillation, the patient must wait until the vision clears before driving or using machinery.

4.8 Undesirable effects

Summary of the safety profile

In clinical trials with travoprost, the most common adverse reactions were ocular hyperaemia and iris hyperpigmentation, occurring in approximately 20% and 6% of patients respectively.

Tabulated list of adverse reactions

The following adverse reactions are classified according to the following convention: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare $< 1/10,000$), or not known (frequency cannot be estimated from the available data). Within each frequency group, adverse reactions are presented in decreasing order of seriousness. The adverse reactions were obtained from clinical studies and post- marketing data with travoprost.

System Organ Class	Frequency	Adverse Reactions
Infections and infestations	Rare	herpes simplex, keratitis herpetic
Immune system disorders	Uncommon	hypersensitivity, seasonal allergy
Psychiatric disorders	Not known	depression, anxiety
Nervous system disorders	Uncommon	Headache, dizziness, visual field defect
	Rare	dysgeusia,
Eye disorders	Very common	ocular hyperaemia
	Common	iris hyperpigmentation, eye pain, ocular

		discomfort, dry eye, eye pruritus, eye irritation
	Uncommon	corneal erosion, uveitis, iritis, anterior chamber inflammation, keratitis, punctate keratitis, photophobia, eye discharge, blepharitis, erythema of eyelid, periorbital oedema, eyelids pruritus, visual acuity reduced, vision blurred, lacrimation increased, conjunctivitis, ectropion, cataract, eyelid margin crusting, growth of eyelashes, eyelash discolouration, asthenopia
	Rare	iritidocyclitis, eye inflammation, photopsia, eczema eyelids, conjunctival oedema, halo vision, conjunctival follicles, hypoaesthesia eye, meibomianitis, anterior chamber pigmentation, myadriasis, eyelash thickening
	Not known	macular oedema, sunken eyes
Ear and labyrinth disorders	Not known	vertigo, tinnitus
	Uncommon	palpitations, d
Cardiac disorders	Rare	heart rate irregular, heart rate decreased
	Not known	chest pain, bradycardia, tachycardia
Vascular disorders	Rare	blood pressure diastolic decreased, blood pressure systolic increased, hypotension, hypertension
Respiratory, thoracic and mediastinal disorders	Uncommon	dyspnoea, asthma, nasal congestion, throat irritation
	Rare	respiratory disorder, oropharyngeal pain, cough, dysphonia
	Not known	asthma aggravated
Gastrointestinal disorders	Rare	peptic ulcer reactivated, gastrointestinal disorder, constipation, dry mouth
	Not known	diarrhoea, abdominal pain, nausea

Skin and subcutaneous tissue disorders	Uncommon	skin hyperpigmentation (periocular) skin discolouration, hair texture abnormal, hypertrichosis
	Rare	dermatitis allergic, dermatitis contact, erythema, rash, hair colour changes, madarosis
	Not known	pruritus, hair growth abnormal
Musculoskeletal, connective tissue and bone disorders	Rare	musculoskeletal pain
	Not known	arthralgia
Renal and urinary disorders	Not known	dysuria, urinary incontinence
General disorders and administrative site conditions	Uncommon	asthenia
Investigations	Not known	prostatic specific antigen increased

Paediatric Population

In a 3 month phase 3 study and a 7 days pharmacokinetic study, involving 102 paediatric patients exposed to Travoprost, the types and characteristics of adverse reactions reported were similar to what has been observed in adult patients. The short-term safety profiles in the different paediatric subsets were also similar (see section 5.1). The most frequent adverse reactions reported in the paediatric population were ocular hyperaemia (16.9%) and growth of eyelashes (6.5%). In a similar 3 month study in adult patients, these events occurred at an incidence of 11.4% and 0.0%, respectively.

Additional adverse drug reactions reported in paediatric patients in the 3 month paediatric study (n=77) compared to a similar trial in adults (n=185) included erythema of eyelid, keratitis, lacrimation increased, and photophobia all reported as single events with an incidence of 1.3% versus 0.0% seen in adults.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL- Dublin 2; Tel: +353 1 6764971; Fax:+353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

4.9 Overdose

No cases of overdose have been reported. A topical overdose is not likely to occur or to be associated with toxicity. A topical overdose of Travoprost may be flushed from the eye(s) with lukewarm water. Treatment of a suspected oral ingestion is symptomatic and supportive.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic Group: Ophthalmologicals -antiglaucoma preparations and miotics- prostaglandin analogues

ATC code: S0 1E E04

Mechanism of action

Travoprost, a prostaglandin F2α analogue, is a highly selective full agonist which has a high affinity for the prostaglandin FP receptor, and reduces the intraocular pressure by increasing the outflow of aqueous humour via trabecular meshwork and uveoscleral pathways. Reduction of the intraocular pressure in man starts about 2 hours after administration and maximum effect is reached after 12 hours. Significant lowering of intraocular pressure can be maintained for periods exceeding 24 hours with a single dose.

Clinical efficacy and safety

Data on adjunctive administration of travoprost with timolol 0.5% and limited data with brimonidine 0.2% were collected during clinical trials that showed an additive effect of travoprost with these glaucoma medications. No clinical data are available on adjunctive use with other ocular hypotensive medications.

Secondary pharmacology

Travoprost significantly increased optic nerve head blood flow in rabbits following 7 days of topical ocular administration (1.4 micrograms, once-daily).

Paediatric population

The efficacy of travoprost in paediatric patients from 2 months to less than 18 years of age was demonstrated in a 12-week, double-masked clinical study of travoprost compared with timolol in 152 patients diagnosed with ocular hypertension or paediatric glaucoma. Patients received either travoprost 0.004% once daily or timolol 0.5% (or 0.25% for subjects younger than 3 years old) twice daily. The primary efficacy endpoint was the intraocular pressure (IOP) change from baseline at Week 12 of the study. Mean IOP reductions in the travoprost and timolol groups were similar (see Table 1).

In the age groups 3 to < 12 years (n=36) and 12 to <18 years (n=26), mean IOP reduction at Week 12 in the travoprost group was similar to that in the timolol group. Mean IOP reduction at Week 12 in the 2 months to < 3 years of age group was 1.8 mmHg in the travoprost group and 7.3 mmHg in the timolol group. IOP reductions for this group were based on only 6 patients in the timolol group and 9 patients in the travoprost group where 4 patients in the travoprost group versus 0 patients in the timolol group had no relevant mean IOP reduction at Week 12. No data are available for children less than 2 months old.

The effect on IOP was seen after the second week of treatment and was consistently maintained throughout the 12 week period of study for all age groups.

Table 1 - Comparison of mean IOP change from baseline (mmHg) at week 12

Travoprost			Timolol		
N	Mean (SE)	N	Mean (SE)	Mean difference ^a	(95 % CI)
53	-6.4	60	-5.8	-0.5	(-2.1, 1.0)
	(1.05)		(0.96)		

SE = Standard Error; CI = Confidence Interval;

^aMean differences is Travoprost–Timolol. Estimates based on least squares means derived from a statistical model that accounts for correlated IOP measurements within patient where primary diagnosis and baseline IOP stratum are in the model.

5.2 Pharmacokinetic properties

Absorption

Travoprost is an ester prodrug. It is absorbed through the cornea where the isopropyl ester is hydrolysed to the active free acid. Studies in rabbits have shown peak concentrations of 20 ng/g of the free acid in aqueous humour one to two hours after topical dosing of Travoprost. Aqueous humour concentrations declined with a half-life of approximately 1.5 hours.

Distribution

Following topical ocular administration of Travoprost to healthy volunteers, low systemic exposure to active free acid was demonstrated. Peak active free acid plasma concentrations of 25pg/ml or less were observed between 10 and 30 minutes post-dose. Thereafter, plasma levels declined rapidly to below the 10 pg/ml assay quantitation limit before 1 hour post-administration. Due to the low plasma concentrations and rapid elimination following topical dosing, the elimination half-life of active free acid in man could not be determined.

Biotransformation

Metabolism is the major route of elimination of both Travoprost and the active free acid. The systemic metabolic pathways parallel those of endogenous prostaglandin $F_{2\alpha}$ which are characterised by reduction of the 13-14 double bond, oxidation of the 15-hydroxyl and β -oxidative cleavages of the upper side chain.

Elimination

Travoprost free acid and its metabolites are mainly excreted by the kidneys. Travoprost has been studied in patients with mild to severe hepatic impairment and in patients with mild to severe renal impairment (creatinine clearance as low as 14 ml/min). No dosage adjustment is necessary in these patients.

Paediatric population

A pharmacokinetic study in paediatric patients aged 2 months to <18 years demonstrated very low plasma exposure to travoprost free acid, with concentrations ranging from below the 10 pg/mL assay limit of quantitation (BLQ) to 54.5 pg/mL. In 4 previous systemic pharmacokinetic studies in adult populations, travoprost free acid plasma concentrations ranged from BLQ to 52.0 pg/mL. While most of the plasma data across all studies was non-quantifiable, making statistical comparisons of systemic exposure across age groups unfeasible, the overall trend shows that plasma exposure to travoprost free acid following topical administration of travoprost is extremely low across all age groups evaluated.

5.3 Preclinical safety data

In ocular toxicity studies in monkeys, administration of Travoprost at a dose of 0.45 microgram, twice a day, was shown to induce increased palpebral fissure. Topical ocular administration of Travoprost to monkeys at concentrations of up to 0.012% to the right eye, twice daily for one year resulted in no systemic toxicity.

Reproduction toxicity studies have been undertaken in rat, mice and rabbit by systemic route. Findings are related to FP receptor agonist activity in uterus with early embryoletality, post-implantation loss, foetotoxicity. In pregnant rat, systemic administration of Travoprost at doses more than 200 times the clinical dose during the period of organogenesis resulted in an increased incidence of malformations. Low levels of radioactivity were measured in amniotic fluid and foetal tissues of pregnant rats administered 3H Travoprost. Reproduction and development studies have demonstrated a potent effect on foetal loss with a high rate observed in rats and mice at plasma levels of 180 pg/ml and 30 pg/ml, respectively corresponding to 1.2 to 6 times the clinical exposure (up to 25 pg/ml).

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Benzalkonium chloride solution

Macrogol glycerol hydroxy stearate 40

Trometamol
 Disodium edetate
 Boric acid (E284)
 Mannitol (E421)
 Sodium hydroxide (for pH adjustment)
 Water for injection

6.2 Incompatibilities

Not applicable.

Specific *in vitro* interaction studies were performed with Travoprost and medicinal products containing thiomersal. No evidence of precipitation was observed.

6.3 Shelf life

Unopened product: 3 years

Discard 4 weeks after first opening.

6.4 Special precautions for storage

Before opening, keep bottle in overwrap pouch in order to protect from moisture.

After first opening, this medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

Translucent polypropylene (PP) 5 ml bottle with a transparent low density polyethylene (LDPE) dropper and a white high density polyethylene (HDPE) tamper-proof screw cap, presented in a polyethylene terephthalate/aluminium/polyethylene (PET/Alu/PE) overwrap. Each bottle contains 2.5 ml eye drops.

Pack sizes:

Cartons containing 1 or 3 bottles.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Actavis Group PTC ehf
 Reykjavíkurvegi 76-78
 220 Hafnarfjörður
 Iceland

8 MARKETING AUTHORISATION NUMBER

PA1380/169/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 14th November 2014

10 DATE OF REVISION OF THE TEXT

August 2015