

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Tacrolimus 1.0mg hard capsules

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each capsule contains 1 mg of tacrolimus (as tacrolimus monohydrate)

Excipients: 48.68 mg of lactose monohydrate

For the full list of excipients see section 6.1

3 PHARMACEUTICAL FORM

Capsule hard

White / White hard gelatin capsules, approximately 11.40mm, size "5" imprinted with "TCR" on cap & "1" on body containing white to off white granular powder

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Prophylaxis of transplant rejection in liver, kidney or heart allograft recipients.

Treatment of allograft rejection resistant to treatment with other immunosuppressive medicinal products.

4.2 Posology and method of administration

Tacrolimus therapy requires careful monitoring by adequately qualified and equipped personnel. This medicinal product should only be prescribed and changes in immunosuppressive therapy, initiated by physicians experienced in immunosuppressive therapy and the management of transplanted patients.

Inadvertent, unintentional or unsupervised switching of immediate- or prolonged-release formulations of tacrolimus is unsafe. This can lead to graft rejection or increased incidence of side effects, including under- or over immunosuppression, due to clinically relevant differences in systemic exposure to tacrolimus. Patients should be maintained on a single formulation of tacrolimus with the corresponding daily dosing regimen; alterations in formulation or regimen should only take place under the close supervision of a transplant specialist (see sections 4.4 and 4.8). Following conversion to any alternative formulation, therapeutic drug monitoring must be performed and dose adjustments made to ensure that systemic exposure to tacrolimus is maintained.

General considerations

The recommended initial dosages presented below are intended to act solely as a guideline. Tacrolimus capsules dosing should primarily be based on clinical assessments of rejection and tolerability in each patient individually aided by blood level monitoring, (see below for recommended target whole blood trough concentrations). If clinical signs of rejection are apparent, alteration of the immunosuppressive regimen should be considered.

Tacrolimus can be administered intravenously or orally. In general, dosing may commence orally; if necessary, by administering the capsule contents suspended in water, via nasogastric tubing.

Tacrolimus capsules is routinely administered in conjunction with other immunosuppressive agents in the initial post-operative period. The Tacrolimus capsules dose may vary depending upon the immunosuppressive regimen chosen.

Method of administration

It is recommended that the oral daily dose be administered in two divided doses (e.g. morning and evening). Capsules

should be taken immediately following removal from the blister. The capsules should be swallowed with fluid (preferably water).

Capsules should generally be administered on an empty stomach or at least 1 hour before or 2 to 3 hours after a meal to achieve maximal absorption (see section 5.2).

Duration of treatment

To suppress graft rejection, immunosuppression must be maintained; consequently, no limit to the duration of oral therapy can be given.

Dosage recommendations – Liver transplantation

Prophylaxis of transplant rejection - adults

Oral Tacrolimus therapy should commence at 0.10 - 0.20 mg/kg/day administered as two divided doses (e.g. morning and evening). Administration should commence approximately 12 hours after the completion of surgery.

If the dose cannot be administered orally as a result of the clinical condition of the patient intravenous administration of 0.01 - 0.05 mg/kg/day should be initiated as a 24-hour continuous infusion.

Prophylaxis of transplant rejection - children

An initial oral dose of 0.30 mg/kg/day should be administered in two divided doses (e.g. morning and evening). If the clinical condition of the patient prevents oral dosing, an initial intravenous dose of 0.05 mg/kg/day should be administered as a 24-hour continuous infusion.

Dose adjustments during post-transplant period in adults and children

Tacrolimus capsules doses are usually reduced in the post-transplant period. In some cases it is possible to withdraw concomitant immunosuppressive therapy leading to Tacrolimus capsules monotherapy. Post-transplant improvement in the condition of the patient may alter the pharmacokinetics of tacrolimus and may necessitate further dose adjustments.

Rejection therapy – adults and children

Increased Tacrolimus capsules doses, supplemental corticosteroid therapy and introduction of short courses of monoclonal/polyclonal antibodies have all been used to manage rejection episodes. If signs of toxicity are noted (e.g. pronounced adverse reactions - see section 4.8) the dose of Tacrolimus capsules may need to be reduced.

For conversion to Tacrolimus capsules, treatment should begin with the initial oral dose recommended for primary immunosuppression.

For information on conversion from ciclosporin to tacrolimus, see below under “Dose adjustments in specific patient populations”.

Dosage recommendations - Kidney transplantation

Prophylaxis of transplant rejection – adults

Oral tacrolimus therapy should commence at 0.20 - 0.30 mg/kg/day administered as two divided doses (e.g. morning and evening). Administration should commence within 24 hours after the completion of surgery.

If the dose cannot be administered orally as a result of the clinical condition of the patient intravenous therapy of 0.05 - 0.10 mg/kg/day should be initiated as a 24-hour continuous infusion.

Prophylaxis of transplant rejection – children

An initial oral dose of 0.30 mg/kg/day should be administered in two divided doses (e.g. morning and evening). If the clinical condition of the patient prevents oral dosing, an initial intravenous dose of 0.075 – 0.100 mg/kg/day should be administered as a 24-hour continuous infusion.

Dose adjustments during post-transplant period in adults and children

Tacrolimus capsules doses are usually reduced in the post-transplant period. In some cases it is possible to withdraw

concomitant immunosuppressive therapy leading to Tacrolimus capsules -based dual therapy. Post-transplant improvement in the condition of the patient may alter the pharmacokinetics of tacrolimus and may necessitate further dose adjustments.

Rejection therapy – adults and children

Increased Tacrolimus capsules doses, supplemental corticosteroid therapy, and introduction of short courses of monoclonal/polyclonal antibodies have all been used to manage rejection episodes. If signs of toxicity are noted (e.g. pronounced adverse reactions - see section 4.8) the dose of Tacrolimus capsules may need to be reduced.

For conversion to Tacrolimus capsules, treatment should begin with the initial oral dose recommended for primary immunosuppression.

For information on conversion from ciclosporin to tacrolimus, see below under “Dose adjustments in specific patient populations”

Dosage recommendations - Heart transplantation

Prophylaxis of transplant rejection – adults

Tacrolimus capsules can be used with antibody induction (allowing for delayed start of Tacrolimus capsules therapy) or alternatively in clinically stable patients, without antibody induction.

Following antibody induction, oral Tacrolimus capsules therapy should commence at a dose of 0.075 mg/kg/day administered as two divided doses (e.g. morning and evening). Administration should commence within 5 days after the completion of surgery, as soon as the patient's clinical condition is stabilised. If the dose cannot be administered orally as a result of the clinical condition of the patient, intravenous therapy with 0.01 to 0.02 mg/kg/day should be initiated as a 24-hour continuous infusion.

An alternative strategy was published where oral tacrolimus was administered within 12 hours post transplantation. This approach was reserved for patients without organ dysfunction (e.g. renal dysfunction). In that case, an initial oral tacrolimus dose of 2 to 4 mg per day was used in combination with mycophenolate mofetil and corticosteroids or in combination with sirolimus and corticosteroids.

Prophylaxis of transplant rejection – children

Tacrolimus capsules have been used in combination with (or without) antibody induction in paediatric heart transplantation.

In patients without antibody induction, if Tacrolimus therapy is initiated intravenously, the recommended starting dose is 0.03 - 0.05 mg/kg/day as a 24-hour continuous infusion targeted to achieve tacrolimus whole blood concentration of 15 - 25 ng/ml. Patients should be converted to oral therapy as soon as clinically practicable. The first dose of oral therapy should be 0.30 mg/kg/day starting 8 to 12 hours after discontinuing intravenous therapy.

Following antibody induction, if Tacrolimus capsules therapy is initiated orally, the recommended starting dose is 0.10 - 0.30 mg/kg/day administered as two divided doses (e.g. morning and evening).

Dose adjustments during post-transplant period in adults and children

Tacrolimus capsules doses are usually reduced in the post-transplant period. Post-transplant improvement in the condition of the patient may alter the pharmacokinetics, of tacrolimus and may necessitate further dose adjustments.

Rejection therapy – adults and children

Increased Tacrolimus capsules doses, supplemental corticosteroid therapy and introduction of short courses of monoclonal/polyclonal antibodies have all been used to manage rejection episodes.

In adults patients converted to Tacrolimus capsules, an initial oral dose of 0.15 mg/kg/day should be administered in two divided doses (e.g. mornings and evenings).

In paediatric patients converted to Tacrolimus, an initial oral dose of 0.20 – 0.30 mg/kg/day should be administered in two divided doses (e.g. mornings and evenings).

For information on conversion from ciclosporin to Tacrolimus, see below under “Dose adjustments in specific patient populations”.

Dosage recommendations - Rejection therapy, other allografts

The dose recommendations for lung, pancreas and intestinal transplantation are based on limited prospective clinical trial data. In lung transplanted patients, Tacrolimus capsules have been used at an initial oral dose of 0.10 - 0.15 mg/kg/day, in pancreas -transplanted patients at an initial oral dose of 0.2 mg/kg/day and in intestinal transplantation at an initial oral dose of 0.3 mg/kg/day

Dosage adjustments in specific patient populations

Race

In comparison to Caucasians, black patients may require higher tacrolimus doses to achieve similar trough levels.

Gender

There is no evidence that male and female patients require different doses to achieve similar trough levels.

Patients with liver impairment

Dose reduction may be necessary in patients with severe liver impairment in order to maintain the blood trough levels within the recommended target range.

Patients with kidney impairment

As the pharmacokinetics of tacrolimus are unaffected by renal function, no dose adjustments should be required. However, owing to the nephrotoxic potential of tacrolimus, careful monitoring of renal function is recommended (including serial serum creatinine concentrations, calculation of creatinine clearance and monitoring of urine output).

Paediatric patients

In general, paediatric patients require doses 1½ - 2 times higher than the adult doses to achieve similar blood levels.

Elderly patients

There is currently no evidence available to indicate that dosing should be adjusted in elderly patients.

Conversion from ciclosporin

Care should be taken when converting patients from ciclosporin-based to tacrolimus based therapy (see sections 4.4 and 4.5). Tacrolimus capsules therapy should be initiated after considering ciclosporin blood concentrations and the clinical condition of the patient. Dosing should be delayed in the presence of elevated ciclosporin blood levels. In practice, Tacrolimus capsules therapy has been initiated 12 - 24 hours after discontinuation of ciclosporin. Monitoring of ciclosporin blood levels should be continued following conversion as the clearance of ciclosporin might be affected.

Target whole blood trough concentrations recommendations

Dosing should primarily be based on clinical assessments of rejection and tolerability in each individual patient.

As an aid to optimise dosing, several immunoassays are available for determining tacrolimus concentrations in whole blood, including a semi-automated microparticle enzyme immunoassay (MEIA). Comparison of concentrations from the published literature to individual values in clinical practice should be assessed with care and knowledge of the assay methods employed. In current clinical practice, whole blood levels are monitored using immunoassay methods.

Blood trough levels of tacrolimus should be monitored during the post-transplantation period. When dosed orally, blood trough levels should be drawn approximately 12 hours post dosing, just prior to the next dose. The frequency of blood level monitoring should be based on clinical needs. As Tacrolimus capsules is a medicinal product with low clearance, adjustments to the dosage regimen may take several days before changes in blood levels are apparent; Blood

trough levels of tacrolimus should be monitored approximately twice weekly during the early post-transplant period and then periodically during maintenance therapy. Blood trough levels of tacrolimus should also be monitored following dose adjustment, changes in the immunosuppressive regimen or following co-administration of substances which may alter tacrolimus whole blood concentrations (see section 4.5).

Clinical studies analysis suggests that the majority of patients can be successfully managed if tacrolimus blood trough levels are maintained below 20 ng/ml. It is necessary to consider the clinical condition of the patient when interpreting whole blood levels.

In clinical practice, whole blood trough levels have generally been in the range 5 - 20 ng/ml in liver transplant recipients and 10 - 20 ng/ml in kidney and heart transplant patients in the early post-transplant period. Subsequently, during maintenance therapy, blood concentrations have generally been in the range of 5 - 15 ng/ml in liver, heart and kidney transplant recipients.

4.3 Contraindications

Hypersensitivity to the active substance, other macrolides or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Monitoring during post-transplantation period

During the initial post-transplantation period, monitoring of the following parameters should be undertaken on a routine basis: blood pressure, ECG, neurological and visual status, fasting blood glucose levels, electrolytes (particularly potassium), liver and renal function tests, haematology parameters, coagulation values and plasma protein determinations. If clinically relevant changes are seen, adjustments of the immunosuppressive regimen should be considered.

Medication errors

Medication errors, including inadvertent, unintentional or unsupervised substitution of immediate- or prolonged-release tacrolimus formulations, have been observed. This has led to serious adverse events, including graft rejection, or other side effects which could be a consequence of either under- or over-exposure to tacrolimus. Patients should be maintained on a single formulation of tacrolimus with the corresponding daily dosing regimen; alterations in formulations or regimen should only take place under the close supervision of a transplant specialist (see sections 4.2 and 4.8).

Herbal preparations

Herbal preparations containing St. John's Wort (*Hypericum perforatum*) or other herbal preparations should be avoided when taking Tacrolimus capsules, due to the risk of interactions that lead to decrease in blood concentrations of tacrolimus and reduced clinical effect of tacrolimus (see section 4.5).

Diarrhoea

Since levels of tacrolimus in blood may significantly change during diarrhoea episodes, extra monitoring of tacrolimus concentrations is recommended during episodes of diarrhoea.

Ciclosporin

The combined administration of ciclosporin and tacrolimus should be avoided and care should be taken when administering tacrolimus to patients who have previously received ciclosporin (see sections 4.2 and 4.5).

Cardiac disorders

Ventricular hypertrophy or hypertrophy of the septum, reported as cardiomyopathies, have been observed on rare occasions. Most cases have been reversible, occurring primarily in children with tacrolimus blood trough concentrations much higher than the recommended maximum levels. Other factors observed to increase the risk of these clinical conditions included pre-existing heart disease, corticosteroid usage, hypertension, renal or hepatic dysfunction, infections, fluid overload and oedema. Accordingly, high-risk patients, particularly young children and

those receiving substantial immunosuppression, should be monitored using such procedures as echocardiography, or ECG pre- and post-transplant (e.g. initially at three months and then at 9-12 months). If abnormalities develop, dose reduction of Tacrolimus capsules therapy or change of treatment to another immunosuppressive agent should be considered. Tacrolimus may prolong the QT interval but at this time lacks substantial evidence for causing Torsades de Pointes. Caution should be exercised in patients with diagnosed or suspected Congenital Long QT Syndrome.

Epstein-Barr virus (EBV) associated lymphoproliferative disorders

Patients treated with Tacrolimus capsules have been reported to develop Epstein-Barr virus (EBV) associated lymphoproliferative disorders. Patients switched to Tacrolimus capsules therapy should not receive anti-lymphocyte treatment Concomitantly. Very young (< 2 years), EBV-VCA-negative children have been reported to have an increased risk of developing a lymphoproliferative disorders. Therefore, in this patient group, the EBV-VCA serology should be ascertained before starting treatment with tacrolimus. During treatment, careful monitoring with EBV-PCR is recommended. Positive EBV-PCR may persist for months and is per se not indicative of lymphoproliferative disease or lymphoma.

Posterior reversible encephalopathy syndrome (PRES)

Patients treated with tacrolimus have been reported to develop posterior reversible encephalopathy syndrome (PRES). If patients taking tacrolimus present with symptoms indicating PRES such as headache, altered mental status, seizures, and visual disturbances, a radiological procedure (e.g. MRI) should be performed. If PRES is diagnosed, adequate blood pressure and seizure control and immediate discontinuation of systemic tacrolimus is advised. Most patients completely recover after appropriate measures are taken.

Pure Red Cell Aplasia

Cases of pure red cell aplasia (PRCA) have been reported in patients treated with tacrolimus. All patients reported risk factors for PRCA such as parvovirus B19 infection, underlying disease or concomitant medications associated with PRCA.”

Risk of opportunistic infections

Patients treated with immunosuppressants, including tacrolimus are at increased risk for opportunistic infections (bacterial, fungal, viral and protozoal). Among these conditions are BK virus associated nephropathy and JC virus associated progressive multifocal leukoencephalopathy (PML). These infections are often related to a high total immunosuppressive burden and may lead to serious or fatal conditions that physicians should consider in the differential diagnosis in immunosuppressed patients with deteriorating renal function or neurological symptoms.

Photosensitivity

As with other immunosuppressive agents, owing to the potential risk of malignant skin changes, exposure to sunlight and UV light should be limited by wearing protective clothing and using a sunscreen with a high protection factor.

Others

As with other potent immunosuppressive compounds, the risk of secondary cancer is unknown (see section 4.8).

Allergic and anaphylactoid reactions have been observed in patients using tacrolimus (see section 4.8)

As Tacrolimus capsules contains lactose, special care should be taken in patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption.

This medicinal product contains less than 1 mmol sodium (23mg) per dose, i.e. ‘essential sodium free’.

4.5 Interaction with other medicinal products and other forms of interaction

Metabolic interactions

Systemically available tacrolimus is metabolised via by hepatic CYP3A4 in the liver. There is also evidence of gastrointestinal metabolism by CYP3A4 in the intestinal wall. Concomitant use of medicinal products or herbal remedies known to inhibit or induce CYP3A4 may affect the metabolism of tacrolimus and thereby increase or decrease tacrolimus blood levels. It is therefore recommended to monitor tacrolimus blood levels whenever substances which have the potential to alter CYP3A metabolism are used concomitantly, and to adjust the dose of tacrolimus as appropriate in order to maintain similar tacrolimus exposure (see sections 4.2 and 4.4).

Inhibitors of metabolism

Clinically the following substances have been shown to increase tacrolimus blood levels:

Strong interactions have been observed with antifungal agents such as ketoconazole, fluconazole, itraconazole and voriconazole, the macrolide antibiotic erythromycin or HIV protease inhibitors (e.g. ritonavir). Concomitant use of these substances may require decreased tacrolimus doses in nearly all patients.

Weaker interactions have been observed with clotrimazole, clarithromycin, josamycin, nifedipine, nicardipine, diltiazem, verapamil, danazol, ethinylestradiol, omeprazole and nefazodone.

In vitro the following substances have been shown to be potential inhibitors of tacrolimus metabolism: bromocriptine, cortisone, dapson, ergotamine, gestodene, lidocaine, mephenytoin, miconazole, midazolam, nilvadipine, norethisterone, quinidine, tamoxifen, troleandomycin.

Grapefruit juice has been reported to increase the blood level of tacrolimus and should therefore be avoided.

Lansoprazole and ciclosporin may potentially inhibit CYP3A4-mediated metabolism of tacrolimus and thereby increase tacrolimus whole blood concentrations.

Inducers of metabolism

Clinically the following substances have been shown to decrease tacrolimus blood levels:

Strong interactions have been observed with rifampicin, phenytoin or St. John's Wort (*Hypericum perforatum*) which may require increased tacrolimus doses in almost all patients. Clinically significant interactions have also been observed with phenobarbital. Maintenance doses of corticosteroids have been shown to reduce tacrolimus blood levels.

High doses prednisolone or methylprednisolone, administered for the treatment of acute rejection have the potential to increase and decrease tacrolimus blood levels.

Carbamazepine, metamizole and isoniazid have the potential to decrease tacrolimus concentrations.

Effect of tacrolimus on the metabolism of other medicinal products

Tacrolimus is a known CYP3A4 inhibitor, thus concomitant use of tacrolimus with medicinal products known to be metabolised by CYP3A4 may affect the metabolism of such medicinal products.

The half-life of ciclosporin is prolonged when tacrolimus is given concomitantly. In addition, synergistic/additive nephrotoxic effects can occur. For these reasons, the combined administration of ciclosporin and tacrolimus is not recommended and care should be taken when administering tacrolimus to patients who have previously received ciclosporin (see sections 4.2 and 4.4).

Tacrolimus has been shown to increase the blood level of phenytoin.

As tacrolimus may reduce the clearance of steroid-based contraceptives leading to increased hormone exposure, particular care should be exercised when deciding upon contraceptive measures.

Limited knowledge of interactions between tacrolimus and statins is available. Available data suggests that the pharmacokinetics of statins are largely unaltered by the co-administration of tacrolimus.

Animal data have shown that tacrolimus could potentially decrease the clearance and increase the half-life of pentobarbital and phenazone.

Other interactions which have led to clinically detrimental effects

Concurrent use of tacrolimus with medicinal products known to have nephrotoxic or neurotoxic effects may increase these effects (e.g. aminoglycosides, gyrase inhibitors, vancomycin, sulfamethoxazole+trimethoprim, NSAIDs, ganciclovir or aciclovir).

Enhanced nephrotoxicity has been observed following the administration of amphotericin B and ibuprofen in conjunction with tacrolimus.

As tacrolimus treatment may be associated with hyperkalaemia or may increase pre-existing hyperkalaemia, high potassium intake, or potassium-sparing diuretics (e.g. amiloride, triamterene or spironolactone) should be avoided.

Immunosuppressants may affect the response to vaccination, and vaccination during treatment with tacrolimus may be less effective. The use of live attenuated vaccines should be avoided.

Protein binding considerations

Tacrolimus is extensively bound to plasma proteins. Possible interactions with other medicinal products known to have high affinity for plasma proteins should be considered (e.g. NSAIDs, oral anticoagulants, or oral antidiabetics).

4.6 Fertility, pregnancy and lactation

Human data show that tacrolimus is able to cross the placenta. Limited data from organ transplant recipients show no evidence of an increased risk of adverse effects on the course and outcome of pregnancy under tacrolimus treatment compared with other immunosuppressive medicinal products. However, cases of spontaneous abortion have been reported. To date, no other relevant epidemiological data are available. Due to the need of treatment, tacrolimus can be considered in pregnant women when there is no safer alternative and when the perceived benefit justifies the potential risk to the foetus. In case of in utero exposure, monitoring of the newborn for the potential adverse effects of tacrolimus is recommended (in particular the effects on the kidneys). There is a risk for premature delivery (<37 weeks) as well as for hyperkalaemia in the newborn, which however normalizes spontaneously.

In rats and rabbits, tacrolimus causes embryofoetal toxicity at doses which demonstrated maternal toxicity (see section 5.3). Tacrolimus affected male fertility in rats (see section 5.3).

Lactation

Human data demonstrate that tacrolimus is excreted in breast milk. As detrimental effects on the newborn cannot be excluded, women should not breast-feed whilst receiving Tacrolimus capsules.

Fertility

A negative effect of tacrolimus on male fertility in the form of reduced sperm counts and motility was observed in rats (see section 5.3).

4.7 Effects on ability to drive and use machines

Tacrolimus may cause visual and neurological disturbances. This effect may be enhanced if Tacrolimus capsules is administered in association with alcohol.

4.8 Undesirable effects

The adverse drug reaction profile associated with immunosuppressive agents is often difficult to establish owing to the underlying disease and the concurrent use of multiple medications.

Many of the adverse drug reactions stated below are reversible and/or respond to dose reduction. Oral administration appears to be associated with a lower incidence of adverse drug reactions compared with intravenous use. Adverse drug reactions are listed below in descending order by frequency of occurrence: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$) and very rare ($< 1/10,000$); not known (cannot be estimated from the available data).

The following adverse reactions have been reported in post-marketing experience:

<u>Infections and infestations</u>
As is well known for other potent immunosuppressive agents, patients receiving

tacrolimus are frequently at increased risk for infections (viral, bacterial, fungal, protozoal). The course of pre-existing infections may be aggravated. Both generalised and localised infections can occur. Cases of BK virus associated nephropathy, as well as cases of JC virus associated progressive multifocal leukoencephalopathy (PML), have been reported in patients treated with immunosuppressants, including tacrolimus.	
<u>Neoplasms benign, malignant and unspecified (including cysts and polyps)</u>	
Patients receiving immunosuppressive therapy are at increased risk of developing malignancies. Benign as well as malignant neoplasms, including EBV-associated lymphoproliferative disorders and skin malignancies have been reported in association with tacrolimus treatment.	
<u>Blood and lymphatic system disorders</u>	
Common:	anaemia, leucopenia, thrombocytopenia, leucocytosis, abnormal red blood cell analysis
Uncommon:	coagulopathies, coagulation and bleeding analyses abnormal, pancytopenia, neutropenia
Rare:	thrombotic thrombocytopenic purpura, hypoprothrombinaemia
<u>Not known:</u>	pure red cell aplasia, agranulocytosis, haemolytic anaemia
<u>Immune system disorders</u>	
Allergic and anaphylactoid reactions have been observed in patients using tacrolimus (see section 4.4)	
<u>Endocrine disorders</u>	
Rare:	Hirsutism
<u>Metabolism and nutrition disorders</u>	
Very common:	hyperglycaemic conditions, diabetes mellitus, hyperkalaemia
Common:	hypomagnesaemia, hypophosphataemia, hypokalaemia, hypocalcaemia, hyponatraemia, fluid overload, hyperuricaemia, appetite decreased, anorexia, metabolic acidoses, hyperlipidaemia, hypercholesterolaemia, hypertriglyceridaemia, other electrolyte abnormalities
Uncommon	dehydration, hypoproteinaemia, hyperphosphataemia, hypoglycaemia
<u>Psychiatric disorders</u>	
Very common:	Insomnia
Common:	anxiety symptoms, confusion and disorientation, depression, depressed mood, mood disorders and disturbances, nightmares, hallucinations, mental disorders
Uncommon:	psychotic disorder
<u>Nervous system disorders</u>	
Very common:	tremor, headache
Common:	seizures, disturbances in consciousness, paraesthesia and dysaesthesia, peripheral neuropathies, dizziness, writing impaired, nervous system disorders
Uncommon:	coma, central nervous system haemorrhages and cerebrovascular accidents, paralysis and paresis, encephalopathy, speech and language abnormalities, amnesia
Rare	hypertonia
Very rare:	myasthenia
<u>Eye disorders</u>	
Common:	blurred vision, photophobia, eye disorders

Uncommon:	cataract
Rare:	blindness
<u>Ear and labyrinth disorders</u>	
Common:	tinnitus
Uncommon:	hypoacusis
Rare:	deafness neurosensory
Very rare:	hearing impaired
<u>Cardiac disorders</u>	
Common:	ischaemic coronary artery disorders, tachycardia
Uncommon:	ventricular arrhythmias and cardiac arrest, heart failures, cardiomyopathies, ventricular hypertrophy, supraventricular arrhythmias, palpitations, ECG investigations abnormal, heart rate and pulse investigations abnormal
Rare:	pericardial effusion
Very rare:	echocardiogram abnormal
<u>Vascular disorders</u>	
Very common:	Hypertension
Common:	haemorrhages, thromboembolic and ischaemic events, peripheral vascular disorders, vascular hypotensive disorders
Uncommon:	infarction, venous thrombosis deep limb, shock
<u>Respiratory, thoracic and mediastinal disorders</u>	
Common:	dyspnoea, parenchymal lung disorders, pleural effusion, pharyngitis, cough, nasal congestion and inflammations
Uncommon:	respiratory failures, respiratory tract disorders, asthma
Rare:	acute respiratory distress syndrome
<u>Gastrointestinal disorders</u>	
Very common:	diarrhoea, nausea
Common:	gastrointestinal inflammatory conditions, gastrointestinal ulceration and perforation, gastrointestinal haemorrhages, stomatitis and ulceration, ascites, vomiting, gastrointestinal and abdominal pain, dyspeptic signs and symptoms, constipation, flatulence, bloating and distension, loose stools, gastrointestinal-signs and symptoms
Uncommon:	ileus paralytic, peritonitis, acute and chronic pancreatitis, blood amylase increased, gastro-oesophageal reflux disease, impaired gastric emptying
Rare:	subileus, pancreatic pseudocysts
<u>Hepatobiliary disorders</u>	
Common:	hepatic enzymes and function abnormalities, cholestasis and jaundice, hepatocellular damage and hepatitis, cholangitis
Rare:	hepatic artery thrombosis, veno-occlusive liver disease
Very rare:	hepatic failure, bile duct stenosis
<u>Skin and subcutaneous disorders</u>	
Common:	pruritus, rash, alopecia, acne, sweating increased
Uncommon:	dermatitis, photosensitivity
Rare:	toxic epidermal necrolysis (Lyell's syndrome)
Very rare:	Stevens-Johnson syndrome
<u>Musculoskeletal and connective tissue disorders</u>	

Common:	arthralgia, muscle cramps, pain in the limb, back pain
Rare:	joint disorders
<u>Renal and urinary disorders</u>	
Very common:	renal <u>impairment</u>
Common:	renal failure, renal failure acute, oliguria, renal tubular necrosis, nephropathy toxic, urinary abnormalities, bladder and urethral symptoms
Uncommon:	anuria, haemolytic uraemic syndrome
Very rare:	nephropathy, haemorrhagic cystitis
<u>Reproductive system and breast disorders</u>	
Uncommon:	dysmenorrhoea and uterine bleeding
<u>General disorders and administration site conditions</u>	
Common:	asthenic conditions, blood alkaline phosphatase increased, febrile disorders, oedema, pain and discomfort, alkaline phosphatase increased, weight increased, body temperature perception disturbed
Uncommon:	multi-organ failure, influenza like illness, temperature intolerance, chest pressure sensation, feeling jittery, feeling abnormal, blood lactate dehydrogenase increased, weight decreased
Rare:	thirst, falls, chest tightness, decreased mobility, ulcers
Very rare:	Fat tissue increased
<u>Injury, poisoning and procedural complications</u>	
Common:	primary graft dysfunction medication errors, including inadvertent, unintentional or unsupervised substitution of immediate- or prolonged-release tacrolimus formulations, have been observed. A number of associated cases of transplant rejection have been reported (frequency cannot be estimated from available data).

4.9 Overdose

Experience with overdose is limited. Several cases, of accidental overdosage have been reported symptoms have included tremor, headache, nausea and vomiting, infections, urticaria, lethargy, increased blood urea nitrogen and elevated serum creatinine concentrations, and increase in alanine aminotransferase levels.

No specific antidote to tacrolimus therapy is available. If overdosage occurs, general supportive measures and symptomatic treatment should be conducted.

Based on its high molecular weight, poor aqueous solubility, and extensive erythrocyte and plasma protein binding, it is anticipated that tacrolimus will not be dialysable. In isolated patients with very high plasma levels, haemofiltration and haemodiafiltration have been effective in reducing toxic concentrations. In cases of oral intoxication, gastric lavage and/or the use of adsorbents (such as activated charcoal) may be helpful, if used shortly after intake.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Calcineurin inhibitors, ATC code: L04AD02

Mechanism of action and pharmacodynamic effects

At the molecular level, the effects of tacrolimus appear to be mediated by binding to a cytosolic protein (FKBP12)

which is responsible for the intracellular accumulation of the compound. The FKBP12-tacrolimus complex specifically and competitively binds to and inhibits calcineurin, leading to a calcium-dependent inhibition of T-cell signal transduction pathways, thereby preventing transcription of a discrete set of lymphokine genes.

Tacrolimus is a highly potent immunosuppressive agent and has proven activity in both *in vitro* and *in vivo* experiments.

In particular, tacrolimus inhibits the formation of cytotoxic lymphocytes, which are mainly responsible for graft rejection. Tacrolimus suppresses T-cell activation and T-helper-cell dependent B-cell proliferation, as well as the formation of lymphokines (such as interleukin-2, interleukin-3, and γ -interferon) and the expression of the interleukin-2 receptor.

Results from published data in other primary organ transplantation

Tacrolimus has evolved into an accepted treatment as a primary immunosuppressive medicinal product following pancreas, lung and intestinal transplantation. In prospective published studies, tacrolimus was investigated as primary immunosuppressant in approximately 175 patients following lung transplantation, 475 patients following pancreas transplantation and 630 patients following intestinal transplantation. Overall, the safety profile of tacrolimus in these published studies appeared to be similar to what was reported in the large studies, where tacrolimus was used as primary treatment in liver, kidney and heart transplantation. Efficacy results of the largest studies in each indication are summarised below.

Lung transplantation

The interim analysis of a recent multi-centre study discussed 110 patients who underwent 1:1 randomisation to either tacrolimus or ciclosporin. Tacrolimus was started as continuous intravenous infusion at a dose of 0.01 to 0.03 mg/kg/day and oral tacrolimus was administered at a dose of 0.05 to 0.3 mg/kg/day. A lower incidence of acute rejection episodes for tacrolimus- versus ciclosporin-treated patients (11.5% versus 22.6%) and a lower incidence of chronic rejection, the bronchiolitis obliterans syndrome (2.86% versus 8.57%) were reported within the first year after transplantation. The one-year patient survival rate was 80.8% in the tacrolimus group and 83% in the ciclosporin group (Treede et al., 3rd ICI San Diego, US, 2004; Abstract 22).

Another randomised study included 66 patients on tacrolimus versus 67 patients on ciclosporin. Tacrolimus was started as a continuous intravenous infusion at a dose of 0.025 mg/kg/day and oral tacrolimus was administered at a dose of 0.15 mg/kg/day with subsequent dose adjustments to target trough levels of 10 to 20 ng/ml. The 1-year patient survival was 83% in the tacrolimus group and 71% in the ciclosporin group, the 2-year survival rates were 76% and 66% respectively. Acute rejection episodes per 100 patient days were numerically fewer in the tacrolimus group (0.85 episodes) than in the ciclosporin group (1.09 episodes). Obliterative bronchiolitis developed in 21.7% of patients in the tacrolimus group compared with 38.0% of patients in the ciclosporin group ($p = 0.025$). Significantly more ciclosporin-treated patients ($n = 13$) required a switch to tacrolimus than tacrolimus-treated patients-to ciclosporin ($n = 2$) ($p = 0.02$) (Keenan et al., *Ann Thoracic Surg* 1995; 60:580).

In an additional two centre study, 26 patients were randomized to the tacrolimus versus 24 patients to the ciclosporin group. Tacrolimus was started as a continuous intravenous infusion at an initial dose of 0.05 mg/kg/day and oral tacrolimus was administered at a dose of 0.1 to 0.3 mg/kg/day with subsequent dose adjustments to target trough levels of 12 to 15 ng/ml. The 1-year survival rates were 73.1% in the tacrolimus versus 79.2% in the ciclosporin group. Freedom from acute rejection was higher in the tacrolimus group at 6 months (57.7% versus 45.8%) and at 1 year after lung transplantation (50% versus 33.3%) (Treede et al., *J Heart Lung Transplant* 2001; 20:511).

The three studies demonstrated similar survival rates. The incidences of acute rejection were numerically lower with tacrolimus in all three studies and one of the studies reported a significantly lower incidence of bronchiolitis obliterans syndrome with tacrolimus.

Pancreas transplantation

A multi-centre study included 205 patients undergoing simultaneous kidney and pancreas transplantation who were randomised to treatment with tacrolimus ($n=103$) or ciclosporin ($n=102$). The initial oral per protocol dose of tacrolimus was 0.2 mg/kg/day with subsequent dose adjustments to target trough levels of 8 to 15 ng/ml by Day 5 and 5 to 10 ng/ml after 6 months. Pancreas survival at 1 year was significantly superior with tacrolimus 91.3% versus 74.5% with ciclosporin ($p < 0.0005$), whereas renal graft survival was similar in both groups. In total 34 patients switched

treatment from ciclosporin to tacrolimus, whereas only 6 tacrolimus patients required alternative therapy (Bechstein et al., Transplantation 2004; 77:1221).

Intestinal transplantation

Published clinical experience from a single centre on the use of tacrolimus for primary treatment following intestinal transplantation showed that the actuarial survival rate of 155 patients (65 intestine alone, 75 liver and intestine, and 25 multivisceral) receiving tacrolimus and prednisone was 75% at 1 year, 54% at 5 years, and 42% at 10 years. In the early years the initial oral dose of tacrolimus was 0.3 mg/kg/day. Results continuously improved with increasing experience over the course of 11 years. A variety of innovations, such as techniques for early detection of Epstein-Barr (EBV) and CMV infections, bone marrow augmentation, the adjunct use of the interleukin-2 antagonist daclizumab, lower initial tacrolimus doses with target trough levels of 10 to 15 ng/ml, and most recently allograft irradiation were considered to have contributed to improved results in this indication over time. (Abu-Elmagd et al., Ann Surg 2001; 234:404).

5.2 Pharmacokinetic properties

Absorption

In man, tacrolimus has been shown to be able to be absorbed throughout the gastrointestinal tract. Following oral administration of Tacrolimus capsules, peak blood concentrations (C_{\max}) of tacrolimus are achieved in approximately 1 - 3 hours. In some patients, tacrolimus appears to be continuously absorbed over a prolonged period, yielding a relatively flat absorption profile. The mean oral bioavailability of tacrolimus is in the range of 20% - 25%.

After oral administration (0.30 mg/kg/day) to liver transplant patients, steady-state concentrations of tacrolimus were achieved within 3 days in the majority of patients.

In healthy subjects, tacrolimus 0.5 mg, tacrolimus 1 mg and tacrolimus 5 mg hard capsules have been shown to be bioequivalent when administered as equivalent dose.

The rate and extent of absorption of tacrolimus is greatest under fasted conditions. The presence of food decreases both the rate and extent of absorption of tacrolimus, the effect being most pronounced after a high fat meal. The effect of a high-carbohydrate meal is less pronounced.

In stable liver transplant patients, the oral bioavailability of Tacrolimus was reduced when it was administered after a meal of moderate fat content (34% of calories). Decreases in AUC (27%) and C_{\max} (50%), and an increase in t_{\max} (173%) in whole blood were evident.

In a study of stable renal transplant patients who were administered tacrolimus immediately after a standard continental breakfast the effect on oral bioavailability was less pronounced. Decreases in AUC (2 to 12%) and C_{\max} (15 to 38%), and an increase in t_{\max} (38 to 80%) in whole blood were evident.

Bile flow does not influence the absorption of tacrolimus.

A strong correlation exists between AUC and whole blood trough levels at steady-state. Monitoring of whole blood trough levels therefore provides a good estimate of systemic exposure.

Distribution and elimination

In man, the disposition of tacrolimus after intravenous infusion may be described as biphasic.

In the systemic circulation, tacrolimus binds strongly to erythrocytes resulting in an approximate 20:1 distribution ratio of whole blood/plasma concentrations in plasma, tacrolimus is highly bound (> 98.8%) to plasma proteins, mainly to serum albumin and alpha-1 acid glycoprotein.

Tacrolimus is extensively distributed in the body. The steady-state volume of distribution based on plasma concentrations is approximately 1300 l (healthy subjects). Corresponding data based on whole blood averaged 47.6 l.

Tacrolimus is a low-clearance substance. In healthy subjects, the average total body clearance (TBC) estimated from whole blood concentrations was 2.25 l/h. In adult liver, kidney and heart transplant patients, values of 4.1 l/h, 6.7 l/h and 3.9 l/h, respectively, have been observed. Paediatric liver transplant recipients have a TBC approximately twice that of adult liver transplant patients. Factors such as low haematocrit and protein levels, which result in an increase in the unbound fraction of tacrolimus, or corticosteroid-induced increased metabolism are considered to be responsible for the higher clearance rates observed following transplantation.

The half-life of tacrolimus is long and variable. In healthy subjects, the mean half-life in whole blood is approximately 43 hours. In adult and paediatric liver transplant patients, it averaged 11.7 hours and 12.4 hours respectively, compared with 15.6 hours in adult kidney transplant recipients. Increased clearance rates contribute to the shorter half-life observed in transplant recipients.

Metabolism and biotransformation

Tacrolimus is widely metabolised in the liver, primarily by cytochrome P450-3A4. Tacrolimus is also considerably metabolised in the intestinal wall. There are several metabolites identified. Only one of these has been shown *in vitro* to have immunosuppressive activity similar to that of tacrolimus. The other metabolites have only weak or no immunosuppressive activity. In systemic circulation only one of the inactive metabolites is present in the systemic circulation at low concentrations. Therefore, metabolites do not contribute to pharmacological activity of tacrolimus.

Excretion

Following intravenous and oral administration of ^{14}C -labelled tacrolimus, most of the radioactivity was eliminated in the faeces. Approximately 2% of the radioactivity was eliminated in the urine. Less than 1% of unchanged tacrolimus was detected in the urine and faeces, indicating that tacrolimus is almost completely metabolised prior to elimination, bile being the principal route of elimination.

5.3 Preclinical safety data

The kidneys and the pancreas were the primary organs affected in toxicity studies performed in rats and baboons. In rats, Tacrolimus caused toxic effects to the nervous system and the eyes. Reversible cardiotoxic effects were observed in rabbits following intravenous administration of tacrolimus.

Embryofoetal toxicity was observed in rats and rabbits and was limited to doses that caused significant toxicity in maternal animals. In rats, female reproductive function, including birth, was impaired at toxic doses and the offspring showed reduced birth weight, viability and growth.

A negative effect of tacrolimus on male fertility in the form of reduced sperm counts and motility was observed in rats.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content:

Lactose monohydrate

Croscarmellose sodium (E468)

Hypromellose (E464)

Magnesium stearate (E470b)

Capsule shell

Gelatin

Titanium dioxide (E 171)

Sodium laurilsulfate

Printing ink of capsule shell: Shellac, propylene glycol, potassium hydroxide, black iron oxide (E172).

6.2 Incompatibilities

Tacrolimus is not compatible with PVC. Tubing, syringes and other equipment used to prepare or administer a suspension of Tacrolimus capsules contents should not contain PVC.

6.3 Shelf life

2 years

6.4 Special precautions for storage

Store below 25°C

Store in the original package in order to protect from moisture.

Hard capsules should be taken immediately following removal from the blister.

6.5 Nature and contents of container

Alu-Alu blister

Packs of 20, 30, 50, 60, 90 and 100 hard capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Accord Healthcare Limited
Sage House, 319 Pinner Road
North Harrow,
Middlesex HAI 4HF
United Kingdom

8 MARKETING AUTHORISATION NUMBER

PA 1390/19/2

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 19 November 2010

10 DATE OF REVISION OF THE TEXT

January 2013