Package leaflet: Information for the user
Naltrexone Hydrochloride 50 mg film-coated tablets
naltrexone hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:
1. What Naltrexone Hydrochloride is and what it is used for
2. What you need to know before you take Naltrexone Hydrochloride
3. How to take Naltrexone Hydrochloride
4. Possible side effects
5. How to store Naltrexone Hydrochloride
6. Contents of the pack and other information

1. What Naltrexone Hydrochloride is and what it is used for

The active ingredient, naltrexone hydrochloride, belongs to a group of medicines other nervous system drugs; drugs used in addictive disorders

What is Naltrexone Hydrochloride used for
Naltrexone hydrochloride is used in combination with other medicines or therapy to help those who are dependent on opioids drugs overcome their addiction

Naltrexone acts by blocking receptors in the brain to block the action of opioids. Individuals will no longer experience the euphoria previously experienced after taking opioids.

Naltrexone hydrochloride is used with a comprehensive treatment program to help those who are dependent on alcohol maintain abstinence (self-denial).

Naltrexone Hydrochloride film-coated tablets does not cause dependency.

2. What you need to know before you take Naltrexone Hydrochloride

Do not take Naltrexone Hydrochloride
- if you are allergic to naltrexone hydrochloride or any of the other ingredients of this medicine (listed in section 6).
- if you have severe kidney problems
- if you have severe liver problems
- if you have an acute liver infection
- if you are dependent on opiates
- if your urine tests positive for opiates
- if you experience withdrawal symptoms after a naloxone injection
- if you are using a medicinal product containing an opioid, for example certain cough medicines, medicines to treat diarrhoea (such as kaolin and morphine) and analgesics (pain killers). Note: Naltrexone hydrochloride does not have a blocking effect on analgesics which do not contain any opioids (such as ibuprofen, paracetamol and acetylsalicylic acid).
- if you take methadone.

If you think any of these apply to you, do not take the tablets. Talk to your doctor first and follow his advice.

Warnings and precautions
Talk to your doctor or pharmacist before taking Naltrexone Hydrochloride

- **Do not** take opiates whilst taking Naltrexone Hydrochloride. Although Naltrexone Hydrochloride will normally block some of the effects (i.e. the highs), if you take high doses of opiates, you may experience breathing difficulties and problems with your circulation (opiate poisoning).
- You should not use Naltrexone Hydrochloride if you are still addicted to opiates as Naltrexone Hydrochloride will cause severe withdrawal symptoms in this situation.
- You must inform every doctor that treats you that you are taking Naltrexone Hydrochloride. Non-opiate based anaesthetics should be used if you require an anaesthetic in an emergency situation. If you have to use opiate containing anaesthetics, you may need higher doses than usual. You may also be more sensitive to the side-effects (breathing difficulties and circulatory problems).
- You must not try to overcome the blocking effect of Naltrexone Hydrochloride with high doses of opiates. There is a risk that the opiates could still be in your body after the effects of Naltrexone Hydrochloride have passed. If this occurs, you could unintentionally overdose with serious consequences.
- Naltrexone is removed from the body by the liver and kidney. Liver problems are common in opiate-dependant individuals. Your doctor will carry out liver function tests before and during treatment.
- If you become pregnant you have to inform your doctor.
- If you go for any blood tests, inform the doctor carrying out the test that you are taking Naltrexone Hydrochloride as these tablets can affect the results of liver function tests.

Consult your doctor if one of the above warnings applies to you or has done so in the past.

Children and adolescents
Naltrexone Hydrochloride should not be used in children and adolescents under 18 years of age, since clinical data in this age-group are lacking. Safe use in children has not been established.

Use in older people
There are insufficient data on the safety and efficacy of naltrexone for this indication in elderly patients.

Other medicines and Naltrexone Hydrochloride
Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.
Some common medicines contain opiates and these may not work when you are taking Naltrexone Hydrochloride. You should inform your doctor if you need cough-mixtures or medicines against diarrhoea or pain since these may contain opiates and the use in conjunction with an opioid-containing medication is contraindicated.
If, despite the contraindication to use in conjunction, opioid containing drugs are needed in emergency cases the suitable dose for pain relief can be higher as usual. Close monitoring through the doctor is absolutely necessary because occurring respiratory depression and other symptoms may be stronger and longer-lasting.

Naltrexone Hydrochloride with food and drink
Taking food and drink has no influence on your treatment with Naltrexone Hydrochloride.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.
The safety of using Naltrexone Hydrochloride during pregnancy has not been demonstrated.

It is not known whether naltrexone is excreted in breast milk. Because the safety of using naltrexone in neonates and children has not been demonstrated, breast-feeding is not advised while using Naltrexone Hydrochloride.

Driving and using machines
Naltrexone may impair the mental and/or physical abilities required for performance of potentially hazardous tasks such as driving a car or operating machinery.

Naltrexone Hydrochloride contains lactose monohydrate
If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Naltrexone Hydrochloride

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 1 tablet per day unless a different dose has been prescribed by your doctor.

- Naltrexone Hydrochloride taken orally with small amount of liquid.
- Before starting to take Naltrexone Hydrochloride, you must not have used any other opiates for at least 7-10 days. Your doctor can use a test to establish whether you are clear of these drugs before you start the treatment. Generally speaking, treatment begins at a dose of 1/2 tablet per day (25 mg), later increased to 1 tablet per day (50 mg).
- Naltrexone Hydrochloride must be used exclusively for the disorder for which your doctor has prescribed this medicine.
- It is important to follow your doctor's instructions closely with respect to the dosage.
- It is important that you take Naltrexone Hydrochloride for the period of time prescribed by your doctor. The treatment can last for three months or longer, according to the judgment of your doctor. Naltrexone Hydrochloride should be combined with other forms of treatment.

If you notice that the effect of Naltrexone Hydrochloride is too strong or not strong enough, consult your doctor or pharmacist.

If you take more Naltrexone Hydrochloride than you should
If you have taken more than the prescribed number of tablets, you should inform your doctor immediately.

If you forget to take Naltrexone Hydrochloride
You can still take the Naltrexone Hydrochloride when you remember.
Do not take a double dose to make up for a forgotten dose.

If you stop taking Naltrexone Hydrochloride
If you consider stopping before the end of the agreed period of treatment, always discuss this with your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.
Naltrexone Hydrochloride can affect your liver function. Your doctor may carry out blood test before you start treatment and at various times during treatment to monitor your liver function.

If you notice any of the following, **stop taking** Naltrexone Hydrochloride and contact your doctor **immediately:**

- Abdominal pain lasting more than a few days
- White bowel movements
- Dark urine
- Yellowing of your eyes

As these may be signs that your liver isn't working well.

If you notice any of the following, tell your doctor immediately:

- Swelling of the face, lips, or tongue
- Skin rash
- Difficulty breathing

As these may be signs of an allergic reaction

Very common (may affect more than 1 in 10 people)

- Difficulty sleeping
- Anxiety or nervousness
- Abdominal pain
- Feeling sick and/or being sick
- Lack of energy or strength
- Joint and/or muscle pain
- Headaches
- Restlessness

Common (may affect up to 1 in 10 people)

- Irritability
- Increased energy
- Dizziness
- Increased or excessive sweating
- Increased tear secretion
- Increased heart rate
- Heart Palpitations
- Change in ECG readings
- Pain in the chest
- Diarrhoea
- Constipation
- Rash
- Delayed ejaculation
- Erectile dysfunction
- Decreased appetite
- Thirst
- Affective disorders
- Chills

Uncommon (may affect up to 1 in 100 people)

- Some infections (e.g. Oral herpes, athlete's foot)
- Swollen/enlarged lymph nodes
- Hallucinations
- Confusional state
- Depression
- Paranoia
- Disorientation
- Nightmare
- Agitation
- Reduced libido
- Abnormal dreams
- Tremor

- Drowsiness
- Blurred vision
- Irritation in eye
- Abnormal intolerance to visual perception of light
- Swelling of eyes
- Eye pain or tiredness
- Strain in eye
- Ear discomfort
- Ear pain
- Ringing of ear
- Vertigo
- Blood pressure fluctuation
- Hot flushes
- Nasal congestion & discomfort
- Sneezing
- Disorders of the mouth and throat
- Sputum increased
- Sinus problems
- Voice disorders
- Shortness of breath/difficulty in breathing
- Coughing
- Yawning
- Runny nose
- Flatulence
- Piles
- Ulcer
- Dry mouth
- Liver disorders (including inflammation of liver)
- Increase in bilirubin level
- Hepatitis
- Seborrhea (mild dermatitis)
- Acne
- Hair loss
- Itching
- Groin pain
- Increased urination
- Painful urination
- Increased appetite
- Weight loss
- Weight gain
- Fever
- Pain
- Sensation of cold in extremities
- Feeling hot

Rare (may affect up to 1 in 1,000 people)
- Decreased number of blood platelets
- Suicidal thoughts
- Attempt to suicide

Very rare (may affect up to 1 in 10,000 people)
- Breakdown of skeletal muscle tissue

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via
HPRA Pharmacovigilance

Website: www.hpra.ie
By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Naltrexone Hydrochloride

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date, which is stated on the label after 'EXP'. The expiry date refers to the last day of that month.
- This medicinal product does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Naltrexone Hydrochloride contains

The active substance is naltrexone hydrochloride.

Each film-coated tablet contains 50 mg naltrexone hydrochloride.

The other ingredients are:
Core Tablet: lactose monohydrate, cellulose microcrystalline, crospovidone, colloidal anhydrous silica, magnesium stearate

Film-coating: hypromellose (E464), macrogol 400, polysorbate 80 (E433), iron oxide yellow (E172), iron oxide red (E172), titanium dioxide (E171)

What Naltrexone Hydrochloride looks like and contents of the pack

Naltrexone Hydrochloride are available as yellow coloured, oval, biconvex, film coated tablets with breakline on one side and plain on other side.
The tablet can be divided into equal halves.

Naltrexone Hydrochloride are available in white opaque PVC/PE/Aclar-Alu blister and Alu –Alu blister packs containing 7, 14, 28, 30, 50 and 56 tablets.
Not all pack sizes may be marketed.

Marketing Authorisation holder & Manufacturer:

Marketing Authorisation holder
Accord Healthcare Ireland Limited,
Euro House, Euro Business Park,
Little Island,
Cork T45 K857,
Ireland

Manufacturer:
Accord Healthcare Polska Sp.z o.o.,
ul. Lutomierska 50,95-200 Pabianice, Poland

Accord Healthcare B.V.,
Winthontlaan 200,
3526 KV Utrecht,
The Netherlands

NL/H/1151/001/II/030, version 01, November 2026

Accord Healthcare Single Member S.A.
64th Km National Road Athens, Lamia, 32009, Greece

This medicinal product is authorised in the Member States of the European Economic Area and the United Kingdom (Northern Ireland) under the following names:

Name of the Member State	**Name of the medicine**
Netherland	Naltrexon Hydrochloride Accord 50 mg filmomhulde tabletten
Belgium	Naltrexone Accord 50 mg filmomhuldetablet
Denmark	Naltrexon Accord
Estonia	Naltrexone Accord
Spain	Naltrexona Accord 50 mg comprimidos recubiertos con película
Finland	Naltreksonihydrokloridi Accord 50 mg Kalvopäävysteinen Tabletti
Germany	Naltrexonhydrochlorid Accord 50 mg Filmtabletten
Ireland	Naltrexone Hydrochloride 50 mg Film coated Tablets
Italy	Naltrexone Accord Healthcare 50 mg compresse rivestite con film
Lithuania	Naltrexone Hydrochloride Accord 50 mg plėvele dengtos tabletės
Latvia	Naltrexone Hydrochloride Accord 50 mg Film coated Tablets
Norway	Naltrexon Accord 50 mg Filmdrasjert tablett
Portugal	Naltrexona Hydrochloride Accord
Poland	Naltrexona Hydrochloride Accord
United Kingdom (Northern Ireland)	Naltrexone Hydrochloride 50 mg Film coated Tablets
Austria	Naltrexone Accord 50 mg Filmtabletten
Bulgaria	Naltrexone Акорд 50 мг филмирани таблетки
Cyprus	Naltrexone Hydrochloride Accord 50 mg Film coated Tablets
Sweden	Naltrexone Accord 50 mg filmdragerad tablet

This leaflet was last revised in November 2026.