

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Topotecan Accord 1mg/ml concentrate for solution for infusion

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

1 ml of concentrate for solution for infusion contains 1 mg topotecan (as hydrochloride).

Each 1 ml vial of concentrate contains 1 mg topotecan (as hydrochloride)

Each 4 ml vial of concentrate contains 4 mg topotecan (as hydrochloride)

For a full list of excipients see section 6.1.

## 3 PHARMACEUTICAL FORM

Concentrate for solution for infusion.

A clear yellow colour solution free from visible foreign particles. pH in the range of 1.5 to 2.5 and Osmolarity in the range of approximately 100 to 40 mOsm/Litre

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Topotecan monotherapy is indicated for the treatment of:

- Patients with metastatic carcinoma of the ovary after failure of first-line or subsequent therapy.
- Patients with relapsed small cell lung cancer [SCLC] for whom re-treatment with the first- line regimen is not considered appropriate (see section 5.1).

Topotecan in combination with cisplatin is indicated for patients with carcinoma of the cervix recurrent after radiotherapy and for patients with Stage IVB disease. Patients with prior exposure to cisplatin require a sustained treatment-free interval to justify treatment with the combination (see section 5.1).

### 4.2 Posology and method of administration

The use of topotecan should be confined to units specialised in the administration of cytotoxic chemotherapy. Topotecan should only be administered under the supervision of a physician experienced in the use of chemotherapy (see section 6.6).

#### **Posology**

When topotecan is used in combination with cisplatin, the full prescribing information for cisplatin should be consulted. Prior to administration of the first course of topotecan, patients must have a baseline neutrophil count of  $\geq 1.5 \times 10^9/l$ , a platelet count of  $\geq 100 \times 10^9/l$  and a haemoglobin level of  $\geq 9$  g/dl (after transfusion if necessary).

#### **Ovarian and Small Cell Lung Carcinoma**

##### **Initial dose**

The recommended dose of topotecan is  $1.5 \text{ mg/m}^2$  body surface area/day administered by intravenous infusion over 30 minutes daily for 5 consecutive days with a 3 week interval between the start of each course. If well tolerated, treatment may continue until disease progression (see sections 4.8 and 5.1).

##### **Subsequent doses**

Topotecan should not be re-administered unless the neutrophil count is  $\geq 1 \times 10^9/l$ , the platelet count is  $\geq 100 \times 10^9/l$ , and the haemoglobin level is  $\geq 9$  g/dl (after transfusion if necessary).

Standard oncology practice for the management of neutropenia is either to administer topotecan with other medicinal products (e.g. G-CSF) or to reduce the dose to maintain neutrophil counts.

If dose reduction is chosen for patients who experience severe neutropenia (neutrophil count  $< 0.5 \times 10^9/l$ ) for 7 days or more or severe neutropenia associated with fever or infection, or who have had treatment delayed due to neutropenia, the dose should be reduced by  $0.25 \text{ mg/m}^2/\text{day}$  to  $1.25 \text{ mg/m}^2/\text{day}$  (or subsequently down to  $1.0 \text{ mg/m}^2/\text{day}$  if necessary).

Doses should be similarly reduced if the platelet count falls below  $25 \times 10^9/l$ . In clinical studies, topotecan was discontinued if the dose had been reduced to  $1.0 \text{ mg/m}^2/\text{day}$  and a further dose reduction was required to manage adverse effects.

### **Cervical Carcinoma**

#### **Initial dose**

The recommended dose of topotecan is  $0.75 \text{ mg/m}^2/\text{day}$  administered as a 30 minute intravenous infusion on days 1, 2 and 3. Cisplatin is administered as an intravenous infusion on day 1 at a dose of  $50 \text{ mg/m}^2/\text{day}$  and following the topotecan dose. This treatment schedule is repeated every 21 days for 6 courses or until progressive disease.

#### **Subsequent doses**

Topotecan should not be re-administered unless the neutrophil count is  $\geq 1.5 \times 10^9/l$ , the platelet count is  $\geq 100 \times 10^9/l$ , and the haemoglobin level is  $\geq 9$  g/dl (after transfusion if necessary).

Standard oncology practice for the management of neutropenia is either to administer topotecan with other medicinal products (e.g. G-CSF) or to reduce the dose to maintain neutrophil counts.

If dose reduction is chosen for patients who experience severe neutropenia (neutrophil count  $< 0.5 \times 10^9/l$ ) for 7 days or more or severe neutropenia associated with fever or infection or who have had treatment delayed due to neutropenia, the dose should be reduced by 20% to  $0.60 \text{ mg/m}^2/\text{day}$  for subsequent courses (or subsequently down to  $0.45 \text{ mg/m}^2/\text{day}$  if necessary).

Doses should be similarly reduced if the platelet count falls below  $25 \times 10^9/l$ .

### **Special populations**

#### *Patients with renal impairment*

#### **Monotherapy (Ovarian and Small cell lung carcinoma)**

There is insufficient experience with the use of topotecan in patients with severely impaired renal function (creatinine clearance  $< 20$  ml/min). Use of topotecan in this group of patients is not recommended (see section 4.4).

Limited data indicate that the dose should be reduced in patients with moderate renal impairment. The recommended monotherapy dose of topotecan in patients with ovarian or small cell lung carcinoma and a creatinine clearance between 20 and 39 ml/min is  $0.75 \text{ mg/m}^2/\text{day}$  for 5 consecutive days.

#### **Combination therapy (Cervical carcinoma):**

In clinical studies with topotecan in combination with cisplatin for the treatment of cervical cancer, therapy was only initiated in patients with serum creatinine less than or equal to  $1.5 \text{ mg/dL}$ . If, during topotecan/cisplatin combination therapy serum creatinine exceeds  $1.5 \text{ mg/dL}$ , it is recommended that the full prescribing information be consulted for any advice on cisplatin dose reduction/continuation. If cisplatin is discontinued, there are insufficient data regarding continuing monotherapy with topotecan in patients with cervical cancer.

#### *Patients with hepatic impairment*

A small number of hepatically impaired patients (serum bilirubin between  $1.5$  and  $10 \text{ mg/dl}$ ) were given intravenous topotecan at  $1.5 \text{ mg/m}^2/\text{day}$  for five days every three weeks. A reduction in topotecan clearance was observed. However, there are insufficient data available to make a dose recommendation for this patient group (see section 4.4).

There is insufficient experience with the use of topotecan in patients with severely impaired hepatic function (serum bilirubin  $\geq 10 \text{ mg/dl}$ ) due to cirrhosis. Topotecan is not recommended to be used in this patient group (see section 4.4).

**Paediatric population**

Currently available data are described in sections 5.1 and 5.2 but no recommendation on a posology can be made.

**Method of Administration**

Topotecan must be further diluted before use (see section 6.6).

**4.3 Contraindications**

- Severe hypersensitivity to the active substance or to any of the excipients
- Breast-feeding (see section 4.6)
- Severe bone marrow depression prior to starting first course, as evidenced by baseline neutrophils  $< 1.5 \times 10^9/l$  and/or a platelet count of  $< 100 \times 10^9/l$ .

**4.4 Special warnings and precautions for use**

Haematological toxicity is dose-related and full blood count including platelets should be determined regularly (see section 4.2).

As with other cytotoxic medicinal products, topotecan can cause severe myelosuppression. Myelosuppression leading to sepsis and fatalities due to sepsis have been reported in patients treated with topotecan (see section 4.8).

Topotecan-induced neutropenia can cause neutropenic colitis. Fatalities due to neutropenic colitis have been reported in clinical studies with topotecan. In patients presenting with fever, neutropenia, and a compatible pattern of abdominal pain, the possibility of neutropenic colitis should be considered.

Topotecan has been associated with reports of interstitial lung disease (ILD), some of which have been fatal (see section 4.8). Underlying risk factors include history of ILD, pulmonary fibrosis, lung cancer, thoracic exposure to radiation and use of pneumotoxic substances and/or colony stimulating factors. Patients should be monitored for pulmonary symptoms indicative of ILD (e.g. cough, fever, dyspnoea and/or hypoxia), and topotecan should be discontinued if a new diagnosis of ILD is confirmed.

Topotecan monotherapy and topotecan in combination with cisplatin are commonly associated with clinically relevant thrombocytopenia. This should be taken into account when prescribing topotecan, e.g. if patients at increased risk of tumour bleeds are considered for therapy.

As would be expected, patients with poor performance status ( $PS > 1$ ) have a lower response rate and an increased incidence of complications such as fever, infection and sepsis (see section 4.8). Accurate assessment of performance status at the time therapy is given is important, to ensure that patients have not deteriorated to PS 3.

There is insufficient experience of the use of topotecan in patients with severely impaired renal function (creatinine clearance  $< 20$  ml/min) or severely impaired hepatic function (serum bilirubin  $\geq 10$  mg/dl) due to cirrhosis. Use of topotecan in these patient groups is not recommended (see section 4.2).

A small number of hepatically impaired patients (serum bilirubin between 1.5 and 10 mg/dl) were given intravenous topotecan at  $1.5 \text{ mg/m}^2$  /day for five days every three weeks. A reduction in topotecan clearance was observed. However, there are insufficient data available to make a dose recommendation for this patient group (see section 4.2).

Topotecan Accord contains sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially "sodium free". However, if a solution of common salt (0.9% w/v sodium chloride solution) is used for the dilution of Topotecan Accord prior to administration then the dose of sodium received would be higher.

**4.5 Interaction with other medicinal products and other forms of interaction**

No *in vivo* human pharmacokinetic interaction studies have been performed.

Topotecan does not inhibit human P450 enzymes (see section 5.2). In a population study using the intravenous route, the co-administration of granisetron, ondansetron, morphine or corticosteroids did not appear to have a significant effect on the pharmacokinetics of total topotecan (active and inactive form).

When combining topotecan with other chemotherapy agents, reduction of the doses of each medicinal product may be required to improve tolerability. However, when combining with platinum agents, there is a distinct sequence-dependent interaction depending on whether the platinum agent is given on day 1 or 5 of the topotecan dosing. If either cisplatin or carboplatin is given on day 1 of the topotecan dosing, a lower dose of each agent must be given to improve tolerability compared to the dose of each agent which can be given if the platinum agent is given on day 5 of the topotecan dosing.

When topotecan (0.75 mg/m<sup>2</sup>/day for 5 consecutive days) and cisplatin (60 mg/m<sup>2</sup>/day on Day 1) were administered in 13 patients with ovarian cancer, a slight increase in AUC (12%, n=9) and C<sub>max</sub> (23%, n=11) was noted on day 5. This increase is considered unlikely to be of clinical relevance.

#### **4.6 Fertility, pregnancy and lactation**

##### Women of childbearing potential / Contraception in males and females

Topotecan has been shown to cause embryo-foetal lethality and malformations in preclinical studies (see section 5.3). As with other cytotoxic medicinal products, topotecan may cause foetal harm and therefore women of child bearing potential should be advised to avoid becoming pregnant during therapy with topotecan.

As with all cytotoxic chemotherapy, patients being treated with topotecan must be advised that they or their partner must use an effective method of contraception.

Women of childbearing potential should use effective contraceptive measures while being treated with topotecan and for 6 months following completion of treatment.

Men are recommended to use effective contraceptive measures and to not father a child while receiving topotecan and for 3 months following completion of treatment.

##### Pregnancy

If topotecan is used during pregnancy, or if the patient becomes pregnant during therapy with topotecan, the patient must be warned of the potential hazards to the foetus.

##### Breastfeeding

Topotecan is contraindicated during breast-feeding (see section 4.3). Although it is not known whether topotecan is excreted in human breast milk, breast-feeding should be discontinued at the start of therapy.

##### Fertility

No effects on male or female fertility have been observed in reproductive toxicity studies in rats (see section 5.3). However, as with other cytotoxic medicinal products topotecan is genotoxic and effects on fertility, including male fertility, cannot be excluded

#### **4.7 Effects on ability to drive and use machines**

No studies of the effects on the ability to drive and use machines have been performed. However, caution should be observed when driving or operating machines if fatigue and asthenia persist.

#### **4.8 Undesirable effects**

In dose-finding studies involving 523 patients with relapsed ovarian cancer and 631 patients with relapsed small cell lung cancer, the dose-limiting toxicity of topotecan monotherapy was found to be haematological. Toxicity was predictable and reversible. There were no signs of cumulative haematological or non-haematological toxicity.

The safety profile of topotecan when given in combination with cisplatin in the cervical cancer clinical studies is consistent with that seen with topotecan monotherapy. The overall haematological toxicity is lower in patients treated with topotecan in combination with cisplatin compared to topotecan monotherapy, but higher than with cisplatin alone.

Additional adverse events were seen when topotecan was given in combination with cisplatin, however, these events were seen with cisplatin monotherapy and were not attributable to topotecan. The prescribing information for cisplatin should be consulted for a full list of adverse events associated with cisplatin use.

The integrated safety data for topotecan monotherapy are presented below.

Adverse reactions are listed below, by system organ class and absolute frequency (all reported events). Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ) and not known (cannot be estimated from the available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

<b>Infections and infestations</b>	
Very common	Infection
Common	Sepsis <sup>1</sup>
<b>Blood and lymphatic system disorders</b>	
Very common	Febrile neutropenia, neutropenia (see "Gastrointestinal disorders"), thrombocytopenia, anaemia, leucopenia
Common	Pancytopenia
Not known	Severe bleeding (associated with thrombocytopenia)
<b>Immune system disorders</b>	
Common	Hypersensitivity reaction including rash
Rare	Anaphylactic reaction, angioedema, urticaria
<b>Metabolism and nutrition disorders</b>	
Very common	Anorexia (which may be severe)
<b>Respiratory, thoracic and mediastinal disorders</b>	
Rare	Interstitial lung disease (some cases have been fatal)
<b>Gastrointestinal disorders</b>	
Very common	Nausea, vomiting and diarrhoea (all of which may be severe), constipation, abdominal pain <sup>2</sup> , mucositis
Not known	Gastrointestinal perforation
<b>Hepatobiliary disorders</b>	
Common	Hyperbilirubinaemia
<b>Skin and subcutaneous tissue disorders</b>	
Very common	Alopecia
Common	Pruritus
<b>General disorders and administration site conditions</b>	
Very common	Pyrexia, asthenia, fatigue
Common	Malaise
Very rare	Extravasation <sup>3</sup>
Not known	Mucosal inflammation
<ol style="list-style-type: none"> <li>1. Fatalities due to sepsis have been reported in patients treated with topotecan (see section 4.4).</li> <li>2. Neutropenic colitis, including fatal neutropenic colitis, has been reported to occur as a complication of topotecan-induced neutropenia (see section 4.4).</li> <li>3. Reactions have been mild and have not generally required specific therapy.</li> </ol>	

The adverse events listed above have the potential to occur with a higher frequency in patients who have a poor performance status (see section 4.4).

The frequencies associated with the haematological and non-haematological adverse events listed below represent the adverse event reports considered to be related/possibly related to topotecan therapy.

#### *Haematological*

**Neutropenia:** Severe (neutrophil count  $< 0.5 \times 10^9/l$ ) during course 1 in 55 % of patients and with duration  $\geq 7$  days in 20% and overall in 77% of patients (39% of courses). In association with severe neutropenia, fever or infection occurred in 16% of patients during course 1 and overall in 23% of patients (6% of courses). Median time to onset of severe neutropenia was 9 days and the median duration was 7 days. Severe neutropenia lasted beyond 7 days in 11% of courses overall. Among all patients treated in clinical studies (including both those with severe neutropenia and those who did not develop severe neutropenia), 11% (4% of courses) developed fever and 26% (9% of courses) developed infection. In addition, 5% of all patients treated (1% of courses) developed sepsis (see section 4.4).

*Thrombocytopenia:* Severe (platelets  $25 \times 10^9/l$ ) in 25 % of patients (8% of courses); moderate (platelets between 25.0 and  $50.0 \times 10^9/l$ ) in 25 % of patients (15% of courses). Median time to onset of severe thrombocytopenia was Day 15 and the median duration was 5 days. Platelet transfusions were given in 4% of courses. Reports of significant sequelae associated with thrombocytopenia including fatalities due to tumour bleeds have been infrequent.

*Anaemia:* Moderate to severe (Hb  $\leq 8.0$  g/dl) in 37% of patients (14% of courses). Red cell transfusions were given in 52% of patients (21% of courses).

#### Non-haematological

Frequently reported non-haematological effects were gastrointestinal such as nausea (52%), vomiting (32%), diarrhoea (18%), constipation (9%) and mucositis (14%). The incidence of severe (Grade 3 or 4) nausea, vomiting, diarrhoea and mucositis was 4, 3, 2 and 1 % respectively.

Mild abdominal pain was reported in 4% of patients.

Fatigue was observed in approximately 25 % and asthenia in 16 % of patients receiving topotecan. Severe (Grade 3 or 4) fatigue and asthenia both occurred with an incidence of 3 %.

Total or pronounced alopecia was observed in 30% of patients and partial alopecia in 15% of patients.

Other severe events that were recorded as related or possibly related to topotecan treatment were anorexia (12%), malaise (3%) and hyperbilirubinaemia (1%).

Hypersensitivity reactions including rash, urticaria, angioedema and anaphylactic reactions have been reported rarely. In clinical studies, rash was reported in 4% of patients and pruritus in 1.5% of patients.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRC Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie)

## **4.9 Overdose**

Overdoses have been reported in patients being treated with intravenous topotecan (up to 10 fold of the recommended dose) and topotecan capsules (up to 5 fold of the recommended dose). The signs and symptoms observed following overdose were consistent with the known undesirable events associated with topotecan (see section 4.8). The primary complications of overdose are bone marrow suppression and mucositis. In addition, elevated hepatic enzymes have been reported with intravenous topotecan overdose.

There is no known antidote for topotecan overdose. Further management should be as clinically indicated or as recommended by the national poisons center, where available.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmaco-therapeutic group: antineoplastic agents, plant alkaloids and other natural products.

ATC-code: L01CE01

#### Mechanism of action

The anti-tumour activity of topotecan involves the inhibition of topoisomerase-I, an enzyme intimately involved in DNA replication as it relieves the torsional strain introduced ahead of the moving replication fork. Topotecan inhibits topoisomerase-I by stabilising the covalent complex of enzyme and strand-cleaved DNA which is an intermediate of the catalytic mechanism. The cellular sequela of inhibition of topoisomerase-I by topotecan is the induction of protein-associated DNA single-strand breaks.

Clinical efficacy and safety*Relapsed Ovarian Cancer*

In a comparative study of topotecan and paclitaxel in patients previously treated for ovarian carcinoma with platinum based chemotherapy (n = 112 and 114, respectively), the response rate (95% CI) was 20.5% (13%, 28%) versus 14% (8%, 20%) and median time to progression 19 weeks versus 15 weeks (hazard ratio 0.7 [0.6, 1.0]), for topotecan and paclitaxel, respectively. Median overall survival was 62 weeks for topotecan versus 53 weeks for paclitaxel (hazard ratio 0.9 [0.6, 1.3]).

The response rate in the whole ovarian carcinoma programme (n = 392, all previously treated with cisplatin or cisplatin and paclitaxel) was 16%. The median time to response in clinical studies was 7.6-11.6 weeks. In patients refractory to, or relapsing within 3 months after cisplatin therapy (n = 186), the response rate was 10%.

These data should be evaluated in the context of the overall safety profile of the medicinal product, in particular of the significant haematological toxicity (see section 4.8).

A supplementary retrospective analysis was conducted on data from 523 patients with relapsed ovarian cancer. Overall, 87 complete and partial responses were observed, with 13 of these occurring during cycles 5 and 6 and 3 occurring thereafter. Of the patients who received more than 6 cycles of therapy, 91% completed the study as planned or were treated until disease progression with only 3% withdrawn for adverse events.

*Relapsed SCLC*

A phase III study (Study 478) compared oral topotecan plus Best Supportive Care [BSC] [n=71] with BSC alone [n=70] in patients who had relapsed following first line therapy [median time to progression [TTP] from first-line therapy: 84 days for oral topotecan + BSC, 90 days for BSC alone] and for whom retreatment with i.v. chemotherapy was not considered appropriate. In the oral topotecan plus BSC group there was a statistically significant improvement in overall survival compared with the BSC alone group (Logrank p=0.0104). The unadjusted hazard ratio for the oral topotecan plus BSC group relative to the BSC alone group was 0.64 (95% CI: 0.45, 0.90). Median survival in patients treated with oral topotecan + BSC was 25.9 weeks [95 % C.I. 18.3, 31.6] compared to 13.9 weeks [95 % C.I. 11.1, 18.6] for patients receiving BSC alone [p=0.0104].

Patient self-reports of symptoms using an unblinded assessment showed a consistent trend for symptom benefit for oral topotecan + BSC.

One Phase II study (Study 065) and one Phase III study (Study 396) were conducted to evaluate the efficacy of oral topotecan versus intravenous topotecan in patients who had relapsed  $\geq 90$  days after completion of one prior regimen of chemotherapy (see Table 1). Oral and intravenous topotecan were associated with similar symptom palliation in patients with relapsed sensitive SCLC in patient self-reports on an unblinded symptom scale assessment in each of these two studies.

**Table 1. Summary of survival, response rate, and time to progression in SCLC patients treated with oral topotecan or intravenous topotecan**

	<b>Study 065</b>		<b>Study 396</b>	
	<b>Oral topotecan</b>	<b>Intravenous topotecan</b>	<b>Oral topotecan</b>	<b>Intravenous topotecan</b>
	<b>(N = 52)</b>	<b>(N = 54)</b>	<b>(N = 153)</b>	<b>(N = 151)</b>
<b>Median survival (weeks)</b>	32.3	25.1	33.0	35.0
(95% CI)	(26.3, 40.9)	(21.1, 33.0)	(29.1, 42.4)	(31.0, 37.1)
Hazard ratio (95% CI)	0.88 (0.59, 1.31)		0.88 (0.7, 1.11)	
<b>Response rate (%)</b>	23.1	14.8	18.3	21.9
(95% CI)	(11.6, 34.5)	(5.3, 24.3)	(12.2, 24.4)	(15.3, 28.5)
<b>Difference in response rate (95% CI)</b>	8.3 (-6.6, 23.1)		-3.6 (-12.6, 5.5)	
<b>Median time to progression (weeks)</b>	14.9	13.1	11.9	14.6
(95% CI)	(8.3, 21.3)	(11.6, 18.3)	(9.7, 14.1)	(13.3, 18.9)
Hazard ratio (95% CI)	0.90 (0.60, 1.35)		1.21 (0.96, 1.53)	

N = total number of patients treated.

CI = Confidence interval.

In another randomised Phase III study which compared intravenous (IV) topotecan to cyclophosphamide, doxorubicin and vincristine (CAV) in patients with relapsed, sensitive SCLC, the overall response rate was 24.3% for topotecan compared to 18.3% for the CAV group. Median time to progression was similar in the two groups (13.3 weeks and 12.3 weeks respectively). Median survivals for the two groups were 25.0 and 24.7 weeks respectively. The hazard ratio for survival with IV topotecan relative to CAV was 1.04 (95% CI: 0.78, 1.40).

The response rate to topotecan in the combined small cell lung cancer programme [n = 480] for patients with relapsed disease sensitive to first-line therapy, was 20.2 %. The median survival was 30.3 weeks (95 % CI: 27.6, 33.4).

In a population of patients with refractory SCLC (those not responding to first-line therapy), the response rate to topotecan was 4.0%.

#### *Cervical Carcinoma*

In a randomised, comparative Phase III trial conducted by the Gynaecological Oncology Group (GOG 0179), topotecan plus cisplatin (n=147) was compared with cisplatin alone (n=146) for the treatment of histologically confirmed persistent, recurrent or Stage IVB carcinoma of the cervix where curative treatment with surgery and/or radiation was not considered appropriate. Topotecan plus cisplatin had a statistically significant benefit in overall survival relative to cisplatin monotherapy after adjusting for interim analyses (Log-rank p =0.033).

**Table 2: Study results Study GOG-0179**

<b>ITT population</b>		
	<b>Cisplatin 50mg/m<sup>2</sup> on day 1, every 21 days</b>	<b>Cisplatin 50mg/m<sup>2</sup> on day 1 + Topotecan 0.75mg/m<sup>2</sup> on days 1-3, every 21 days</b>
<b>Survival (months)</b>	<b>(n= 146)</b>	<b>(n = 147)</b>
Median (95% CI)	6.5 (5.8, 8.8)	9.4 (7.9, 11.9)
Hazard ratio (95% CI)	0.76 (0.59-0.98)	
Log rank p-value	0.033	
<b>Patients without Prior Cisplatin Chemoradiotherapy</b>		
	<b>Cisplatin</b>	<b>Topotecan/Cisplatin</b>
<b>Survival (months)</b>	<b>(n= 46)</b>	<b>(n = 44)</b>
Median (95% CI)	8.8 (6.4, 11.5)	15.7 (11.9, 17.7)
Hazard ratio (95% CI)	0.51 (0.31, 0.82)	
<b>Patients with Prior Cisplatin Chemoradiotherapy</b>		
	<b>Cisplatin</b>	<b>Topotecan/Cisplatin</b>
Survival (months)	(n= 72)	(n = 69)
Median (95% CI)	5.9 (4.7, 8.8)	7.9 (5.5, 10.9)
Hazard ratio (95% CI)	0.85 (0.59, 1.21)	

In patients (n=39) with recurrence within 180 days after chemoradiotherapy with cisplatin, the median survival in the topotecan plus cisplatin arm was 4.6 months (95% CI: 2.6, 6.1) versus 4.5 months (95% CI: 2.9, 9.6) for the cisplatin arm with a hazard ratio of 1.15 (0.59, 2.23). In those patients (n=102) with recurrence after 180 days, the median survival in the topotecan plus cisplatin arm was 9.9 months (95% C.I.: 7, 12.6) versus 6.3 months (95% C.I.: 4.9, 9.5) for the cisplatin arm with a hazard ratio of 0.75 (0.49, 1.16).

#### *Paediatric population*

Topotecan was also evaluated in the paediatric population; however, only limited data on efficacy and safety are available.

In an open-label study involving children (n = 108, age range: infant to 16 years) with recurrent or progressive solid tumours, topotecan was administered at a starting dose of 2.0 mg/m<sup>2</sup> given as a 30-minute infusion for 5 days repeated every 3 weeks for up to one year depending on response to therapy. Tumour types included were Ewing's Sarcoma/primitive neuroectodermal tumour, neuroblastoma, osteoblastoma, and rhabdomyosarcoma. Antitumour activity was demonstrated primarily in patients with neuroblastoma. Toxicities of topotecan in paediatric patients with recurrent and refractory solid tumours were similar to those historically seen in adult patients. In this study, forty-six (43%) patients received G-CSF over 192 (42.1%) courses; sixty-five (60%) received transfusions of Packed Red Blood Cells and fifty (46%) of platelets over 139 and 159 courses (30.5% and 34.9%), respectively. Based on the dose-limiting toxicity of myelosuppression, the maximum tolerated dose

(MTD) was established at 2.0 mg/m<sup>2</sup>/day with G-CSF and 1.4 mg/m<sup>2</sup>/day without G-CSF in a pharmacokinetic study in paediatric patients with refractory solid tumours (see section 5.2).

## 5.2 Pharmacokinetic properties

### Distribution:

Following intravenous administration of topotecan at doses of 0.5 to 1.5 mg/m<sup>2</sup> as a 30 minute infusion daily for five days, topotecan demonstrated a high plasma clearance of 62 l/h (SD 22), corresponding to approximately 2/3 of liver blood flow. Topotecan also had a high volume of distribution, about 132 l (SD 57), and a relatively short half-life of 2-3 hours. Comparison of pharmacokinetic parameters did not suggest any change in pharmacokinetics over the 5 days of dosing. Area under the curve increased approximately in proportion to the increase in dose. There is little or no accumulation of topotecan with repeated daily dosing and there is no evidence of a change in the pharmacokinetics after multiple doses. Preclinical studies indicate plasma protein binding of topotecan is low (35%) and distribution between blood cells and plasma was fairly homogeneous.

### Biotransformation

The elimination of topotecan has only been partly investigated in man. A major route of clearance of topotecan was by hydrolysis of the lactone ring to form the ring-opened carboxylate.

Metabolism accounts for <10% of the elimination of topotecan. An N-desmethyl metabolite, which was shown to have similar or less activity than the parent in a cell-based assay, was found in urine, plasma, and faeces. The mean metabolite:parent AUC ratio was < 10% for both total topotecan and topotecan lactone. An O-glucuronidation metabolite of topotecan and N-desmethyl topotecan has been identified in the urine.

### Elimination

Overall recovery of topotecan-related material following five daily doses of topotecan was 71 to 76 % of the administered IV dose. Approximately 51% was excreted as total topotecan and 3 % was excreted as N-desmethyl topotecan in the urine. Faecal elimination of total topotecan accounted for 18 % while faecal elimination of N-desmethyl topotecan was 1.7%. Overall, the N-desmethyl metabolite contributed a mean of less than 7 % (range 4-9 %) of the total topotecan related material accounted for in the urine and faeces. The topotecan-O-glucuronide and N-desmethyl topotecan-O-glucuronide in the urine were less than 2.0 %.

*In vitro* data using human liver microsomes indicate the formation of small amounts of N-demethylated topotecan. *In vitro*, topotecan did not inhibit human P450 enzymes CYP1A2, CYP2A6, CYP2C8/9, CYP2C19, CYP2D6, CYP2E, CYP3A or CYP4A nor did it inhibit the human cytosolic enzymes dihydropyrimidine or xanthine oxidase.

When given in combination with cisplatin (cisplatin day 1, topotecan days 1 to 5), the clearance of topotecan was reduced on day 5 compared to day 1 (19.1 L/h/m<sup>2</sup> compared to 21.3 L/h/m<sup>2</sup> [n=9]) (see section 4.5).

### Special populations

#### Hepatic impairment

Plasma clearance in patients with hepatic impairment (serum bilirubin between 1.5 and 10 mg/dl) decreased to about 67 % when compared with a control group of patients. Topotecan half-life was increased by about 30 % but no clear change in volume of distribution was observed. Plasma clearance of total topotecan (active and inactive form) in patients with hepatic impairment only decreased by about 10 % compared with the control group of patients.

#### Renal impairment

Plasma clearance in patients with renal impairment (creatinine clearance 41-60 ml/min.) decreased to about 67 % compared with control patients. Volume of distribution was slightly decreased and thus half-life only increased by 14 %. In patients with moderate renal impairment topotecan plasma clearance was reduced to 34 % of the value in control patients. Mean half-life increased from 1.9 hours to 4.9 hours.

#### Age/weight

In a population study, a number of factors including age, weight and ascites had no significant effect on clearance of total topotecan (active and inactive form).

#### Paediatric population

The pharmacokinetics of topotecan given as a 30-minute infusion for 5 days were evaluated in two studies. One study included a dose range of 1.4 mg/m<sup>2</sup> to 2.4 mg/m<sup>2</sup> in children (aged 2 up to 12 years, n = 18), adolescents (aged 12 up to 16 years, n = 9), and young adults (aged 16 to 21 years, n = 9) with refractory solid tumours. The second study included a dose range of 2.0 mg/m<sup>2</sup> to 5.2 mg/m<sup>2</sup> in children (n = 8), adolescents (n = 3), and young adults (n = 3) with leukaemia. In these studies, there were no apparent differences in the pharmacokinetics of topotecan among children, adolescents, and young adult patients with solid tumours or leukaemia, but data are too limited to draw definite conclusions.

### 5.3 Preclinical safety data

Resulting from its mechanism of action, topotecan is genotoxic to mammalian cells (mouse lymphoma cells and human lymphocytes) in vitro and mouse bone marrow cells in vivo. Topotecan was also shown to cause embryo-foetal lethality when given to rats and rabbits.

In reproductive toxicity studies with topotecan in rats there was no effect on male or female fertility; however, in females super-ovulation and slightly increased pre-implantation loss were observed

The carcinogenic potential of topotecan has not been studied.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Tartaric acid (E334)  
Hydrochloric acid (for pH adjustment) (E507)  
Sodium hydroxide (for pH adjustment)  
Water for injections

### 6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### 6.3 Shelf life

*Unopened Vials*  
36 months.

*Diluted solution*

Chemical and physical in-use stability has been demonstrated for 30 days at 25°C under normal light conditions and at 2-8°C when protected from light. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer 24 hours at 2 to 8°C, unless dilution has taken place in controlled and validated aseptic conditions.

### 6.4 Special precautions for storage

Store below 25°C. Keep the vial in the outer carton in order to protect from light.  
For storage conditions after dilution of the medicinal product, see section 6.3.

### 6.5 Nature and contents of container

Topotecan 1 mg/ml, 1 ml is supplied in 2 ml Type I amber glass vial and are closed with 13 mm flurotec rubber stopper and sealed with 13 mm aluminium flip off royal blue seal.

Topotecan 1 mg/ml, 4 ml is supplied in 5 ml Type I amber glass vial and are closed with 13 mm flurotec rubber stopper and sealed with 13 mm aluminium flip off royal blue seal.

Topotecan Infusion is available in cartons containing 1 vial and 5 vials.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal

Topotecan is provided as a sterile concentrate containing 1 mg Topotecan in 1 ml solution and 4 mg Topotecan in 4 ml solution.

Parenteral products should be visually inspected for particulate matter and discoloration prior to administration. The medicinal product is a clear yellow colour solution. If visible particles are observed, the product should not be administered. Further dilution with either sodium chloride 9 mg/ml (0.9%) solution for injection or glucose 50 mg/ml (5%) solution for injection is required, to obtain a final concentration of between 25 and 50 micrograms/ml prior to administration to the patient.

The normal procedures for proper handling and disposal of anticancer medicinal products should be adopted, namely:

- Personnel should be trained to reconstitute the medicinal product.
- Pregnant staff should be excluded from working with this medicinal product.
- Personnel handling this medicinal product during reconstitution should wear protective clothing including mask, goggles and gloves.
- All items for administration or cleaning, including gloves, should be placed in high-risk, waste disposal bags for high-temperature incineration.
- Accidental contact with the skin or eyes should be treated immediately with copious amounts of water. If there is lasting irritation, a doctor should be consulted.
- Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## 7 MARKETING AUTHORISATION HOLDER

Accord Healthcare Ireland Ltd.  
Euro House  
Euro Business Park  
Little Island  
Cork T45 K857  
Ireland

## 8 MARKETING AUTHORISATION NUMBER

PA2315/121/001

## 9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 4th May 2012

Date of last renewal: 27th February 2017

## 10 DATE OF REVISION OF THE TEXT

September 2025