

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Glucosamine sulfate 750mg Film-coated Tablets

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 942 mg glucosamine sulfate sodium chloride equivalent to 750 mg glucosamine sulfate or 589 mg glucosamine.

### Excipient(s) with known effect

Each tablet contains 75.9 mg (3.3mmol) of sodium, 3.0 mg Lactose monohydrate and 0.36 mg Lecithin soya (E322)

For a full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Film-coated tablet.

Off-white oblong shaped film-coated tablet, 8x19mm.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic Indications

Glucosamine sulfate tablets are indicated for relief of symptoms in mild to moderate osteoarthritis of the knee.

### 4.2 Posology and method of administration

Method of administration:

Glucosamine sulfate tablets should be swallowed whole.

Tablets can be taken with or without food.

#### *Adults and the elderly:*

One Glucosamine sulfate tablet should be taken twice daily.

Or

Two Glucosamine sulfate tablets to be taken once daily.

Glucosamine is not indicated for the treatment of acute painful symptoms. Relief of symptoms (especially pain relief) may not be experienced until after several weeks of treatment and in some cases even longer. If no relief of symptoms is experienced after 2-3 months, continued treatment with glucosamine should be reevaluated.

Additional information on special populations:

#### *Paediatric population*

Safety and efficacy has not been established in children and adolescents, therefore, Glucosamine sulfate tablets should not be used in persons under the age of 18 years.

#### *Elderly*

No specific studies have been performed in the elderly, but according to clinical experience dosage adjustment is not required when treating otherwise healthy, elderly patients.

#### *Impaired renal and/or liver function*

In patients with impaired renal and/or liver function no dose recommendations can be given, since no studies have been performed.

### 4.3 Contraindications

Hypersensitivity to the active substance, glucosamine (or any of its derivatives), sulfates or to any of the excipients listed in section 6.1.

Glucosamine sulfate tablets must not be used in patients who are allergic to shellfish as the active ingredient is obtained from shellfish.

Glucosamine sulfate tablets contains soya lecithin. Persons allergic to soya or peanut should therefore not use this medicinal product.

### 4.4 Special warnings and precautions for use

The presence of other joint disease, which would require alternative treatment, should be excluded.

In patients with impaired glucose tolerance, monitoring of the blood glucose levels and, where relevant, insulin requirements is recommended before start of treatment and periodically during treatment.

In patients with a known risk factor for cardiovascular disease, monitoring of the blood lipid levels is recommended since hypercholesterolemia has been observed in a few patients treated with glucosamine.

A report on exacerbated asthma symptoms triggered after initiation of glucosamine therapy has been described (symptoms resolved after withdrawal of glucosamine). Asthmatic patients starting on glucosamine should therefore be aware of potential worsening of asthma symptoms.

Glucosamine sulfate tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

This medicinal product contains 75.9 mg sodium per film-coated tablet, equivalent to 3.8 % of the WHO recommended maximum daily intake of 2 g sodium for an adult.

### 4.5 Interaction with other medicinal products and other forms of interactions

There are limited data on possible drug interactions with glucosamine, but increments in the INR parameter have been reported with oral vitamin K antagonists. Patients treated with oral vitamin K antagonists should therefore be closely monitored at the time of initiation or termination of glucosamine therapy.

Close monitoring of blood sugar levels is recommended for diabetics on hypoglycaemic agents.

Concurrent treatment with glucosamine may increase the absorption and serum concentrations of tetracyclines, but the clinical relevance of this interaction is probably limited.

Due to limited documentation on potential drug interactions with glucosamine, one should generally be aware of altered response or concentration of concurrently used medical products.

### 4.6 Fertility, pregnancy and lactation

#### Pregnancy

There are inadequate data concerning the use of glucosamine in pregnant women. From animal studies only insufficient data are available. Glucosamine should not be used during pregnancy.

**Breast feeding**

There is no data available on the excretion of glucosamine in breastmilk. The use of glucosamine during breast feeding is therefore not recommended as there is no data on the safety of the child.

**4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive or use machines have been performed. If dizziness or drowsiness is experienced, car driving and the operating of machinery is not recommended.

**4.8 Undesirable effects**

**The most common adverse reactions associated with treatment with glucosamine are nausea, abdominal pain, indigestion, constipation and diarrhoea. In addition, headache, tiredness, rash itching, and flushing have been reported. The reported adverse reactions are usually mild and transitory.**

<b>Medra System Organ Class</b>	<b>Common (<math>\geq 1/100</math> to <math>&lt; 1/10</math>)</b>	<b>Uncommon (<math>\geq 1/1,000</math> to <math>&lt; 1/100</math>)</b>	<b>Rare (<math>\geq 1/10,000</math> to <math>&lt; 1/1,000</math>)</b>	<b>Not known (cannot be estimated from the available data)</b>
<b>Nervous system disorders</b>	Headache Tiredness			Dizziness
<b>Respiratory, thoracic and mediastinal disorders</b>				Asthma / Asthma aggravated
<b>Gastrointestinal disorders</b>	Nausea Abdominal pain Indigestion Diarrhoea Constipation			Vomiting
<b>Skin and subcutaneous tissue disorders</b>		Rash Itching Flushing		Angiodema Urticaria
<b>Metabolism and nutrition disorders</b>				Diabetes mellitus inadequate control Hypercholesterolaemia
<b>General disorders and administration site conditions</b>				Oedema/peripheral oedema

Cases of Hypercholesterolemia, Asthma, aggravated and Diabetes mellitus inadequate control have been reported, but causality has not been established.

Glucosamine sulfate tablets may cause Hepatic enzyme elevation and rarely jaundice.

**Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance. Website: [www.hpra.ie](http://www.hpra.ie)

**4.9 Overdose**

Signs and symptoms of accidental or intentional overdose with glucosamine might include headache, dizziness, disorientation, arthralgia, nausea, vomiting, diarrhoea or constipation.

In cases of overdose, treatment with glucosamine should be discontinued and standard supportive measures should be adopted as required.

In clinical trials one of five healthy young subjects experienced headache following infusion of glucosamine up to 30 g. In addition, one case of overdose has been reported in a 12-year old female who took orally 28 g of glucosamine hydrochloride. She developed arthralgia, vomiting and disorientation. The patient fully recovered.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Other anti-inflammatory and anti-rheumatic agents, non-steroidal anti-inflammatory drugs. ATC code: M01AX05

Glucosamine is an endogenous substance, a normal constituent of the polysaccharide chains of cartilage matrix and synovial fluid glucosaminoglycans. In vitro and in vivo studies have shown glucosamine stimulates the synthesis of physiological glycosaminoglycans and proteoglycans by chondrocytes and of hyaluronic acid by synoviocytes. The mechanism of action of glucosamine in humans is unknown. The period to onset of response cannot be assessed.

### **5.2 Pharmacokinetic properties**

Glucosamine is a relatively small molecule (molecular mass 179), which is easily dissolved in water and soluble in hydrophilic organic solvents. The available information on the pharmacokinetics of glucosamine is limited. The absolute bioavailability is unknown. The distribution volume is approximately 5 litres and the half-life after intravenous administration is approximately 2 hours. Approximately 38% of an intravenous dose is excreted in the urine as unchanged substance.

### **5.3 Preclinical safety data**

D-glucosamine has low acute toxicity. Animal experimental data relating to toxicity during repeated administration, reproduction toxicity, mutagenicity and carcinogenicity is lacking for glucosamine.

Results from in vitro studies and in vivo studies in animals have shown that glucosamine reduces insulin secretion and induces insulin resistance, probably via glucokinase inhibition in the beta cells. The clinical relevance is unknown.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Tablet:

Microcrystalline cellulose 101  
Microcrystalline cellulose 102  
Lactose monohydrate  
Pregelatinised maize starch  
Crospovidone  
Stearic acid

Coating:

Poly(vinyl) alcohol  
Titanium dioxide (E171)  
Talc (E553b)  
Lecithin soya (E322)  
Macrogol 3350

## 6.2 Incompatibilities

Not applicable.

## 6.3 Shelf life

3 years.

After first opening of the tablet container the medicinal product should be used within 6 months.

## 6.4 Special precautions for storage

This medicinal product does not require any special temperature storage conditions. Store in the original package in order to protect from moisture.

## 6.5 Nature and contents of container

Cartons containing PVdC coated PVC/Al blisters: Pack Size: 8, 10, 12, 14, 20, 28, 30, 56, 60, 112, 120, 168, 180, 336, 360 film-coated tablets.

OR

Cartons containing HDPE containers fitted with a tamper-evident HDPE screw cap.

Pack Size: 8, 10, 12, 14, 20, 28, 30, 56, 60, 112, 120, 168, 180, 336, 360 film-coated tablets.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal

No special requirements.

## 7 MARKETING AUTHORISATION HOLDER

FMC Pharma Ltd  
Unit 622, Northern Extension  
Co. Waterford  
Ireland

## 8 MARKETING AUTHORISATION NUMBER

PA1521/003/002

## 9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 26th August 2011

**10 DATE OF REVISION OF THE TEXT**

February 2021