

PACKAGE LEAFLET: INFORMATION FOR THE USER

Temgesic Injection 1 ml

Buprenorphine hydrochloride

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any of the side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Temgesic is and what it is used for
2. What you need to know before you use Temgesic Injection
3. How to use Temgesic Injection
4. Possible side effects
5. How to store Temgesic Injection
6. Contents of the pack and other information

1. WHAT TEMGESIC INJECTION IS AND WHAT IT IS USED FOR

Temgesic Injection 1 ml is a strong pain killer that helps lessen moderate or severe pain.

2. BEFORE YOU USE TEMGESIC INJECTION

Do not use Temgesic

- If you are allergic (hypersensitive) to buprenorphine (the active ingredient in Temgesic)-or to other opiates-or to any other component of the product.

Warnings and precautions

Talk to your doctor or pharmacist before using Temgesic Injection if you have:

- asthma or other breathing problems or are taking medicines which may make your breathing slower or weaker,
- an addiction or have ever been addicted to medicines or substances,
- problems with your liver, such as hepatitis,
- any kidney disease,
- low blood pressure,
- recently suffered from head injury or brain disease,
- thyroid problems,
- adrenocortical disorder such as Addison's disease,
- a urinary disorder such as an enlarged prostate or narrowing of the urethra.
- prostatic disease,

- are elderly
- a history of gallbladder disease
- acute abdominal pain
- alcohol problems
- curvature of the spine
- depression or other conditions that are treated with antidepressants. The use of these medicines together with Temgesic can lead to serotonin syndrome, a potentially life-threatening condition (see "Other medicines and Temgesic").

Important things to be aware of:

- **Misuse, abuse and diversion**

This medicine can be a target for people who abuse prescription medicines, and should be kept in a safe place to protect it from theft. **Do not give this medicine to anyone else.** It can cause death or otherwise harm them.

- **Breathing problems**

Some people have died from respiratory failure (inability to breathe) because they misused this medicine or took it in combination with other central nervous system depressants, such as alcohol, benzodiazepines (tranquilisers), or other opioids.

- **Sleep-related breathing disorders**

Temgesic can cause sleep-related breathing disorders such as sleep apnoea (breathing pauses during sleep) and sleep related hypoxemia (low oxygen level in the blood). The symptoms can include breathing pauses during sleep, night awakening due to shortness of breath, difficulties to maintain sleep or excessive drowsiness during the day. If you or another person observe these symptoms, contact your doctor. A dose reduction may be considered by your doctor.

- **Tolerance, dependence, and addiction**

This medicine contains buprenorphine which is an opioid medicine. Repeated use of opioids can result in the drug being less effective (you become accustomed to it, known as tolerance).

Repeated use of Temgesic can also lead to dependence, abuse, and addiction, which may result in life-threatening overdose. The risk of these side effects can increase with a higher dose and longer duration of use.

Dependence or addiction can make you feel that you are no longer in control of how much medicine you need to take or how often you need to take it.

The risk of becoming dependent or addicted varies from person to person. You may have a greater risk of becoming dependent on or addicted to Temgesic if:

- You or anyone in your family have ever abused or been dependent on alcohol, Prescription medicines or illegal drugs ("addiction").
- You are a smoker.
- You have ever had problems with your mood (depression, anxiety, or a personality disorder) or have been treated by a psychiatrist for other mental illnesses.

If you notice any of the following signs whilst taking Temgesic, it could be a sign that you have become dependent or addicted:

- You need to take the medicine for longer than advised by your doctor
- You need to take more than the recommended dose
- You might feel that you need to carry on taking your medicine, even when it doesn't help to relieve your pain.

- You are using the medicine for reasons other than prescribed, for instance, ‘to stay calm’ or ‘help you sleep’
- You have made repeated, unsuccessful attempts to quit or control the use of the medicine
- When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again (‘withdrawal effects’)

If you notice any of these signs, speak to your doctor to discuss the best treatment Pathway for you, including when it is appropriate to stop and how to stop safely.

- **Withdrawal symptoms**

This product can cause withdrawal symptoms if you are also taking other opioids such as heroin or methadone. Temgesic can also cause withdrawal symptoms if you stop taking it abruptly.

- **Diagnosis of unrelated medical conditions**

This medicine may mask pain symptoms that could assist in the diagnosis of some diseases. Do not forget to advise your doctor if you take this medicine.

Other medicines and Temgesic Injection

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Some medicines may increase the side effects of Temgesic and may cause various serious reactions. Do not take any medicines whilst taking Temgesic without talking to your doctor, especially:

- Benzodiazepines (used to treat anxiety or sleep disorders) such as diazepam, temazepam, or alprazolam. Your doctor will prescribe the correct dose for you. Concomitant use of Temgesic and sedative medicines such as benzodiazepines or related medicines increases the risk of drowsiness, difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible. However if your doctor does prescribe Temgesic together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor. Please tell your doctor about all sedative medicines you are taking, and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.
- **Other medicines that may make you feel sleepy which are** used to treat illnesses such as anxiety, sleeplessness, convulsions/seizures, or pain. These types of medicines will reduce your alertness levels making it difficult for you to drive and use machines. They may also cause a decreased heart rate and ability to breathe and may cause unconsciousness, which is very serious. Below is a list of examples of these types of medicines:
 - Other opioid containing medicines such as methadone, certain pain killers and cough suppressants.
 - Anti-depressants (used to treat depression) such as isocarboxazide, phenelzine, selegeline, tranylcypromine and valproate may increase the effects of this medicine.

- Sedative H₁ receptor antagonists (used to treat allergic reactions) such as diphenhydramine and chlorphenamine.
- Antihistamines or antiemetics (used to treat allergies, travel sickness or nausea)
- Barbiturates (used to cause sleep or sedation) such as phenobarbital and secobarbital.
- Tranquilisers (used to cause sleep or sedation) such as chloral hydrate.
- Anti-depressants, such as moclobemide, tranylcypromine, citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline, duloxetine, venlafaxine, amitriptyline, doxepine, or trimipramine. These medicines may interact with Temgesic and you may experience symptoms such as involuntary, rhythmic contractions of muscles, including the muscles that control movement of the eye, agitation, hallucinations, coma, excessive sweating, tremor, exaggeration of reflexes, increased muscle tension, body temperature above 38°C. Contact your doctor when experiencing such symptoms.
- Antipsychotics or neuroleptics (used to treat psychiatric disorders)
- Muscle relaxants
- Medicines to treat Parkinson's disease
- Clonidine (used to treat high blood pressure) may extend the effects of this medicine.
- Naltrexone may prevent the therapeutic effects of Temgesic. If currently taking this medicine followed by concomitant use of naltrexone, you may experience a sudden onset of prolonged and intense withdrawal.
- Anti-retrovirals (used to treat AIDS) such as ritonavir, nelfinavir and indinavir may increase the effects of this medicine.
- Some antifungal agents (used to treat fungal infections) such as ketoconazole, itraconazole and certain antibiotics (macrolides) may extend the effects of this medicine.
- Some medicines may decrease the effect of Temgesic. These include medicines used to treat epilepsy or pain due to nerve problems/neuropathic pain (such as gabapentin, pregabalin, phenobarbital, Phenothiazines carbamazepine and phenytoin), and medicines used to treat tuberculosis (rifampicin).
- Anaesthetics (used during surgery such as halothane).

Athletes must be aware that this medicine may cause a positive reaction to 'anti-doping' tests.

Temgesic Injection with food, drink and alcohol

Do not take Temgesic together with alcohol. Alcohol may increase drowsiness and may increase the risk of respiratory failure if taken with Temgesic. Do not take Temgesic together with alcohol.

Pregnancy and breast-feeding

The risks of using Temgesic in pregnant women are not known. Tell your doctor if you are pregnant or intend to become pregnant. Your doctor will decide if your treatment should be continued with an alternative medicine.

When taken during pregnancy, particularly late pregnancy, medicines like Temgesic may cause drug withdrawal symptoms including problems with breathing in your newborn baby. This may not appear until several days after birth.

Do not breast-feed whilst taking this medicine, since Temgesic passes into breast milk.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Temgesic may cause drowsiness, particularly if you also drink alcohol or take sedative medicines. Do not drive, use any tools or machines, or perform dangerous activities until you know how this medicine affects you.

3. HOW TO USE TEMGESIC INJECTION

Always take Temgesic exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. **Do not inject more than your doctor has told you to.**

Before starting treatment and regularly during treatment, your doctor will discuss with you what you may expect from using Temgesic, when and how long you need to take it, when to contact your doctor, and when you need to stop.

Adults and children over 12 years of age

The usual dose is 300 – 600 micrograms of buprenorphine every six to eight hours, or as required. Inject Temgesic into a muscle or slowly into a vein, as instructed by your doctor.

Children between 6 years and 12 years of age

The dose depends on their weight and your doctor will tell you the correct dose for your child.

This product will not be given to infants and children under 6 months of age.

If you take more Temgesic Injection than you should

If you or someone else take too much of this medicine, you must go or be taken immediately to an emergency centre or hospital for treatment as overdose with Temgesic may cause serious and life-threatening breathing problems.

Symptoms of overdose may include feeling sleepy, low blood pressure (hypotension), nausea (feeling sick), vomiting, constricted pupils (small pupils characterized by blurred vision and difficulty focusing) and you may breathe much slower than is normal for you.

If you forget to take Temgesic Injection

If you miss an injection, take it as soon as you remember unless it is time for your next injection. Do not take a double dose to make up for a missed dose.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately or you must go or be taken immediately to an emergency centre or hospital if you experience any of the following symptoms after taking this medicine. These symptoms can be serious and Temgesic may cause serious and life-threatening breathing problems:

- If you start to breathe more slowly or weakly than usual. This may be a sign of respiratory depression induced by Temgesic (affects an unknown number of people).
- If you get any sudden wheezing, difficulty breathing, swelling of the eyelids, face or lips, rash or itching especially those covering your whole body. These may be signs of an allergic reaction (rare side effect that may affect up to 1 in 1,000 people).
- If you start to feel faint, as this may be a sign of low blood pressure (common side effect that may affect up to 1 in 10 people).
- If you get sudden constriction of the airways (bronchial spasm) (affects an unknown number of people).
- If you get a sensitivity reaction with fever, skin rash, swelling and sometimes decreased blood pressure (anaphylactic shock) (affects an unknown number of people).
- If you get a swollen face, tongue, or throat, or difficulties to swallow, hives and difficulties to breathe (angioedema) (affects an unknown number of people).

Other side effects that may occur with Temgesic are presented below:

<i>Very common side effects (may affect more than 1 in 10 people):</i>
Sedation, dizziness, nausea (feeling sick), loss of balance and/or giddiness (vertigo).
<i>Common side effects (may affect up to 1 in 10 people):</i>
Headache, miosis (small pupils), hypoventilation (shallow or slow breathing), vomiting, hyperhidrosis (increased sweating), low blood pressure.
<i>Uncommon side effects (may affect up to 1 in 100 people):</i>
State of confusion, euphoric mood (exaggerated feeling of well-being), nervousness, depression, psychotic disorder (mental disorder causing abnormal thinking and perceptions), hallucination, depersonalisation (not feeling like yourself), dysarthria (speech disorder), paraesthesia (pins and needles), coma, tremor, blurred vision, diplopia (double vision), visual impairment, conjunctivitis (infection of the eye), tinnitus (ringing of the ears), tachycardia (fast heartbeat) bradycardia (slow heartbeat), cyanosis (blue or purple discoloration of the skin), second degree atrioventricular block (heart condition), hypertension (high blood pressure), pallor (pale skin), dyspnoea (shortness of breath), apnoea (pauses in breathing), dry mouth, constipation, dyspepsia (indigestion), flatulence, pruritus (itching), rash, urinary retention, asthenia (feeling weak), fatigue, malaise (feeling of discomfort).

<i>Rare side effects (may effect up to 1 in 1000 people):</i>
Decreased appetite, dysphoria (feeling unwell or unhappy), agitation, convulsion (fits), abnormal coordination, diarrhoea, urticaria (hives), allergy (hypersensitivity).
<i>Not known (frequency can not be estimated from the available data):</i>
Drug dependence, somnolence, drug ineffective, drug interaction, dental caries.

Experience of the use of Temgesic in patients have also identified the following reactions and/or side effects:

Serious allergic reaction, confusional state, drug dependence, hallucination, sleepiness, dizziness, headache, low blood pressure, very slow breathing rate (respiratory depression), shortness of breath (Bronchospasm), nausea, vomiting, itching, rash, excessive sweating, swelling under the skin, medicine not working, medicine affected by another medicine, tiredness.

Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

HPRA Pharmacovigilance
 Earlsfort Terrace
 IRL - Dublin 2
 Tel: +353 1 6764971
 Fax: +353 1 6762517
 Website: www.hpra.ie
 e-mail: <mailto:medsafety@hpra.ie>

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. HOW TO STORE TEMGESIC INJECTION

Keep this medicine out of the sight and reach of children.
 Store this medicine in a safe and secure place, where other people cannot access it. It can cause serious harm and be fatal to people who may take this medicine by accident, or intentionally when it has not been prescribed for them.

Store below 30°C

Store in the original package to protect from light.

Do not use Temgesic Injection after the expiry date which is stated on the pack (EXP month/year). The expiry date refers to the first day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Temgesic Injection contains

- The active substance is 300 micrograms buprenorphine hydrochloride
- The other ingredients are dextrose, hydrochloric acid and water for injections.

What Temgesic Injection looks like and contents of the pack

Temgesic Injection comes in 1 ml ampoules and in cartons containing a plastic tray of five ampoules.

Marketing Authorisation Holder

EUMEDICA Pharmaceuticals GmbH
Basler Straße 126
79540 Lörrach
Germany

Manufacturer

Indivior Europe Limited
27 Windsor Place
Dublin 2
D02 DK44
Ireland

This leaflet was last revised in August 2024.