

Package leaflet: Information for the user

***OxyContin*[®] 15 mg, 30 mg, 60 mg, 120 mg and 160 mg prolonged release tablets**
Oxycodone hydrochloride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What ***OxyContin*** is and what it is used for
2. What you need to know before you take ***OxyContin***
3. How to take ***OxyContin***
4. Possible side effects
5. How to store ***OxyContin***
6. Contents of the pack and other information

1. What *OxyContin* is and what it is used for

These tablets have been prescribed for you by your doctor to relieve severe pain over a period of 12 hours. They contain the active ingredient oxycodone which is a strong analgesic ('painkiller') which belongs to a group of medicines called opioids.

2. What you need to know before you take *OxyContin*

Do not take *OxyContin* if you:

- are allergic (hypersensitive) to oxycodone or any of the ingredients of the tablets (listed in section 6) or have previously had an allergic reaction when taking other strong analgesics or painkillers (such as morphine or other opioids);
- have breathing problems, such as severe chronic obstructive lung disease, severe bronchial asthma or severe respiratory depression. Symptoms may include breathlessness, coughing or breathing more slowly and weakly than expected;
- have a head injury that causes a severe headache or makes you feel sick. This is because the tablets may make these symptoms worse or hide the extent of the head injury;
- have a condition where the bowel does not work properly (paralytic ileus), your stomach empties more slowly than it should (delayed gastric emptying) or you have severe sudden pain in your abdomen (acute abdomen);
- have a heart problem after long-term lung disease (cor pulmonale);

Warnings and precautions

Talk to your doctor or pharmacist before taking these tablets if you:

- are elderly or weakened;
- have an under-active thyroid gland (hypothyroidism);
- have myxoedema (a thyroid disorder, with dryness, coldness, and swelling ['puffiness'] of the skin, affecting the face and limbs)
- have a severe headache or feel sick as this may indicate that the pressure in your skull is increased;

- have low blood pressure (hypotension);
- have a mental disorder as a result of an intoxication (toxic psychosis);
- have inflammation of the pancreas (which may cause severe pain in the abdomen and back) or problems with your gall bladder or bile duct;
- have a blockage of the gut or an inflammatory bowel disorder;
- have colicky abdominal pain or discomfort;
- have an enlarged prostate gland, which causes difficulty in passing urine (in men);
- have poor adrenal gland function (your adrenal gland is not working properly) for example Addison's disease;
- have breathing problems such as severely impaired pulmonary function, chronic obstructive airways disease, severe lung disease or reduced respiratory reserve. Symptoms may include breathlessness and coughing;
- have kidney or liver problems;
- are or have ever been addicted to alcohol or drugs, or have a known opioid dependence;
- have withdrawal symptoms such as agitation, anxiety, palpitations, shaking or sweating upon stopping taking alcohol or drugs;
- suffer from seizures, fits or convulsions;
- are feeling light-headed or faint.
- need to take increasingly higher doses of ***OxyContin*** to gain the same level of pain relief (tolerance);
- have an increase in sensitivity to pain;
- are taking a type of medicine known as a monoamine oxidase inhibitor (examples include tranylcypromine, phenelzine, isocarboxazid, moclobemide and linezolid), or you have taken this type of medicine in the last two weeks.

If you are going to have an operation, please tell the doctor at the hospital that you are taking these tablets.

The tablets should never be crushed, or injected as this may lead to serious side effects, which may be fatal.

You may experience hormonal changes while taking these tablets. Your doctor may want to monitor these changes.

Other medicines and *OxyContin*

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. If you take these tablets with some other medicines, the effect of these tablets or the other medicine may be changed.

Tell your doctor or pharmacist if you are taking:

- a type of medicine known as a monoamine oxidase inhibitor or you have taken this type of medicine in the last two weeks (see 'Warnings and precautions');
- medicines to help you sleep or stay calm (for example hypnotics or sedatives, including benzodiazepines);
- medicines to treat depression (for example paroxetine and fluoxetine);
- medicines to treat psychiatric or mental disorders (such as phenothiazines or neuroleptic drugs);
- other strong analgesics ('painkillers');
- muscle relaxants;
- medicines to treat high blood pressure;
- quinidine (a medicine to treat a fast heartbeat);
- cimetidine (a medicine for stomach ulcers, indigestion or heartburn);

- medicines to treat fungal infections (such as ketoconazole, voriconazole, itraconazole, or posaconazole);
- medicines to treat infections (such as clarithromycin, erythromycin or telithromycin).
- a specific type of medicine known as a protease inhibitor to treat HIV (examples include boceprevir, ritonavir, indinavir, nelfinavir or saquinavir);
- rifampicin to treat tuberculosis;
- carbamazepine (a medicine to treat seizures, fits or convulsions and certain pain conditions);
- phenytoin (a medicine to treat seizures, fits or convulsions);
- a herbal remedy called St John's Wort (also known as *Hypericum perforatum*);
- antihistamines;
- medicines to treat Parkinson's disease.

Also, tell your doctor if you have recently been given an anaesthetic.

Taking *OxyContin* with food, drink and alcohol

These tablets may be taken with or without food.

Drinking alcohol whilst taking *OxyContin* tablets may make you feel more sleepy or increase the risk of serious side effects such as shallow breathing with a risk of stopping breathing, and loss of consciousness. It is recommended not to drink alcohol while you are taking *OxyContin* tablets. You should avoid drinking grapefruit juice during your treatment with *OxyContin*.

Pregnancy and breastfeeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking any medicine.

Pregnancy

You should not use these tablets during pregnancy and labour unless you have been specifically told by your doctor. Depending on the dose and duration of therapy with oxycodone, slow and shallow breathing (respiratory depression) or withdrawal symptoms may occur in the newborn infant.

Breastfeeding

These tablets should not be used while breastfeeding because the active ingredient can pass into breast milk.

Driving and using machines

These tablets may cause a number of side effects such as drowsiness which could affect your ability to drive or use machinery (see section 4 for a full list of side effects). These are usually most noticeable when you first start taking the tablets, or when changing to a higher dose. If you are affected you should not drive or use machinery.

***OxyContin* contains lactose**

These tablets contain lactose which is a form of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking these tablets.

3. How to take *OxyContin*

Always take these tablets exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. The label on your medicine will tell you how many tablets to take and how often.

Do not exceed the dose recommended by your doctor.

Swallow your tablets whole with a glass of water. **Do not chew, crush or dissolve them.**

You should take your tablets every 12 hours. For instance, if you take a tablet at 8 o'clock in the morning, you should take your next tablet at 8 o'clock in the evening.

The tablets should never be crushed or injected as this may lead to serious side effects, which may be fatal. You must only take these tablets by mouth.

***OxyContin* tablets are designed to work properly over 12 hours when swallowed whole. If a tablet is broken, crushed, dissolved or chewed, the entire 12-hour dose may be absorbed rapidly into your body. This can be dangerous, causing serious problems such as an overdose, which may be fatal.**

Adults (20 years of age and over)

The usual starting dose is one 10 mg tablet every 12 hours. However, your doctor will prescribe the dose required to treat your pain. If you find that you are still in pain whilst taking these tablets discuss this with your doctor.

Children and adults under 20 years of age

Children and adults under 20 years of age should not take the tablets.

Patients with kidney or liver problems

Please tell your doctor if you suffer from kidney or liver problems as they may prescribe you an alternative medicine or a lower dose depending upon your condition.

If you take more *OxyContin* than you should or if someone accidentally swallows your tablets

Call your doctor or hospital straight away.

An overdose may result in:

- a reduction in size of pupils in the eye
- breathing more slowly or weakly than expected (respiratory depression)
- drowsiness or loss of consciousness
- low muscle tone (hypotonia)
- reduced pulse rate
- a fall in blood pressure
- difficulty in breathing due to fluid on the lungs (pulmonary oedema).

In severe cases an overdose may lead to unconsciousness or even death. When seeking medical attention, make sure that you take this leaflet and any remaining tablets with you to show to the doctor.

If you have taken too many tablets under no circumstances should you put yourself in a situation that requires you to be alert e.g. driving a car.

If you forget to take *OxyContin*

If you remember within 4 hours of the time your tablet was due, take your tablet straight away. Take your next tablet at your normal time. If you are more than 4 hours late, please call your doctor or pharmacist for advice. Do not take a double dose to make up for a forgotten tablet.

If you stop taking *OxyContin*

You should not suddenly stop taking these tablets unless your doctor tells you to. If you want to stop taking your tablets, discuss this with your doctor first. They will tell you how to do this, usually by reducing the dose gradually so you do not experience unpleasant effects. Withdrawal symptoms such as agitation, anxiety, palpitations, shaking or sweating may occur if you suddenly stop taking these tablets.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, these tablets can cause side effects, although not everybody gets them.

This medicine can cause allergic reactions, although serious allergic reactions are reported in rare cases. Tell your doctor immediately if you get any sudden wheeziness, difficulties in breathing, swelling of the eyelids, face or lips, rash or itching especially those covering your whole body.

The most serious side effect is a condition where you breathe more slowly or weakly than expected (respiratory depression – a typical hazard of an opioid overdose).

As with all strong analgesics or painkillers, there is a risk that you may become addicted or reliant on these tablets.

Very common: may affect more than 1 in 10 people

- Constipation (your doctor can prescribe a laxative to overcome this problem).
- Feeling or being sick (this should normally wear off after a few days, however your doctor can prescribe an anti-sickness medicine if it continues to be a problem).
- Drowsiness (this is most likely when you start taking your tablets or when your dose is increased, but it should wear off after a few days).
- Dizziness.
- Headache.
- Itchy skin.

Common: may affect up to 1 in 10 people

- Dry mouth, loss of appetite, indigestion, abdominal pain or discomfort, diarrhoea.
- Confusion, depression, a feeling of unusual weakness, shaking, lack of energy, tiredness, anxiety, nervousness, difficulty in sleeping, abnormal thoughts or dreams.
- Wheezing or difficulty in breathing, shortness of breath.
- Difficulty in passing urine.
- Rash.
- Sweating, high temperature.

Uncommon: may affect up to 1 in 100 people

- A condition where you breathe more slowly or weakly than expected (respiratory depression).
- Difficulty in swallowing, belching, hiccups, wind, a condition where the bowel does not work properly (ileus), inflammation of the stomach, gastrointestinal disorders (e.g. upset stomach), changes in taste, mouth ulcers, sore mouth.
- A condition which causes abnormal production of antidiuretic hormone (syndrome of inappropriate antidiuretic hormone secretion).
- A feeling of dizziness or 'spinning' (vertigo), hallucinations, mood swings, a feeling of extreme happiness, agitation, generally feeling unwell, loss of memory, difficulty in speaking, reduced sensitivity to pain or touch, tingling or numbness, seizures, fits or

convulsions, abnormal manner or style of walking, feeling detached from oneself, being unusually overactive, reduced consciousness, unusual muscle stiffness or slackness, involuntary muscle contractions.

- Impotence, decreased sexual drive, low levels of sex hormones in the blood ('hypogonadism', seen in a blood test).
- Flushing of the skin.
- Dehydration, weight change, thirst, swelling of the hands, ankles or feet.
- Dry skin.
- Tear disorder, blurred vision, reduction in size of the pupils in the eye.
- A need to take increasingly higher doses of the tablets to gain the same level of pain relief (tolerance).
- A ringing or buzzing sound in the ears.
- Swelling and irritation inside the nose, nose bleeds, voice alteration.
- Chills.
- Chest pain.
- Inability to fully empty the bladder
- A worsening in liver function tests (seen in a blood test).
- Withdrawal symptoms (see section 3 'If you stop taking *OxyContin*').

Rare: may affect up to 1 in 1,000 people

- A feeling of 'faintness' especially on standing up.
- Low blood pressure.
- Hives.

Not known: frequency cannot be estimated from the available data

- Sudden wheeziness, difficulties in breathing, swelling of the eyelids, face or lips, rash or itching especially those covering your whole body.
- Tooth decay.
- Colicky abdominal pain or discomfort.
- A blockage in the flow of bile from the liver. This can cause itchy skin, yellow skin, very dark urine and very pale stools.
- Absence of menstrual periods.
- An increase in sensitivity to pain.
- Aggression.
- Long term use of *OxyContin* during pregnancy may cause life-threatening withdrawal symptoms in the newborn. Symptoms to look for in the baby include irritability, hyperactivity and abnormal sleep pattern, high pitched cry, shaking, being sick, diarrhea and not putting on weight.

You may see the remains of the tablets in your faeces. This should not affect how the tablets work.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

5. How to store *OxyContin*

Keep this medicine out of the sight and reach of children. Accidental overdose by a child is dangerous and may be fatal.

Do not use this medicine after the expiry date which is stated on the blister and carton. The expiry date refers to the last day of that month. EXP 08 2020 means that you should not take the tablets after the last day of August 2020.

Do not store your tablets above 25°C.

Do not take your tablets if they are broken or crushed as this can be dangerous and can cause serious problems such as overdose.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What *OxyContin* contains

The active ingredient is oxycodone hydrochloride. Each tablet contains 15 mg, 30 mg, 60 mg, 120 mg or 160 mg of oxycodone hydrochloride.

The tablet cores contain:

- Lactose monohydrate
- Povidone
- Ammoniomethacrylate polymer
- Sorbic acid
- Triacetin
- Stearyl alcohol
- Talc
- Magnesium stearate

The tablet coatings contain:

- Hypromellose (E464)
- Titanium dioxide (E171)
- Macrogol

In addition, the tablet coatings contain the following:

15 mg – iron oxide (E172)

30 mg, 60 mg and 120 mg - polysorbate 80 (E433), and iron oxide (E172)

160 mg – polysorbate 80 (E433) and indigo carmine (E132)

What *OxyContin* looks like and the contents of the pack

OxyContin tablets are marked OC on one side and the strength on the other (15, 30 etc). All strengths, apart from the 160 mg tablet strength, are round, bi-convex tablets.

Approximate tablet diameters: 15, 30 and 60 mg – 8 mm; 120 mg - 10 mm. The 160 mg strength tablets are capsule shaped and are approximately 16 mm in length by 6 mm in diameter.

The tablets are all film coated in the following colours: 15 mg - grey, 30 mg - brown, 60 mg - red, 120 mg - purple and 160 mg - blue.

In each box there are 28, 56 or 112 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

The tablets are made by Bard Pharmaceuticals Limited at Cambridge Science Park, Milton Road, Cambridge CB4 0GW, UK.

The marketing authorisation holder is: Mundipharma Pharmaceuticals Limited Millbank House, Arkle Road, Sandyford, Dublin 18, Ireland.

This leaflet is also available in large print, Braille or as an audio CD. To request a copy, please call the RNIB Medicine Information Line on:

0044 1733 37 53 70

You will need to give details of the product name and reference number.

These are as follows:

Product name: OxyContin

Reference number: 1688/5/1

This leaflet was last revised in July 2017

® *OxyContin*, MUNDIPHARMA and the ‘mundipharma’ device (logo) are Registered Trade Marks.

© 2008-2014 Napp Pharmaceuticals Limited.