

Package leaflet: Information for the user

Amlodipine Masuu 5mg Tablets

Amlodipine Masuu 10mg Tablets

Amlodipine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Amlodipine Masuu is and what it is used for
2. What you need to know before you take Amlodipine Masuu
3. How to take Amlodipine Masuu
4. Possible side effects
5. How to store Amlodipine Masuu
6. Contents of the pack and other information

1. What Amlodipine Masuu is and what it is used for

Amlodipine Masuu contains the active substance amlodipine which belongs to a group of medicines called calcium antagonists.

Amlodipine Masuu is used to treat high blood pressure (hypertension) or a certain type of chest pain called angina, a rare form of which is Prinzmetal's or variant angina.

In patients with high blood pressure your medicine works by relaxing blood vessels, so that blood passes through them more easily. In patients with angina Amlodipine Masuu works by improving blood supply to the heart muscle which then receives more oxygen and as a result chest pain is prevented. Your medicine does not provide immediate relief of chest pain from angina.

2. What you need to know before you take Amlodipine Masuu

Do not take Amlodipine Masuu

- If you are allergic to the active substance, any other calcium antagonists or any of the other ingredients of this medicine (listed in section 6). This may be itching, reddening of the skin or difficulty in breathing.
- If you have severe low blood pressure (hypotension).
- If you have narrowing of the aortic heart valve (aortic stenosis) or cardiogenic shock (a condition where your heart is unable to supply enough blood to the body).
- If you suffer from heart failure after a heart attack.

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Amlodipine Masuu.

You should inform your doctor if you have or have had any of the following conditions:

- Recent heart attack
- Heart failure
- Severe increase in blood pressure (Hypertensive crisis)
- Liver disease
- You are elderly and your dose needs to be increased

Children and adolescents

Amlodipine Masuu has not been studied in children under the age of 6 years. Amlodipine Masuu should only be used for hypertension in children and adolescents from 6 years to 17 years of age (see section 3).

For more information, talk to your doctor.

Other medicines and Amlodipine Masuu

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Amlodipine Masuu may affect or be affected by other medicines, such as:

- ketoconazole, itraconazole (anti-fungal medicines)
- ritonavir, indinavir, nelfinavir (so called protease inhibitors used to treat HIV)
- rifampicin, erythromycin, clarithromycin (antibiotics)
- hypericum perforatum (St. John's Wort)
- verapamil, diltiazem (heart medicines)
- dantrolene (infusion for severe body temperature abnormalities)
- simvastatin (a cholesterol lowering medicine)

Amlodipine Masuu may lower your blood pressure even more if you are already taking other medicines to treat your high blood pressure.

Amlodipine Masuu with food and drink

Grapefruit juice and grapefruit should not be consumed by people who are taking Amlodipine Masuu. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active ingredient amlodipine, which can cause an unpredictable increase in the blood pressure lowering effect of Amlodipine Masuu.

Pregnancy and breast-feeding

Pregnancy

The safety of amlodipine in human pregnancy has not been established. If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Breast-feeding

It is not known whether amlodipine is passed into breast milk. If you are breast-feeding or about to start breast-feeding you must tell your doctor before taking Amlodipine Masuu.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Amlodipine Masuu may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

3. How to take Amlodipine Masuu

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 5 mg Amlodipine Masuu once daily. The dose can be increased to 10 mg Amlodipine Masuu once daily.

Your medicine can be used before or after food and drinks. You should take your medicine at the same time each day with a drink of water. Do not take Amlodipine Masuu with grapefruit juice.

Use in children and adolescents

For children and adolescents (6-17 years old), the recommended usual starting dose is 2.5 mg a day. The maximum recommended dose is 5 mg a day.

For the 2.5 mg dose (currently not available), the patient should take other medicinal products available on the market.

It is important to keep taking the tablets. Do not wait until your tablets are finished before seeing your doctor.

If you take more Amlodipine Masuu than you should

Taking too many tablets may cause your blood pressure to become low or even dangerously low. You may feel dizzy, lightheaded, faint or weak. If blood pressure drop is severe enough shock can occur. Your skin could feel cool and clammy and you could lose consciousness. Seek immediate medical attention if you take too many Amlodipine Masuu tablets.

If you forget to take Amlodipine Masuu

Do not worry. If you forget to take a tablet, leave out that dose completely. Take your next dose at the right time. Do not take a double dose to make up for a forgotten dose.

If you stop taking Amlodipine Masuu

Your doctor will advise you how long to take your medicine. Your condition may return if you stop using your medicine before you are advised.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Visit your doctor immediately if you experience any of the following very rare, severe side effects after taking this medicine.

- Sudden wheeziness, chest pain, shortness of breath or difficulty in breathing
- Swelling of eyelids, face or lips
- Swelling of the tongue and throat which causes great difficulty breathing
- Severe skin reactions including intense skin rash, hives, reddening of the skin over your whole body, severe itching, blistering, peeling and swelling of the skin, inflammation of mucous membranes (Stevens Johnson Syndrome) or other allergic reactions
- Heart attack, abnormal heart beat
- Inflamed pancreas which may cause severe abdominal and back pain accompanied with feeling very unwell

The following **common side-effects** have been reported. If any of these cause you problems or if they **last for more than one week**, you should **contact your doctor**.

Common: may affect up to 1 in 10 people

- Headache, dizziness, sleepiness (especially at the beginning of treatment)
- Palpitations (awareness of your heart beat), flushing
- Abdominal pain, feeling sick (nausea)
- Ankle swelling (oedema), tiredness

Other side-effects that have been reported include the following list. If any of these get serious, or if you notice any side-effects not listed in this leaflet, please tell your doctor or pharmacist.

Uncommon: may affect up to 1 in 100 people

- Mood changes, anxiety, depression, sleeplessness
- Trembling, taste abnormalities, fainting, weakness
- Numbness or tingling sensation in your limbs; loss of pain sensation
- Visual disturbances, double vision, ringing in the ears
- Low blood pressure
- Sneezing/running nose caused by inflammation of the lining of the nose (rhinitis)
- Altered bowel habits, diarrhoea, constipation, indigestion, dry mouth, vomiting (being sick)
- Hair loss, increased sweating, itchy skin, red patches on skin, skin discoloration
- Disorder in passing urine, increased need to urinate at night, increased number of times of passing urine
- Inability to obtain an erection, discomfort or enlargement of the breasts in men
- Weakness, pain, feeling unwell
- Joint or muscle pain, muscle cramps, back pain
- Weight increase or decrease

Rare: may affect up to 1 in 1,000 people

- Confusion

Very rare: may affect up to 1 in 10,000 people

- Decreased numbers of white blood cells, decrease in blood platelets which may result in unusual bruising or easy bleeding (red blood cell damage)
- Excess sugar in blood (hyperglycaemia)

- A disorder of the nerves which can cause weakness, tingling or numbness
- Cough, swelling of the gums
- Abdominal bloating (gastritis)
- Abnormal liver function, inflammation of the liver (hepatitis), yellowing of the skin (jaundice), liver enzyme increase which may have an effect on some medical tests
- Increased muscle tension
- Inflammation of blood vessels, often with skin rash
- Sensitivity to light
- Disorders combining rigidity, tremor, and/or movement disorders

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly to:

HPRA Pharmacovigilance
 Earlsfort Terrace
 IRL - Dublin 2
 Tel: +353 1 6764971
 Fax: +353 1 6762517
 Website: www.hpra.ie
 e-mail: medsafety@hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Amlodipine Masuu

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after 'EXP'. The expiry date refers to the last day of that month.

Store in the original package in order to protect from light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Amlodipine Masuu contains

- The active substance is amlodipine besilate. Each tablet contains amlodipine besilate equivalent to 5mg amlodipine.

- The active substance is amlodipine besilate. Each tablet contains amlodipine besilate equivalent to 10mg amlodipine.

- The other ingredients are Cellulose Microcrystalline, Sodium Starch Glycolate, Silica, Colloidal Anhydrous and Magnesium stearate.

What Amlodipine Masuu looks like and contents of the pack

5 mg Tablets: White to off-white, round biconvex, uncoated tablets plain on both sides.

10 mg Tablets: White to off white, round biconvex, uncoated tablets with a score line on one side and plain on the other side. The tablet can be divided into equal doses.

White opaque PVC/PVdC- Alu Blister

5mg:
20, 28, 30, 50, 60, 100 tablets

10mg:
14, 20, 28, 30, 50, 60, 100 tablets

Not all pack sizes may be marketed.

Marketing Authorisation Holder

PharmSol Europe Limited
The Victoria Centre Unit 2
Lower Ground Floor
Valletta Road
Mosta, MST 9012
Malta

Manufacturer

Fairmed Healthcare GmbH
Maria-Goeppert-Straße 3
23562 Lübeck
Germany

Fine Foods Pharmaceuticals N.M.T. S.p.A.
Via Berlino, 39
24040 Zingonia-Verdellino, BG, Italy

This medicinal product is authorised in the Member States of the EEA under the following names:

Portugal	Amlodipina Masuu
Austria	Amlodipin PharmSol 5mg, 10mg Tabletten
Germany	Amlodipin Fair-Med Healthcare 5mg, 10mg Tabletten
Spain	Amlodipino PharmSol 5mg, 10mg Comprimidos EFG
Ireland	Amlodipine Masuu 5mg, 10mg Tablets
Poland	Amlodipine PharmSol

This leaflet was last revised in.