

Package leaflet: Information for the use

## **Linezolid 2 mg/ml solution for infusion**

Linezolid

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet**

1. What Linezolid is and what it is used for
2. What you need to know before you use Linezolid
3. How to use Linezolid
4. Possible side effects
5. How to store Linezolid
6. Contents of the pack and other information

### **1. What Linezolid is and what it is used for**

Linezolid is an antibiotic of the oxazolidinones group that works by stopping the growth of certain bacteria (germs) that cause infections. It is used to treat pneumonia and some infections in the skin or under the skin. Your doctor will have decided if Linezolid is suitable to treat your infection.

### **2. What you need to know before you use Linezolid**

#### **You should not be treated with Linezolid:**

- if you are allergic to linezolid or any of the other ingredients of this medicine (listed in section 6).
- if you are taking or have taken within the last 2 weeks any medicines known as monoamine oxidase inhibitors (MAOIs: for example phenelzine, isocarboxazid, selegiline, moclobemide). These medications may be used to treat depression or Parkinson's disease;
- if you are breast-feeding. This is because Linezolid passes into breast milk and could affect the baby.

### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Linezolid.

Linezolid may not be suitable for you if you answer **yes** to any of the following questions. In this case tell your doctor as he/she will need to check your general health and your

blood pressure before and during your treatment or may decide that another treatment is better for you.

Ask your doctor if you are not sure whether these categories apply to you.

- Do you have high blood pressure, whether or not you are taking medicines for this?
- Have you been diagnosed with an overactive thyroid?
- Do you have a tumour of the adrenal glands (phaeochromocytoma) or carcinoid syndrome (caused by tumours of the hormone system with symptoms of diarrhoea, flushing of the skin, wheezing)?
- Do you suffer from manic depression, schizoaffective disorder, mental confusion or other mental problems?
- Do you have a history of hyponatraemia (low blood sodium levels) or do you take medicines that lower blood sodium levels e.g. certain diuretics (also called “water tablets”) such as hydrochlorothiazide?
- Do you take any opioids?

The use of certain medicines, including antidepressants and opioids, together with Linezolid can lead to serotonin syndrome, a potentially life-threatening condition (see section 2 “Other medicines and Linezolid” and section 4).

### **Take special care with Linezolid**

Tell your doctor before you are treated with this medicine if you

- are elderly
- bruise and bleed easily
- are anaemic (have low red blood cells)
- are prone to getting infections
- have a history of seizures
- have liver problems or kidney problems particularly if you are on dialysis
- have diarrhoea

Tell your doctor immediately if during treatment you suffer from

- problems with your vision such as blurred vision, changes in colour vision, difficulty in seeing detail or if your field of vision becomes restricted.
- loss of sensitivity in your arms or legs or a sensation of tingling or pricking in your arms or legs.
- you may develop diarrhoea while taking or after taking antibiotics, including Linezolid. If this becomes severe or persistent or you notice that your stool contains blood or mucus, you should stop taking Linezolid immediately and consult your doctor. In this situation, you should not take medicines that stop or slow bowel movement.
- recurrent nausea or vomiting, abdominal pain or rapid breathing

- unexplained muscle pain, tenderness, or weakness, and/or dark urine. These can be signs of a serious condition called rhabdomyolysis (muscle breakdown), which can lead to kidney damage.
- Feeling sick and unwell with muscle weakness, headache, confusion and memory impairment which may indicate hyponatremia (low blood sodium levels)

### **Other medicines and Linezolid**

There is a risk that Linezolid may sometimes interact with certain other medicines to cause side effects such as changes in blood pressure, temperature or heart rate.

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

**Tell your doctor if you are taking or have taken within the last 2 weeks** the following medicines as Linezolid **must not** be taken if you are already taking these medicines or have taken them recently. (See also Section 2 above ‘Do not take Linezolid.’)

- monoamine oxidase inhibitors (MAOIs for example phenelzine, isocarboxazid, selegiline, moclobemide). These may be used to treat depression or Parkinson’s disease

Also tell your doctor if you are taking the following medicines. Your doctor may still decide to give you Linezolid, but will need to check your general health and your blood pressure before and during your treatment. In other cases, your doctor may decide that another treatment is better for you.

- Decongestant cold or flu remedies containing pseudoephedrine or phenylpropanolamine.
- Some medicines used to treat asthma such as salbutamol, terbutaline, fenoterol.
- Certain antidepressants known as tricyclics or SSRIs (selective serotonin reuptake inhibitors). There are many of these, including amitriptyline, citalopram, clomipramine, dosulepin, doxepin, fluoxetine, fluvoxamine, imipramine, lofepramine, paroxetine, sertraline.
- Medicines used to treat migraine such as sumatriptan and zolmitriptan.
- Medicines used to treat sudden, severe allergic reactions such as adrenaline (epinephrine).
- Medicines which increase your blood pressure, such as noradrenaline (norepinephrine), dopamine and dobutamine
- Opioids e.g., pethidine - used to treat moderate to severe pain.
- Medicines used to treat anxiety disorders, such as buspirone.
- Medicines that stop blood clotting, such as warfarin.
- An antibiotic called rifampicin

### **Linezolid with food, drink and alcohol**

- You can take Linezolid either before, during or after a meal.
- Avoid eating large amounts of mature cheese, yeast extracts, or soya bean extracts e.g. soy sauce and drinking alcohol, especially draught beers and wine. This is because linezolid may react with a substance called tyramine which is naturally present in some foods. This interaction may cause an increase in your blood pressure.
- If you develop a throbbing headache after eating or drinking, tell your doctor or pharmacist immediately.

### **Pregnancy, breast-feeding and fertility**

The effect of Linezolid in pregnant women is not known. Therefore it should not be taken in pregnancy unless advised by your doctor. If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should not breast-feed when taking Linezolid because it passes into breast milk and could affect the baby.

### **Driving and using machines**

Linezolid may make you feel dizzy or experience problems with your vision. If this happens, do not drive or operate any machinery. Remember that if you are unwell your ability to drive or operate machinery may be affected.

### **Linezolid contains glucose**

Each 1 ml of Linezolid contains 45.7 mg glucose (13.7 g glucose in one bag/bottle).

**This should be taken into account in patients with diabetes mellitus.**

### **Linezolid contains sodium**

Each 1 ml of Linezolid contains 0.38 mg sodium (main component of cooking/table salt) (114 mg sodium in one bag/bottle). The sodium in one bag is equivalent to 5.7% of the recommended maximum daily dietary intake of sodium for an adult.

**To be taken into consideration by patients on a controlled sodium diet.**

## **3. How to use Linezolid**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

### **Adults**

This medicine will be given to you through a drip (by infusion into a vein) by a doctor or healthcare professional.

The recommended dose for adults (18 years and older) is 300 ml (600 mg linezolid) twice daily which is given directly into the blood stream (intravenously) by a drip over a period of 30 to 120 minutes.

If you are on kidney dialysis, you should be given Linezolid after your dialysis treatment.

A course of treatment usually lasts 10 to 14 days, but can last up to 28 days. The safety and effectiveness of this medicine have not been established for treatment periods longer than 28 days. Your doctor will decide how long you should be treated.

While you are taking Linezolid, your doctor should perform regular blood tests to monitor your blood count.

Your doctor should monitor your eyesight if you take Linezolid for more than 28 days.

#### **Use in children and adolescents**

Linezolid is not normally used to treat children and adolescents (under 18 years old).

#### **If you receive more Linezolid than you should**

If you are concerned that you may have been given too much Linezolid, tell your doctor or a nurse at once.

#### **If you miss a dose of Linezolid**

As you will be given this medicine under close supervision, it is very unlikely that you will miss a dose. If you think that you have missed a dose of treatment, tell a doctor or nurse at once. Do not take a double dose to make up for a forgotten dose.

### **4. Possible side effects**

Like all medicines Linezolid can cause side-effects although not everybody gets them.

**Tell your doctor, nurse or pharmacist immediately** if you notice any of these side effects during your treatment with Linezolid:

The serious side effects (with frequency in brackets) of Linezolid are:

- Severe skin disorder (uncommon), swelling, particularly around the face and neck (uncommon), wheezing and/or difficulty breathing (rare). This may be the sign of an allergic reaction and it may be necessary for you to stop taking Linezolid. Skin reactions such as a raised purple rash due to inflammation of the blood vessels (rare), red sore skin and flaking (dermatitis) (uncommon), rash (common), itching (common).
- Problems with your vision (uncommon) such as blurred vision (uncommon), changes in colour vision (not known), difficulty in seeing detail (not known), or if your field of vision becomes restricted (rare).
- Severe diarrhoea containing blood and/or mucus (antibiotic associated colitis including pseudomembranous colitis), which in rare circumstances may develop into complications that are life-threatening (uncommon).

- Recurrent nausea or vomiting, abdominal pain or rapid breathing (rare).
- Fits or seizures (uncommon) have been reported with Linezolid.
- Serotonin syndrome (not known): You should let your doctor know if you experience agitation, confusion, delirium, rigidity, tremor, incoordination, seizure, rapid heartbeat, severe breathing problems and diarrhoea (suggestive of serotonin syndrome) while also taking antidepressants known as SSRI's or opioids (see section 2).
- Unexplained bleeding or bruising, which may be due to changes in the numbers of certain cells in the blood which may affect blood clotting or lead to anaemia (common).
- Changes in numbers of certain cells in the blood which may affect your ability to fight infection (uncommon) some signs of infection include: any fever (common), sore throat (uncommon), mouth ulcers (uncommon) and tiredness (uncommon).
- Rhabdomyolysis (rare): Signs and symptoms include unexplained muscle pain, tenderness, or weakness, and/or dark urine. These can be signs of a serious condition called rhabdomyolysis (muscle breakdown), which can lead to kidney damage.
- Inflammation of the pancreas (uncommon).
- Convulsions (uncommon).
- Transient ischaemic attacks (temporary disturbance of blood flow to the brain causing short term symptoms such as loss of vision, leg and arm weakness, slurring of speech and loss of consciousness) (uncommon).
- "Ringing" in the ears (tinnitus) (uncommon).

Numbness, tingling or blurred vision have been reported by patients who have been given Linezolid for more than 28 days. If you experience difficulties with your vision you should consult your doctor as soon as possible.

#### **Other side effects include:**

##### **Common (may affect up to 1 in 10 people)**

- Fungal infections especially vaginal or oral "thrush"
- Headache
- Metallic taste in the mouth
- Diarrhoea, nausea or vomiting
- Changes in some blood test results including those measuring proteins, salts or enzymes which measure your kidney or liver function or blood sugar levels
- Difficulty in sleeping
- Increased blood pressure
- Anaemia (low red blood cell)
- Dizziness
- Localised or general abdominal pain
- Constipation
- Indigestion

- Localised pain
- Reduction in platelets

**Uncommon (may affect up to 1 in 100 people)**

- Inflammation of the vagina or genital area in women
- Sensations such as tingling or feeling numb
- Swollen, sore, or discoloured tongue
- Dry mouth
- Pain at and around the place where the infusion (drip) was given
- Inflammation of the veins (including where the infusion (drip) was given)
- A need to urinate more often
- Chills
- Feeling thirsty
- Increased sweating
- Hyponatraemia (low blood sodium levels)
- Kidney failure
- Abdominal bloating
- Injection site pain
- Increase in creatinine
- Stomach pain
- Changes in heart rate (e.g., increase rate)
- Decrease of the blood cell count
- Weakness and/or sensory changes

**Rare (may affect up to 1 in 1,000 people)**

- Superficial tooth discolouration, removable with professional dental cleaning (manual descaling)

**The following side effects have also been reported (Not known: frequency cannot be estimated from the available data)**

- Alopecia (hair loss)

**Reporting of side effects**

If you get any side effects, talk to your doctor pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects, you can help provide more information on the safety of this medicine.

**For the UK:**

Yellow Card Scheme.

Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App store.

**For Ireland:**

HPRA Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie)

**5. How to store Linezolid**

Keep this medicine out of the sight and reach of children.

Freeflex bag: Do not use this medicine after the expiry date which is stated on the carton, bags and overwrap after EXP. The expiry date refers to the last day of that month.

Hospital Staff will make sure that Linezolid is not used after the “Use by” date printed on the bag and that it is given to you as soon as the seal is broken. They will also visually inspect the solution prior to use and only clear solution, without particles will be used. They will also make sure that the solution is kept correctly in its box and foil wrapping in order to protect from light and out of the reach and sight of children until it is needed.

KabiPac bottle: Do not use this medicine after the expiry date which is stated on the label on the bottle and carton after EXP. The expiry date refers to the last day of the month.

Hospital Staff will make sure that Linezolid is not used after the “Use by” date printed on the bottle and that it is given to you as soon as removed from outer carton. They will also visually inspect the solution prior to use and only clear solution, without particles will be used. They will also make sure that the solution is kept correctly in its carton box in order to protect from light and out of the reach and sight of children until it is needed.

**After opening:**

Chemical and physical in-use stability has been demonstrated for 24 hours at 2-8°C and 25°C.

From a microbiological point of view, unless the method of opening precludes the risk of microbial contamination, the product should be used immediately. If not used immediately, in-use storage times and conditions are the responsibility of the user

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. Contents of the pack and other information****What Linezolid contains**

-The active substance is linezolid. Each 1ml of solution contains 2 mg linezolid.

-The other ingredients are glucose monohydrate (a type of sugar), sodium citrate, citric acid, hydrochloric acid or sodium hydroxide and water for injections.



### **What Linezolid looks like and contents of the pack**

Freeflex bag:

**Linezolid** is presented as a clear, practically free from particles, colourless to yellow solution in single infusion bags containing 300 ml (600mg linezolid) of solution.

The bags are supplied in boxes of 10, 30 or 50 bags.

KabiPac bottle:

**Linezolid** is presented as a clear, practically free from particles, colourless to yellow or slightly brownish solution in single infusion bottle containing 300 ml (600mg linezolid) of solution.

The bottles are supplied in boxes of 10, 30 or 50 bottles.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

#### **Marketing Authorisation Holder**

##### **For UK**

Fresenius Kabi Limited  
Cestrian Court, Eastgate Way,  
Manor Park, Runcorn, Cheshire  
WA7 1NT, UK

##### **For Ireland**

Fresenius Kabi Deutschland GmbH  
Else-Kroener-Strasse 1  
61352 Bad Homburg v.d.H  
Germany

#### **Manufacturer**

Fresenius Kabi Polska Sp. z o.o.  
ul. Sienkiewicza 25  
PL-99-300 Kutno  
Poland  
Or

Fresenius Kabi Norge AS  
Svinesundsveien 80  
NO-1788 Halden,  
Norway

**This medicinal product is authorised in the Member States of the European Economic Area and in the United Kingdom (Northern Ireland) under the following names:**

Austria	Linezolid Kabi 2 mg/ ml Infusionslösung
Belgium	Linezolid Fresenius Kabi 2 mg/ml, oplossing voor infusie
Bulgaria	Linezolid Kabi 2 mg/ml инфузионен разтвор
Czech republic	Linezolid Kabi 2 mg/ ml
Croatia	Linezolid Kabi 2 mg/ml otopina za infuziju
Denmark	Linezolid Fresenius Kabi
Estonia	Linezolid Fresenius Kabi
France	Linezolid Kabi 2 mg/ml, solution pour perfusion
Germany	Linezolid Kabi 2 mg/ ml Infusionslösung
Greece	Linezolid Kabi
Hungary	Linezolid Fresenius Kabi, 2 mg/ ml oldatos infúzió
Ireland	Linezolid 2 mg/ ml solution for infusion
Italy	Linezolid Kabi
Luxembourg	Linezolid Kabi 2 mg/ ml Infusionslösung
Netherlands	Linezolid Fresenius Kabi 2 mg/ml, oplossing voor infusie
Norway	Linezolid Fresenius Kabi
Poland	Linezolid Kabi
Portugal	Linezolid Kabi
Romania	Linezolid Kabi 2 mg/ml soluție perfuzabilă
Slovenia	Linezolid Kabi 2 mg/ml raztopina za infundiranje
Slovakia	Linezolid Kabi 2 mg/ ml
Spain	Linezolid Kabi 2 mg/ml solución para perfusión
United Kingdom	Linezolid 2 mg/ ml solution for infusion

**This leaflet was last revised in April 2025.**

-----  
The following information is intended for healthcare professionals only:

**Linezolid 2mg/ml solution for infusion**  
**Linezolid**

**IMPORTANT:** Refer to Summary of Product Characteristics before prescribing.

Linezolid is not active against infections caused by Gram negative pathogens. Specific therapy against Gram negative organisms must be initiated concomitantly if co-infection with a Gram negative pathogen is documented or suspected.

**Description Freeflex bags:**

Single use, ready-to-use, latex-free, multilayered polyolefine film infusion Freeflex bags sealed inside a foil laminate overwrap. The bag holds 300 ml solution and is packaged in a box. Each box contains 10, 30 or 50 infusion bags.

Linezolid infusion contains linezolid 2 mg/ml in an isotonic, clear, practically free from particles, colourless to yellow solution. Other ingredients are: glucose monohydrate, sodium citrate, citric acid, hydrochloric acid or sodium hydroxide, water for injections

**Description KabiPac bottles:**

Single use, ready-to-use, low-density polyethylene bottles (KabiPac) as primary packaging closed with a cap containing a rubber disc to allow insertion of the needle. The bottle holds 300 ml solution and is packaged in a single carton box to protect from light. Each collective box contains 10, 30 or 50 infusion bags.

Linezolid infusion contains linezolid 2 mg/ml in an isotonic, clear, practically free from particles, colourless to yellow or slightly brownish solution. Other ingredients are: glucose monohydrate, sodium citrate, citric acid, hydrochloric acid or sodium hydroxide, water for injections.

**Dosage and method of administration**

Linezolid should only be initiated in a hospital environment and after consultation with a relevant specialist such as a microbiologist or an infectious diseases specialist.

Patients who commence treatment on the parenteral formulation may be switched to either oral presentation when clinically indicated. In such circumstances, no dose adjustment is required as linezolid has an oral bioavailability of approximately 100 %.

The solution for infusion should be administered over a period of 30 to 120 minutes.

The recommended linezolid dosage should be administered intravenously IV twice daily.

**Recommended dosage and duration for adults:**

The duration of treatment is dependent on the pathogen, the site of infection and its severity, and on the patient's clinical response.

The following recommendations for duration of therapy reflect those used in the clinical trials. Shorter treatment regimens may be suitable for some types of infection but have not been evaluated in clinical trials.

The maximum treatment duration is 28 days. The safety and effectiveness of linezolid have not yet been established for treatment periods longer than 28 days.

No increase in the recommended dosage or duration of treatment is required for infections associated with concurrent bacteraemia. The dose recommendation for the solution for infusion are as follows:

Infections	Dosage and route for twice daily administration	Duration of treatment
Nosocomial pneumonia	600 mg twice daily	10-14 Consecutive Days
Community acquired pneumonia		
Complicated skin and soft tissue infections	600 mg twice daily	

**Paediatric population:**

The safety and efficacy of linezolid in children aged (< 18 years old) has not been established. Currently available data are described in section 4.8, 5.1, and 5.2 of the SmPC but no recommendation on a posology can be made.

**Elderly:** No dose adjustment is required.

**Renal impairment:** No dose adjustment is required.

Severe renal impairment (i.e. CLCR < 30 ml/min): No dose adjustment is required. Due to the unknown clinical significance of higher exposure (up to 10-fold) to the two primary metabolites of linezolid in patients with severe renal insufficiency, linezolid should be used with special caution in these patients and only when the anticipated benefit is considered to outweigh the theoretical risk.

As approximately 30 % of a linezolid dose is removed during 3 hours of haemodialysis, Linezolid 2ml/ml solution for infusion should be given after dialysis in patients receiving such treatment. The primary metabolites of linezolid are removed to some extent by haemodialysis, but the concentrations of these metabolites are still very considerably higher following dialysis than those observed in patients with normal renal function or mild to moderate renal insufficiency. Therefore, linezolid should be used with special caution in patients with severe renal insufficiency who are undergoing dialysis, and only when the anticipated benefit is considered to outweigh the theoretical risk.

To date, there is no experience of linezolid administration to patients undergoing continuous ambulatory peritoneal dialysis (CAPD) or alternative treatments for renal failure (other than haemodialysis).

**Hepatic impairment:** Patients with mild to moderate hepatic insufficiency (Child-Pugh class A or B): No dose adjustment is required.

Severe hepatic impairment (Child-Pugh class C): As linezolid is metabolised by a non-enzymatic process, impairment of hepatic function would not be expected to significantly alter its metabolism and, therefore, no dose adjustment is recommended. However, there are limited clinical data and it is recommended that linezolid should be used in such patients only when the anticipated benefit is considered to outweigh the theoretical risk (see sections 4.4 and 5.2).

### **Contraindications**

Patients hypersensitive to linezolid or any of the excipients.

Linezolid should not be used in patients taking any medicinal product which inhibits monoamine oxidases A or B (e.g. phenelzine, isocarboxazid, selegiline, moclobemide) or within two weeks of taking any such medicinal product.

Unless there are facilities available for close observation and monitoring of blood pressure, linezolid should not be administered to patients with the following underlying clinical conditions or on the following types of concomitant medications:

- Patients with uncontrolled hypertension, phaeochromocytoma, carcinoid, thyrotoxicosis, bipolar depression, schizoaffective disorder, acute confusional states.
- Patients taking any of the following medications: Serotonin re-uptake inhibitors, tricyclic antidepressants, serotonin 5-HT<sub>1</sub> receptor agonists (triptans), directly and indirectly acting sympathomimetic agents (including the adrenergic bronchodilators, pseudoephedrine and phenylpropanolamine), vasopressor agents (e.g. adrenaline / epinephrine, noradrenaline / norepinephrine), dopaminergic agents (e.g. dopamine, dobutamine), pethidine or buspirone.

Breast-feeding should be discontinued prior to and throughout administration (see section 4.6 of SmPC).

### **Special warnings and precautions for use**

#### Myelosuppression

Myelosuppression (including anaemia, leucopenia, pancytopenia and thrombocytopenia) has been reported in patients receiving linezolid. In cases where the outcome is known, when linezolid was discontinued, the affected haematologic parameters have risen toward pretreatment levels. The risk of these effects appears to be related to the duration of treatment. Elderly patients treated with linezolid may be at greater risk of experiencing blood dyscrasias than younger patients. Thrombocytopenia may occur more commonly in patients with severe renal insufficiency, whether or not on dialysis, and in patients with

moderate to severe hepatic impairment. Therefore, close monitoring of blood counts is recommended in patients who: have pre-existing anaemia, granulocytopenia or thrombocytopenia; are receiving concomitant medications that may decrease haemoglobin levels, depress blood counts or adversely affect platelet count or function; have severe renal insufficiency or moderate to severe hepatic impairment; receive more than 10-14 days of therapy. Linezolid should be administered to such patients only when close monitoring of haemoglobin levels, blood counts and platelet counts is possible.

If significant myelosuppression occurs during linezolid therapy, treatment should be stopped unless it is considered absolutely necessary to continue therapy, in which case intensive monitoring of blood counts and appropriate management strategies should be implemented.

In addition, it is recommended that complete blood counts (including haemoglobin levels, platelets, and total and differentiated leucocyte counts) should be monitored weekly in patients who receive linezolid regardless of baseline blood count.

In compassionate use studies, a higher incidence of serious anaemia was reported in patients receiving linezolid for more than the maximum recommended duration of 28 days. These patients more often required blood transfusion. Cases of anaemia requiring blood transfusion have also been reported post marketing, with more cases occurring in patients who received linezolid therapy for more than 28 days.

Cases of sideroblastic anaemia have been reported post-marketing. Where time of onset was known, most patients had received linezolid therapy for more than 28 days. Most patients fully or partially recovered following discontinuation of linezolid with or without treatment for their anaemia.

#### Mortality imbalance in a clinical trial in patients with catheter-related Gram positive bloodstream infections

Excess mortality was seen in patients treated with linezolid, relative to vancomycin/dicloxacillin/oxacillin, in an open-label study in seriously ill patients with intravascular catheter-related infections [78/363 (21.5%) vs 58/363 (16.0%)]. The main factor influencing the mortality rate was the Gram positive infection status at baseline. Mortality rates were similar in patients with infections caused purely by Gram positive organisms (odds ratio 0.96; 95% confidence interval: 0.58-1.59) but were significantly higher ( $p=0.0162$ ) in the linezolid arm in patients with any other pathogen or no pathogen at baseline (odds ratio 2.48; 95% confidence interval: 1.38-4.46). The greatest imbalance occurred during treatment and within 7 days following discontinuation of study drug. More patients in the linezolid arm acquired Gram negative pathogens during the study and died from infection caused by Gram negative pathogens and polymicrobial infections. Therefore, in complicated skin and soft tissue infections linezolid should only be used in patients with known or possible co-infection with Gram negative organisms if there are no alternative treatment options available. In these circumstances treatment against Gram negative organisms must be initiated concomitantly.

### Antibiotic-associated diarrhoea and colitis

Antibiotic-associated diarrhoea and antibiotic-associated colitis, including pseudomembranous colitis and *Clostridioides difficile*-associated diarrhoea, has been reported in association with the use of nearly all antibiotics including linezolid and may range in severity from mild diarrhoea to fatal colitis. Therefore, it is important to consider this diagnosis in patients who develop serious diarrhoea during or after the use of linezolid. If antibiotic-associated diarrhoea or antibiotic-associated colitis is suspected or confirmed, ongoing treatment with antibacterial agents, including linezolid, should be discontinued and adequate therapeutic measures should be initiated immediately. Drugs inhibiting peristalsis are contraindicated in this situation.

### Lactic acidosis

Lactic acidosis has been reported with the use of linezolid. Patients who develop signs and symptoms of metabolic acidosis including recurrent nausea or vomiting, abdominal pain, a low bicarbonate level, or hyperventilation while receiving linezolid should receive immediate medical attention. If lactic acidosis occurs, the benefits of continued use of linezolid should be weighed against the potential risks.

### Mitochondrial dysfunction

Linezolid inhibits mitochondrial protein synthesis. Adverse events, such as lactic acidosis, anaemia and neuropathy (optic and peripheral), may occur as a result of this inhibition; these events are more common when the drug is used longer than 28 days.

### Serotonin syndrome

Spontaneous reports of serotonin syndrome associated with the co-administration of linezolid and serotonergic agents, including antidepressants such as selective serotonin reuptake inhibitors (SSRIs) and opioids have been reported (see section 4.5 of SmPC). Co-administration of linezolid and serotonergic agents is therefore contraindicated (see section 4.3 of SmPC) except where administration of linezolid and concomitant serotonergic agents is essential. In those cases patients should be closely observed for signs and symptoms of serotonin syndrome such as cognitive dysfunction, hyperpyrexia, hyperreflexia and incoordination. If signs or symptoms occur physicians should consider discontinuing either one or both agents; if the concomitant serotonergic agent is withdrawn, discontinuation symptoms can occur.

### Rhabdomyolysis

Rhabdomyolysis has been reported with the use of linezolid. Linezolid should be used with caution in patients with pre-disposing factors for rhabdomyolysis. If signs or symptoms of rhabdomyolysis are observed, linezolid should be discontinued and appropriate therapy initiated.

### Hyponatraemia and SIADH

Hyponatraemia and/or Syndrome of Inappropriate Antidiuretic Hormone Secretion (SIADH) have been observed in some patients treated with linezolid. It is recommended that serum sodium levels are monitored regularly in patients at risk of hyponatraemia such as elderly patients or

patients taking medicines that may lower blood sodium levels (e.g. thiazide diuretics such as hydrochlorothiazide).

#### Peripheral and optic neuropathy

Peripheral neuropathy, as well as optic neuropathy and optic neuritis sometimes progressing to loss of vision, have been reported in patients treated with Linezolid 2mg/ml solution for infusion; these reports have primarily been in patients treated for longer than the maximum recommended duration of 28 days.

All patients should be advised to report symptoms of visual impairment, such as changes in visual acuity, changes in colour vision, blurred vision, or visual field defect. In such cases, prompt evaluation is recommended with referral to an ophthalmologist as necessary. If any patients are taking Linezolid for longer than the recommended 28 days, their visual function should be regularly monitored.

If peripheral or optic neuropathy occurs, the continued use of Linezolid should be weighed against the potential risks.

There may be an increased risk of neuropathies when linezolid is used in patients currently taking or who have recently taken antimycobacterial medications for the treatment of tuberculosis.

#### Convulsions

Convulsions have been reported to occur in patients when treated with Linezolid. In most of these cases, a history of seizures or risk factors for seizures was reported. Patients should be advised to inform their physician if they have a history of seizures.

#### Monoamine oxidase inhibitors

Linezolid is a reversible, non-selective inhibitor of monoamine oxidase (MAOI); however, at the doses used for antibacterial therapy, it does not exert an anti-depressive effect. There are very limited data from drug interaction studies and on the safety of linezolid when administered to patients with underlying conditions and/or on concomitant medications which might put them at risk from MAO inhibition. Therefore, linezolid is not recommended for use in these circumstances unless close observation and monitoring of the recipient is possible.

#### Use with tyramine-rich foods

Patients should be advised against consuming large amounts of tyramine rich foods.

#### Superinfection

The effects of linezolid therapy on normal flora have not been evaluated in clinical trials.

The use of antibiotics may occasionally result in an overgrowth of non-susceptible organisms. For example, approximately 3% of patients receiving the recommended linezolid doses experienced drug-related candidiasis during clinical trials. Should superinfection occur during therapy, appropriate measures should be taken.



### Special populations

Linezolid should be used with special caution in patients with severe renal insufficiency and only when the anticipated benefit is considered to outweigh the theoretical risk (see sections 4.2 and 5.2).

It is recommended that linezolid should be given to patients with severe hepatic insufficiency only when the perceived benefit outweighs the theoretical risk.

### Impairment of fertility

Linezolid reversibly decreased fertility and induced abnormal sperm morphology in adult male rats at exposure levels approximately equal to those expected in humans; possible effects of linezolid on the human male reproductive system are not known.

### Clinical trials

The safety and effectiveness of linezolid when administered for periods longer than 28 days have not been established.

Controlled clinical trials did not include patients with diabetic foot lesions, decubitus or ischaemic lesions, severe burns or gangrene. Therefore, experience in the use of linezolid in the treatment of these conditions is limited.

### Excipients

#### Glucose

Each ml of the solution contains 45.7 mg (i.e. 13.7 g/300 ml) glucose. This should be taken into account in patients with diabetes mellitus or other conditions associated with glucose intolerance.

#### Sodium

Each ml of solution also contains 0.38 mg (114 mg/300 ml) sodium, equivalent to 0.02% of the WHO maximum recommended daily intake (RDI) of 2 g sodium for an adult. The sodium content should be taken into account in patients on a controlled sodium diet.

Linezolid solution for infusion may be further prepared for administration with sodium-containing solutions (see sections 4.2, 6.2, and 6.6) and this should be considered in relation to the total sodium from all sources that will be administered to the patient.

## **Interactions**

### Monoamine oxidase inhibitors

Linezolid is a reversible, non-selective inhibitor of monoamine oxidase (MAOI). There are very limited data from drug interaction studies and on the safety of linezolid when administered to patients on concomitant medications that might put them at risk from MAO inhibition. Therefore, linezolid is not recommended for use in these circumstances unless close observation and monitoring of the recipient is possible.

#### Potential interactions producing elevation of blood pressure

In normotensive healthy volunteers, linezolid enhanced the increases in blood pressure caused by pseudoephedrine and phenylpropanolamine hydrochloride. Co-administration of linezolid with either pseudoephedrine or phenylpropanolamine resulted in mean increases in systolic blood pressure of the order of 30-40 mm Hg, compared with 11-15 mm Hg increases with linezolid alone, 14-18 mm Hg with either pseudoephedrine or phenylpropanolamine alone and 8-11 mm Hg with placebo. Similar studies in hypertensive subjects have not been conducted. It is recommended that doses of drugs with a vasopressive action, including dopaminergic agents, should be carefully titrated to achieve the desired response when coadministered with linezolid.

#### Potential serotonergic interactions

The potential drug-drug interaction with dextromethorphan was studied in healthy volunteers. Subjects were administered dextromethorphan (two 20 mg doses given 4 hours apart) with or without linezolid. No serotonin syndrome effects (confusion, delirium, restlessness, tremors, blushing, diaphoresis, hyperpyrexia) have been observed in normal subjects receiving linezolid and dextromethorphan.

Post marketing experience: there has been one report of a patient experiencing serotonin syndrome-like effects while taking linezolid and dextromethorphan which resolved on discontinuation of both medications.

During clinical use of linezolid with serotonergic agents, including antidepressants such as selective serotonin reuptake inhibitors (SSRIs) and opioids, cases of serotonin syndrome have been reported. Therefore, while coadministration is contraindicated (see section 4.3 of SmPC), management of patients for whom treatment with linezolid and serotonergic agents is essential, is described in special warnings and precautions for use.

#### Use with tyramine-rich foods

No significant pressor response was observed in subjects receiving both linezolid and less than 100 mg tyramine. This suggests that it is only necessary to avoid ingesting excessive amounts of food and beverages with a high tyramine content (e.g. mature cheese, yeast extracts, undistilled alcoholic beverages and fermented soya bean products such as soy sauce).

#### Drugs metabolised by cytochrome P450

Linezolid is not detectably metabolised by the cytochrome P450 (CYP) enzyme system and it does not inhibit any of the clinically significant human CYP isoforms (1A2, 2C9, 2C19, 2D6, 2E1, 3A4). Similarly, linezolid does not induce P450 isoenzymes in rats. Therefore, no CYP450-induced drug interactions are expected with linezolid.

#### Rifampicin

The effect of rifampicin on the pharmacokinetics of linezolid was studied in sixteen healthy adult male volunteers administered linezolid 600 mg twice daily for 2.5 days with and without rifampicin 600 mg once daily for 8 days. Rifampicin decreased the linezolid

C<sub>max</sub> and AUC by a mean 21% [90% CI, 15, 27] and a mean 32% [90% CI, 27, 37], respectively. The mechanism of this interaction and its clinical significance are unknown.

#### Warfarin

When warfarin was added to linezolid therapy at steady-state, there was a 10% reduction in mean maximum INR on co-administration with a 5% reduction in AUC INR. There are insufficient data from patients who have received warfarin and linezolid to assess the clinical significance, if any, of these findings.

### **Fertility, pregnancy and lactation**

#### Pregnancy

There are limited data from the use of linezolid in pregnant women. Studies in animals have shown reproductive toxicity. A potential risk for humans exists.

Linezolid should not be used during pregnancy unless clearly necessary i.e. only if the potential benefit outweighs the theoretical risk.

#### Breast-feeding

Animal data suggest that linezolid and its metabolites may pass into breast milk and, accordingly, breastfeeding should be discontinued prior to and throughout administration.

#### Fertility

In animal studies, linezolid caused a reduction in fertility.

### **Effects on ability to drive and use machines**

Patients should be warned about the potential for dizziness or symptoms of visual impairment whilst receiving Linezolid and should be advised not to drive or operate machinery if any of these symptoms occurs.

### **Undesirable effects**

The table below provides a listing of adverse drug reactions with frequency based on all-causality data from clinical studies that enrolled more than 6,000 adult patients who received the recommended linezolid doses for up to 28 days.

Those most commonly reported were diarrhoea (8.9%), nausea (6.9%), vomiting (4.3%) and headache (4.2%), .

The most commonly reported drug-related adverse events which led to discontinuation of treatment were headache, diarrhoea, nausea and vomiting. About 3 % of patients discontinued treatment because they experienced a drug-related adverse event.

Additional adverse reactions reported from post-marketing experience are included in the table with frequency category 'Not known', since the actual frequency cannot be estimated from the available data.

The following undesirable effects have been observed and reported during treatment with linezolid with the following frequencies: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to

<1/10); uncommon ( $\geq 1/1,000$  to <1/100); rare ( $\geq 1/10,000$  to <1/1,000); very rare (<1/10,000); Not known (cannot be estimated from the available data)

<b>System Organ Class</b>	<b>Common (<math>\geq 1/100</math> to &lt;1/10)</b>	<b>Uncommon (<math>\geq 1/1,000</math> to &lt;1/100)</b>	<b>Rare (<math>\geq 1/10,000</math> to &lt;1/1,000)</b>	<b>Frequency not known (cannot be estimated from available data)</b>
<b>Infections and infestations</b>	candidiasis, oral candidiasis, vaginal candidiasis, fungal infections	antibiotic associated colitis, including pseudomembranous colitis* vaginites		
<b>Blood and the lymphatic system disorders</b>	thrombocytopenia*, anaemia*†	pancytopenia*, leucopenia*, neutropenia, eosinophilia	sideroblastic anaemia*	myelosuppression*,
<b>Immune system disorders</b>			anaphylaxis	
<b>Metabolism and nutrition disorders</b>		hyponatraemia	lactic acidosis*	
<b>Psychiatric disorders</b>	insomnia			
<b>Nervous system disorders</b>	headache, taste perversion (metallic taste), dizziness	convulsions*, peripheral neuropathy * hypoesthesia, paraesthesia		serotonin syndrome**,
<b>Eye disorders</b>		optic neuropathy*, blurred vision*	changes in visual field defect*	, optic neuritis*, loss of vision*, changes in visual acuity*, changes in colour vision*
<b>Ear and labyrinth disorders</b>		tinnitus		
<b>Cardiac disorders</b>		arrhythmia (tachycardia)		
<b>Vascular disorders</b>	hypertension	transient ischaemic attacks, phlebitis, thrombophlebitis		
<b>Gastrointestinal disorders</b>	diarrhoea, nausea, vomiting, localised or general abdominal pain, constipation, dyspepsia	pancreatitis, gastritis, abdominal distention, dry mouth, glossitis, loose stools, stomatitis, tongue discolouration or disorder	superficial tooth discolouration	
<b>Hepato- biliary disorders</b>	abnormal liver function test; increased AST, ALT	increased total bilirubin		

	or alkaline phosphatase			
<b>Skin and subcutaneous tissue disorders</b>	pruritus, rash	angioedema, urticaria, dermatitis bullous, dermatitis, diaphoresis,	toxic epidermal necrolysis#, Stevens-Johnson syndrome#, hypersensitivity vasculitis	alopecia
<b>Musculoskeletal and connective tissue disorders</b>			rhabdomyolysis*	
<b>Renal and urinary disorders</b>	increased BUN	renal failure polyuria, increased creatinine		
<b>Reproductive system and breast disorders</b>		vulvovaginal disorder		
<b>General disorders and administration site conditions</b>	fever, localised pain	chills, fatigue, injection site pain, increased thirst,		
<b>Investigations</b>	<u>Chemistry</u> Increased LDH, creatine kinase, lipase, amylase or non fasting glucose. Decreased total protein, albumin, sodium or calcium. Increased or decreased potassium or bicarbonate.  <u>Haematology</u> Increased neutrophils or eosinophils. Decreased haemoglobin, haematocrit or red blood cell count. Increased or decreased platelet or white blood cell counts.	<u>Chemistry</u> Increased sodium or calcium. Decreased non fasting glucose. Increased or decreased chloride.  <u>Haematology</u> Increased reticulocyte count. Decreased neutrophils.		

\* See section Special warnings and precautions for use

\*\* See sections Contraindications and Interactions

# ADR frequency estimated using “The Rule of 3”

† See below

The following adverse reactions to linezolid were considered to be serious in rare cases: localised abdominal pain, transient ischaemic attacks and hypertension.

†In controlled clinical trials where linezolid was administered for up to 28 days, 2.0% of the patients reported anaemia. In a compassionate use program of patients with life-threatening infections and underlying co-morbidities, the percentage of patients who developed anaemia when receiving linezolid for  $\leq 28$  days was 2.5% (33/1326) as compared with 12.3% (53/430) when treated for  $>28$  days. The proportion of cases reporting drug-related serious anaemia and requiring blood transfusion was 9% (3/33) in patients treated for  $\leq 28$  days and 15% (8/53) in those treated for  $>28$  days.

#### Paediatric population

Safety data from clinical studies based on more than 500 paediatric patients (from birth to 17 years) do not indicate that the safety profile of linezolid for paediatric patients differs from that for adult patients.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via:

#### **For the UK:**

Yellow Card Scheme.

Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App store.

#### **For Ireland:**

HPRA Pharmacovigilance

Website: [www.hpra.ie](http://www.hpra.ie)

#### **Overdose**

No specific antidote is known.

No cases of overdose have been reported. However, the following information may prove useful:

Supportive care is advised together with maintenance of glomerular filtration. Approximately 30% of a linezolid dose is removed during 3 hours of haemodialysis, but no data are available for the removal of linezolid by peritoneal dialysis or haemoperfusion.

## **Instructions for use and handling**

For single use only.

For Freeflex bags:

Remove overwrap only when ready to use, then check for minute leaks by squeezing the bag firmly. If the bag leaks, do not use as sterility may be impaired. The solution should be visually inspected prior to use and only clear solutions, without particles should be used. Do not use these bags in series connections. Any unused solution must be discarded. Do not reconnect partially used bags.

For KabiPac Bottles:

Remove from outer carton only when ready to use, The solution should be visually inspected prior to use and only clear solutions, without particles should be used. Do not use these bottles in series connections. Any unused solution must be discarded. No special requirements for disposal. Any unused product or waste should be disposed of in accordance with local requirements.

Do not reconnect partially used bottles.

Linezolid 2mg/ml solution for infusion is compatible with the following solutions: glucose 50 mg/ml (5%), intravenous infusion sodium chloride 9 mg/ml (0,9%) intravenous infusion, Ringer-lactate solution for injection (Hartmann's solution for injection).

## **Incompatibilities**

Additives should not be introduced into this solution. If linezolid is to be given concomitantly with other drugs, each drug should be given separately in accordance with its own directions for use. Similarly, if the same intravenous line is to be used for sequential infusion of several drugs, the line should be flushed prior to and following linezolid administration with a compatible infusion solution.

Linezolid 2mg/ml solution for infusion is known to be physically incompatible with the following compounds: amphotericin B, chlorpromazine hydrochloride, diazepam, pentamidine isethionate, erythromycin lactobionate, phenytoin sodium and sulphamethoxazole / trimethoprim. Additionally, it is chemically incompatible with ceftriaxone sodium.

## **Shelf Life**

Chemical and physical in-use stability has been demonstrated for 24 hours at 2-8°C and 25°C. From a microbiological point of view, unless the method of opening precludes the risk of microbial contamination, the product should be used immediately. If not used immediately, in-use storage times and conditions are the responsibility of the user.

## **Special Precautions for Storage**

Freeflex bag: Store in the original package (overwrap and carton) until ready to use in order to protect from light.

KabiPac Bottle: Keep the bottle in the outer carton until ready to use in order to protect from light.