

## Summary of Product Characteristics

### 1 NAME OF THE MEDICINAL PRODUCT

Capozide 50mg/25mg Tablets

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 50mg captopril and 25mg hydrochlorothiazide.

Excipients: Contains lactose. Product imported from Spain contains Sunset Yellow (E110)

*For a full list of excipients, see section 6.1.*

### 3 PHARMACEUTICAL FORM

Tablet

*Product imported from the U.K.*

White, oval, biconvex tablets with a breakline on one side and 'SQUIBB 390' on the other.

*Product imported from Spain*

Peach, oval, biconvex tablets with a breakline on one side and '390' on the other.

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic Indications

Treatment of essential hypertension.

This fixed dose combination is indicated in patients whose blood pressure is not adequately controlled by captopril alone or hydrochlorothiazide alone.

#### 4.2 Posology and method of administration

Capozide can be administered in a single or two divided doses/day with or without food in patients whose blood pressure is not adequately controlled by captopril alone or hydrochlorothiazide alone. A maximum daily dose of 100mg captopril/30mg hydrochlorothiazide should not be exceeded. If satisfactory reduction of blood pressure has not been achieved, additional antihypertensive medication may be added (*see 4.5*).

*Adults:* The administration of the fixed combination of captopril and hydrochlorothiazide is usually recommended after dosage titration with the individual components. The usual maintenance dose is 50/25mg, once a day, in the morning. When clinically appropriate a direct change from monotherapy to the fixed combination may be considered.

The 25/25mg\* strength may be used once a day for patients whose blood pressure is not adequately controlled by hydrochlorothiazide 25mg monotherapy and before titration of the captopril component. The 50/25mg and 25/25mg strengths are intended to be used once daily, as two tablets would result in an inappropriately high dose of hydrochlorothiazide (50mg/day). The 50/15mg strength may be administered to start the fixed combination in patients whose blood pressure is not adequately controlled by 50mg captopril monotherapy, and/or when a lower dose of hydrochlorothiazide is preferred.

**Renal impairment:** Creatinine clearance between 30 and 80 ml/min: the initial dose is usually 25/12.5mg once a day, in the morning.

The combination captopril/hydrochlorothiazide is contraindicated in patients with severe renal impairment (creatinine

clearance <30ml/min).

**Special populations:** In salt/volume depleted patients, elderly patients and diabetic patients, the usual starting dose is 25/12.5mg once a day.

**Children:** The safety and efficacy of Capozide in children has not been established.

\* Not available in Ireland

### 4.3 Contraindications

- History of hypersensitivity to captopril, to any of the excipients or any other ACE inhibitor.
- History of hypersensitivity to hydrochlorothiazide or other sulphonamide-derived drugs.
- History of angioedema associated with previous treatment with previous ACE inhibitor therapy.
- Hereditary/idiopathic angioneurotic oedema.
- Severe renal impairment (creatinine clearance <30 ml/min)
- Severe hepatic impairment
- Second and third trimester of pregnancy (*see 4.6*).
- Lactation (*see 4.6*)

### 4.4 Special warnings and precautions for use

#### CAPTOPRIL

**Hypotension:** Rarely hypotension is observed in uncomplicated hypertensive patients. Symptomatic hypotension is more likely to occur in hypertensive patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea, vomiting or haemodialysis. Volume and/or sodium depletion should be corrected before the administration of an ACE inhibitor and a lower starting dose should be considered.

As with any antihypertensive agent, excessive blood pressure lowering in patients with ischaemic cardiovascular or cerebrovascular disease may increase the risk of myocardial infarction or stroke. If hypotension develops, the patient should be placed in a supine position. Volume repletion with intravenous normal saline may be required.

**Renovascular hypertension:** There is an increased risk of hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with ACE inhibitors. Loss of renal function may occur with only mild changes in serum creatinine. In these patients, therapy should be initiated under close medical supervision with low doses, careful titration and monitoring of renal function.

**Angioedema:** Angioedema of the extremities, face, lips, mucous membranes, tongue, glottis or larynx may occur in patients treated with ACE inhibitors, particularly during the first weeks of treatment. However, in rare cases, severe angioedema may develop after long-term treatment with ACE inhibitor. Treatment should be discontinued promptly. Angioedema involving the tongue, glottis or larynx may be fatal. Emergency therapy should be instituted. The patient should be hospitalised and observed for at least 12 to 24 hours and should not be discharged until complete resolution of symptoms has occurred.

**Cough:** Cough has been reported with the use of ACE inhibitors. Characteristically, the cough is non-productive, persistent and resolves after discontinuation of therapy.

**Hepatic failure:** Rarely, ACE inhibitors have been associated with a syndrome that starts with cholestatic jaundice and progresses to fulminant hepatic necrosis and (sometimes) death. The mechanism of this syndrome is not understood. Patients receiving ACE inhibitors who develop jaundice or marked elevations of hepatic enzymes should discontinue the ACE inhibitors and receive appropriate medical follow-up.

**Hyperkalaemia:** Elevations in serum potassium have been observed in some patients treated with ACE inhibitors, including captopril. Patients at risk for the development of hyperkalaemia include those with renal insufficiency, diabetes mellitus, or those using concomitant potassium-sparing diuretics, potassium supplements or potassium-containing salt substitutes; or those patients taking other drugs associated with increased in serum potassium (e.g. heparin). If concomitant use of the above mentioned agents is deemed appropriate, regular monitoring of serum potassium is recommended.

**Aortic and mitral valve stenosis/Obstructive hypertrophic cardiomyopathy/Cardiogenic shock:** ACE inhibitors should be used with caution in patients with left ventricular valvular and outflow tract obstruction and avoided in cases of cardiogenic shock and haemodynamically significant obstruction.

**Neutropenia/Agranulocytosis:** Neutropenia/agranulocytosis, thrombocytopenia and anaemia have been reported in patients receiving ACE inhibitors, including captopril. In patients with normal renal function and no other complicating factors, neutropenia occurs rarely.

Captopril should be used with extreme caution in patients with collagen vascular disease, immunosuppressant therapy, treatment with allopurinol or procainamide, or a combination of these complicating factors, especially if there is pre-existing impaired renal function. Some of these patients developed serious infections which in a few instances did not respond to intensive antibiotic therapy.

If captopril is used in such patients, it is advised that white blood cell count and differential counts should be performed prior to therapy, every 2 weeks during the first 3 months of captopril therapy, and periodically thereafter. During treatment all patients should be instructed to report any sign of infection (e.g. sore throat, fever) when a differential white blood cell count should be performed. Captopril and other concomitant medication (*see 4.5*) should be withdrawn if neutropenia (neutrophils less than  $1000/\text{mm}^3$ ) is detected or suspected

In most patients neutrophil counts rapidly return to normal upon discontinuing captopril.

**Proteinurea:** Proteinurea may occur particularly in patients with existing renal function impairment or on relatively high doses of ACE inhibitors.

Total urinary proteins greater than 1 g per day were seen in about 0.7% of patients receiving captopril. The majority of patients had evidence of prior renal disease or had received relatively high doses of captopril (in excess of 150mg/day), or both. Nephritic syndrome occurred in about one-fifth of proteinuric patients. In most cases, proteinuria subsided or cleared with six months whether or not captopril was continued. Parameters of renal function, such as BUN and creatinine, were seldom altered in the patients with proteinuria.

Patients with prior renal disease should have urinary protein estimations (dip-stick on first morning urine) prior to treatment, and periodically thereafter.

**Anaphylactoid reactions during desensitisation:** Sustained life-threatening anaphylactoid reactions have been rarely reported for patients undergoing desensitising treatment with hymenoptera venom while receiving another ACE inhibitor. In the same patients, these reactions were avoided when the ACE inhibitor was temporarily withheld, but they reappeared upon inadvertent rechallenge. Therefore, caution should be used in patients with ACE inhibitors undergoing such desensitisation procedures.

**Anaphylactoid reactions during high-flux dialysis/lipoprotein apheresis membrane exposure:** Anaphylactoid reactions have been reported in patients haemodialysed with high-flux dialysis membranes or undergoing low-density lipoprotein apheresis with dextran sulphate absorption. In these patients, consideration should be given to using a different type of dialysis membrane or a different class of medication.

**Surgery/Anaesthesia:** Hypotension may occur in patients undergoing major surgery or during treatment with anaesthetic agents that are known to lower blood pressure. If hypotension occurs, it may be corrected by volume expansion.

**Diabetic patients:** The glycaemia levels should be closely monitored in diabetic patients previously treated with oral antidiabetic drugs or insulin, namely during the first month of treatment with an ACE inhibitor.

As with other angiotensin converting enzyme inhibitors, Capozide is apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of higher prevalence of low-renin states in the black hypertensive population.

## HYDROCHLOROTHIAZIDE

**Renal impairment:** In patients with renal disease, thiazides may precipitate azotaemia. Cumulative effects of the drug may develop in patients with impaired renal function. If progressive renal impairment becomes evident, as indicated by a rising non-protein nitrogen, careful reappraisal of therapy is necessary, with consideration given to discontinuing diuretic therapy (*see 4.3*)

**Hepatic impairment:** Thiazides should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma (*see 4.3*)

**Metabolic and endocrine effects:** Thiazide therapy may impair glucose tolerance. In diabetic patients dosage adjustments of insulin or oral hypoglycaemic agents may be required. Latent diabetes mellitus may become manifest during thiazide therapy. Increases in cholesterol and triglyceride levels have been associated with thiazide diuretic therapy.

Hyperuricaemia may occur or frank gout may be precipitated in certain patients receiving thiazide therapy.

**Electrolyte imbalance:** As for any patient receiving diuretic therapy, periodic determination of serum electrolytes should be performed at appropriate intervals.

Thiazides, including hydrochlorothiazide, can cause fluid or electrolyte imbalance (hypokalaemia, hyponatraemia and hypochloraemic alkalosis). Warning signs of fluid or electrolyte imbalance are dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pain or cramps, muscular fatigue, hypotension, oliguria, tachycardia and gastrointestinal disturbances such as nausea or vomiting. Although hypokalaemia may develop with the use of thiazide diuretics, concurrent therapy with captopril may reduce diuretic-induced hypokalaemia.

The risk of hypokalaemia is greatest in patients with cirrhosis of the liver, inpatients experiencing brisk diuresis, in patients who are receiving inadequate oral intake of electrolytes and in patients receiving concomitant therapy with corticosteroids or ACTH (*see 4.5*).

Dilutional hyponatraemia may occur in oedematous patients in hot weather. Chloride deficit is generally mild and usually does not require treatment.

Thiazide may decrease urinary calcium excretion and cause an intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism. Marked hypercalcaemia may be evidence of hidden hyperparathyroidism. Thiazides should be discontinued before carrying out tests for parathyroid function. Thiazides have been shown to increase the urinary excretion of magnesium, which may result in hypomagnesaemia.

**Anti-doping test:** Hydrochlorothiazide contained in this medication could produce a positive analytical result in an anti-doping test.

**Other:** Sensitivity reactions may occur in patients with or without a history of allergy or bronchial asthma. The possibility of exacerbation or activation of systemic lupus erythematosus has been reported.

## CAPTOPRIL/HYDROCHLOROTHIAZIDE COMBINATION

**Pregnancy:** Capozide is not recommended during the first trimester of pregnancy (*see 4.6*). If treatment is discontinued due to pregnancy, the prescriber should decide whether treatment of hypertension should be continued.

**Risk of hypokalaemia:** The combination of an ACE inhibitor with a thiazide diuretic does not rule out the occurrence of hypokalaemia. Regular monitoring of kalaemia should be performed.

**Combination with lithium:** Capozide is not recommended in association with lithium due to the potentiation of lithium toxicity.

**Lactose:** Capozide contains lactose. Therefore, it should not be used in cases of congenital galactosaemia, glucose and galactose malabsorption or lactase deficiency syndromes (rare metabolic diseases).

## 4.5 Interaction with other medicinal products and other forms of interaction

### CAPTOPRIL

**Potassium sparing diuretics or potassium supplements:** ACE inhibitors attenuate diuretic induced potassium loss. Potassium sparing diuretics (e.g. spironolactone, triamterene or amiloride), potassium supplements, or potassium-containing salt substitutes may lead to significant increases in serum potassium. If concomitant use is indicated because of demonstrated hypokalaemia they should be used with caution and with frequent monitoring of serum potassium (*see 4.4*).

**Diuretics:** (thiazides or loop diuretics): Prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with captopril (*see 4.4*). The hypotensive effects can be reduced by discontinuation of the diuretic, by increasing volume or salt intake or by initiating therapy with a low dose of captopril. However, no clinically significant drug interactions have been found in specific studies with hydrochlorothiazide or furosemide.

**Other antihypertensive agents:** Captopril has been safely co-administered with other commonly used anti-hypertensive agents (e.g. beta-blockers and long-acting calcium channel blockers). Concomitant use of these agents may increase the hypotensive effects of captopril. Nitroglycerine and other nitrates, or other vasodilators, should be used with caution.

**Treatment of acute myocardial infarction:** Captopril may be used concomitantly with acetylsalicylic acid (at cardiological doses), thrombolytics, beta-blockers and/or nitrates in patients with myocardial infarction.

**Tricyclic antidepressants/Antipsychotics:** ACE inhibitors may enhance the hypotensive effects of certain tricyclic antidepressants and antipsychotics (*see 4.4*). Postural hypotension may occur.

**Allopurinol, procainamide, cytostatic or immuno-suppressant agents:** Concomitant administration with ACE inhibitors may lead to an increased risk of leucopenia, especially when the latter area used at higher than currently recommended doses.

**Sympathomimetics:** May reduce the antihypertensive effects of ACE inhibitors; patients should be carefully monitored.

**Antidiabetics:** Pharmacological studies have shown the ACE inhibitors, including captopril, can potentiate the blood glucose-reducing effects of insulin and oral antidiuretics, such as sulphonylurea, in diabetes. Should this very rare interaction occur, it may be necessary to reduce the dose of the antidiabetic during simultaneous treatment with ACE inhibitors.

### HYDROCHLOROTHIAZIDE

**Amphotericin B (parenteral), carbenoxolone, corticosteroids, corticotrophin (ACTH) or stimulant laxative:** hydrochlorothiazide may intensify electrolyte imbalance, particularly hypokalaemia.

**Calcium salts:** Increased serum calcium levels due to decreased excretion may occur when administered concurrently with thiazide diuretics.

**Cardiac glycosides:** Enhanced possibility of digitalis toxicity associated with thiazide induced hypokalaemia.

**Cholestyramine resin and colestipol:** May delay or decrease absorption of hydrochlorothiazide. Sulphonamide diuretics should be taken at least one hour before or four to six hours after these medications.

**Nondepolarising muscle relaxants (e.g. tubocurarine chloride):** effects of these agents may be potentiated by hydrochlorothiazide.

**Drugs associated with torsades de pointes:** Because of the risk of hypokalaemia, caution should be used when hydrochlorothiazide is coadministered with drugs associated with torsades de pointes, e.g. some antiarrhythmics, some antipsychotics and other drugs known to induce torsades de pointes.

## CAPTOPRIL/HYDROCHLOROTHIAZIDE COMBINATION

**Lithium:** Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with ACE inhibitors. Concomitant use of thiazide diuretics may increase the risk of lithium toxicity and enhance the already increased risk of lithium toxicity with ACE inhibitors. The combination of captopril and hydrochlorothiazide with lithium is therefore not recommended and careful monitoring of serum lithium levels should be performed if the combination proves necessary.

**Non-steroidal anti-inflammatory medicinal products:** It has been described that non-steroidal anti-inflammatory medicinal products (NSAIDs) and ACE inhibitors exert an additive effect on the increase in serum potassium, whereas renal function may decrease. These effects are, in principle, reversible. Rarely, acute renal failure may occur, particularly in patients with compromised renal function such as the elderly or dehydrated. Chronic administration of NSAIDs may reduce the antihypertensive effect of an ACE inhibitor. The administration of NSAIDs may reduce the diuretic, natriuretic and antihypertensive effects of thiazide diuretics.

## Clinical chemistry

Captopril may cause a false-positive urine test for acetone. Hydrochlorothiazide may cause diagnostic interference of the bentiromide test. Thiazides may decrease serum PBI (Protein Bound Iodine) levels without signs of thyroid disturbance.

## 4.6 Fertility, pregnancy and lactation

### Pregnancy:

Capozide is not recommended during the first trimester of pregnancy. When a pregnancy is planned or confirmed, an alternative treatment should be initiated as soon as possible. Controlled studies with ACE inhibitors have not been done in humans, but a limited number of cases of first trimester exposures has now shown malformations.

Capozide is contraindicated during the second and third trimesters of pregnancy. Prolonged captopril exposure during the second and third trimesters is known to induce toxicity in foetuses (decreased renal function oligohydramnios, skull ossification retardation) and in neonates (neonatal renal failure, hypotension, hyperkalaemia) (*see also* 5.3).

Hydrochlorothiazide, in cases of prolonged exposure during the third trimester of pregnancy, may cause a foetoplacental ischaemia and risk of growth retardation, moreover, rare cases of hypoglycaemia and thrombocytopenia in neonates have been reported in case of exposure near term.

Hydrochlorothiazide can reduce plasma volume as well as the uteroplacental blood flow.

Should exposure to Capozide have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.

### Lactation:

Capozide is contraindicated in the lactation period.

Both captopril and hydrochlorothiazide are excreted in human milk. Thiazides during breast feeding by lactating mothers

have been associated with a decrease or even suppression of lactation. Hypersensitivity to sulphonamide-derived drugs, hypokalaemia and nuclear icterus might occur.

Because of the potential for serious adverse effects in nursing infants from both drugs, a decision should be made whether to discontinue nursing or to discontinue therapy, taking into account the importance of this therapy to the mother.

## 4.7 Effects on ability to drive and use machines

As with other antihypertensives, the ability to drive and use machines may be reduced, e.g. at the start of the treatment or when the dose is modified, and also when used in combination with alcohol, but these effects depend on the individual's susceptibility.

## 4.8 Undesirable effects

### CAPTOPRIL

Undesirable effects reported for captopril and/or ACE inhibitor therapy include:

#### Blood and lymphatic disorders:

*Very rare:* Neutropenia/agranulocytosis (*see 4.4*), pancytopenia, particularly in patients with renal dysfunction (*see 4.4*), anaemia (including aplastic and haemolytic), thrombocytopenia, lymphadenopathy, eosinophilia, auto-immune diseases and/or positive ANA-titres.

#### Metabolism and lymphatic disorders:

*Rare:* Anorexia

*Very rare:* Hyperkalaemia hypoglycaemia (*see 4.4*)

#### Psychiatric disorders:

*Common:* Sleep disorders

*Very rare:* Confusion, depression

#### Nervous system disorders:

*Common:* Taste impairment, dizziness

*Rare:* Drowsiness, headache and paraesthesia

*Very rare:* Cerebrovascular incidents, including stroke, and syncope

#### Eye disorders:

*Very rare:* Blurred vision

#### Cardiac disorder:

*Uncommon* Tachycardia or tachyarrhythmia, angina pectoris, palpitations

*Very rare:* Cardiac arrest, cardiogenic shock

#### Vascular disorders:

*Uncommon* Hypotension (*see 4.4*), Raynaud syndrome, flush, pallor

#### Respiratory, thoracic and mediastinal disorders:

*Common:* Dry, irritating (non-productive) cough (*see 4.4*) and dyspnoea

*Very rare:* Bronchospasm, rhinitis, allergic alveolitis/eosinophilic pneumonia

#### Gastrointestinal disorders:

*Common:* Nausea, vomiting, gastric irritations, abdominal pain, diarrhoea, constipation, dry mouth

*Rare:* Stomatitis/apthous ulcerations

*Very rare:* Glossitis, peptic ulcer, pancreatitis

**Hepato-biliary disorders:**

*Very rare:* Impaired hepatic function and cholestasis (including jaundice), hepatitis including necrosis, elevated liver enzymes and bilirubin.

**Skin and subcutaneous tissue disorders:**

*Common:* Pruritus with or without a rash, rash and alopecia

*Uncommon:* Angioedema (*see 4.4*)

*Very rare:* Urticaria, Stevens Johnson syndrome, erythema multiforme, photosensitivity, erythroderma, pemphigoid reactions and exfoliative dermatitis.

**Musculoskeletal, connective tissue and bone disorders:**

*Very rare:* Myalgia, arthralgia

**Renal and urinary disorders:**

*Rare:* Renal function disorders, including renal failure, polyuria, oliguria, increased urine frequency

*Very rare:* Nephritic syndrome

**Reproductive system and breast disorders:**

*Very rare:* Impotence, gynaecomastia

**General disorders:**

*Uncommon:* Chest pain, fatigue, malaise

*Very rare:* Fever

**Investigations:**

*Very rare:* Proteinuria, eosinophilia, increase of serum potassium, decrease of serum sodium, elevation of BUN, serum creatinine and serum bilirubin, decreases in haemoglobin, haematocrit, leucocytes, thrombocytes, positive ANA-titre, elevated ESR.

**HYDROCHLOROTHIAZIDE****Infections and infestations:** Sialadenitis

**Blood and lymphatic system disorders:** Leucopenia, neutropenia/agranulocytosis, thrombocytopenia, aplastic anaemia, haemolytic anaemia, bone marrow depression.

**Metabolism and nutrition disorders:** Anorexia, hyperglycaemia, glycosuria, hyperuricaemia, electrolyte imbalance (including hyponatraemia and hypokalaemia), increases in cholesterol and triglycerides.

**Psychiatric disorders:** Restlessness, depression, sleep disturbances

**Nervous system disorders:** Loss of appetite, paraesthesia, light-headedness

**Eye disorders:** Xanthopsia, transient blurred vision

**Ear and labyrinth disorders:** Vertigo

**Cardiac disorders:** Postural hypotension, cardiac arrhythmias

**Vascular disease:** Necrotising angitis (vasculitis, cutaneous vasculitis)

**Respiratory, thoracic and mediastinal disorders:** Respiratory distress (including pneumonitis and pulmonary oedema).

**Gastrointestinal disorders:** Gastric irritation, diarrhoea, constipation, pancreatitis.

Hepato-biliary disorders: jaundice (intrahepatic cholestatic jaundice).



**Skin and subcutaneous tissue disorders:** Photosensitivity reactions, rash, cutaneous lupus erythematosus-like reactions, reactivation of cutaneous lupus erythematosus-like reactions, reactivation of cutaneous lupus erythematosus, urticaria, anaphylactic reactions, toxic epidermal necrolysis.

**Musculoskeletal, connective tissue and bone disorders:** Muscle spasm

**Renal and urinary disorders:** Fever, weakness

## 4.9 Overdose

Symptoms of overdosage are: increased diuresis, electrolyte imbalance, severe hypotension, depression of consciousness (including coma), convulsions, paresis, cardiac arrhythmias, bradycardia, renal failure.

Measures to prevent absorption (e.g. gastric lavage, administration of absorbing agents and sodium sulphate within 30 minutes of intake) and to hasten elimination should be applied if ingestion is recent. If hypotension occurs, the patient should be placed in the shock position and sodium chloride and volume supplementation should be given rapidly. Treatment with angiotensin-II can be considered. Bradycardia or extensive vagal reactions should be treated by administering atropine. The use of a pacemaker may be considered. Constant monitoring of water, electrolyte and acid base balance, and blood glucose is essential. In the event of hypokalaemia, potassium substitution is necessary.

Captopril may be removed from circulation by haemodialysis. The degree to which hydrochlorothiazide is removed by haemolysis has not been established.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

**Pharmacotherapeutic group:** ACE (Angiotensin-Converting-Enzyme) inhibitors, combinations, ATC code: C09BA01.

Capozide is a combination of an ACE inhibitor, captopril, and an antihypertensive diuretic, hydrochlorothiazide. The combination of these agents has an additive antihypertensive effect, reducing blood pressure to a greater degree than either alone.

Captopril is an angiotensin converting enzyme (ACE) inhibitor, i.e. it inhibits ACE, the enzyme involved in the conversion of angiotensin I to angiotensin II- a vasoconstrictor which also stimulates aldosterone secretion by the adrenal cortex.

Such inhibition leads to:

Reduced aldosterone secretion increased plasma rennin activity, since aldosterone no longer exerts negative feedback, a drop in total peripheral resistance (with a preferential effect on muscles and kidneys) which is not accompanied by water and sodium retention or reflex tachycardia during long-term treatment. Captopril also exerts its antihypertensive effect in subjects with low or normal rennin concentrations.

Captopril is effective at all stages of hypertension, i.e. mild, moderate or severe. A reduction in supine and standing systolic and diastolic blood pressures is observed.

After a single dose, the antihypertensive effect is evident fifteen minutes post-dose and reaches a maximum between 1h and 1.5 h after administration of the drug. Its duration of action is dose-dependent and varies from 6 to 12 hours.

Blood pressure becomes normalised (seated DBP < 90mmHg) in patients after two weeks to one month of treatment and the drug retains its effectiveness over the course of time. Patients are also classified as responders if seated DBP decreased by 10% or more from baseline-BP.

Rebound hypertension does not occur when treatment is discontinued.

The treatment of hypertension with captopril leads to an increase in arterial compliance, a rise in renal blood flow without any significant drop in the glomerular filtration rate and a decrease in left ventricular hypertrophy.

Hydrochlorothiazide is a thiazide diuretic which acts by inhibiting the reabsorption of sodium in the cortical diluting segment of renal tubules. It increases the excretion of sodium and chloride in urine and, to a lesser extent, the excretion of potassium and magnesium thereby increasing urinary output and exerting an antihypertensive effect.

The time to onset of diuretic activity is approximately 2 hours. Diuretic activity reaches a peak after 4 hours and is maintained for 6 to 12 hours. Above a certain dose, thiazide diuretics reach a plateau in terms of therapeutic effect whereas adverse reactions continue to multiply. When treatment is ineffective, increasing the dose beyond recommended doses serves no useful purpose and often gives rise to adverse reactions.

The concomitant administration of captopril and hydrochlorothiazide in clinical trials led to greater reductions in blood pressure than when either of the products was administered alone.

The administration of captopril inhibits the rennin angiotensin aldosterone system and tends to reduce hydrochlorothiazide-induced potassium loss.

Combination of an ACE inhibitor with a thiazide diuretic produces a synergistic effect and also lessens the risk of hypokalaemia provoked by the diuretic alone.

## 5.2 Pharmacokinetic properties

Captopril is quickly absorbed after oral administration and maximum serum concentrations are obtained around one hour after administration. Minimum mean absorption is approximately 75%. Peak plasma concentrations are reached within 60-90 minutes. The presence of food in the gastrointestinal tract reduces absorption by about 30-40%. Approximately 25-30% of the circulating drug is bound to plasma proteins. The apparent elimination half-life of unchanged captopril in blood is about 2 hours. Greater than 95% of the absorption dose is eliminated in the urine within 24 hours; 40-50% is unchanged drug and the remainder are inactive disulphide metabolites (captopril disulphide and captopril cysteine disulphide). Impaired renal function could result in drug accumulation. Studies in animals indicate that captopril does not cross the blood-brain barrier to any significant extent. Oral absorption of hydrochlorothiazide is relatively rapid. The mean plasma half-life in fasted individuals has been reported to be 5 to 15 hours. Hydrochlorothiazide is eliminated rapidly by the kidney and excreted unchanged (>95%) in the urine.

## 5.3 Preclinical safety data

Animal studies performed during organogenesis with captopril and/or hydrochlorothiazide have not shown any teratogenic effect but captopril has produced foetal toxicity in several species, including foetal mortality during late pregnancy, growth retardation and postnatal mortality in the rat. Preclinical data reveal no other specific hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicology, genotoxicity and carcinogenicity.

# 6 PHARMACEUTICAL PARTICULARS

## 6.1 List of excipients

Excipients present in the product imported from the UK:

Lactose  
Magnesium stearate  
Pregelatinised maize starch  
Microcrystalline cellulose  
Stearic acid

Excipients present in the product imported from Spain:

Lactose  
Magnesium stearate  
Pregelatinised maize starch  
Microcrystalline cellulose  
Stearic acid  
Sunset yellow colouring agent (E110)

## 6.2 Incompatibilities

Not applicable.

## 6.3 Shelf Life

The shelf life expiry date for this product shall be the date on the container and outer package of the product in the country of origin.

## 6.4 Special precautions for storage

Do not store above 30°C.  
Store in the original package.

## 6.5 Nature and contents of container

Blisters packs of 28 or 30 tablets contained in an outer cardboard carton.

## 6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

## 7 PARALLEL PRODUCT AUTHORISATION HOLDER

PCO Manufacturing  
Unit 10, Ashbourne Business Park  
Rath  
Ashbourne  
Co. Meath

## 8 PARALLEL PRODUCT AUTHORISATION NUMBER

PPA 0465/043/001

## 9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 25 September 1996

Date of last renewal: 25 September 2006

## 10 DATE OF REVISION OF THE TEXT

April 2011